

Last time we found

$$e \text{ values: } \lambda_{nm} = \left(\frac{n\pi}{L}\right)^2 + \left(\frac{m\pi}{H}\right)^2, \quad n, m = 1, 2, \dots$$

$$e \text{ functions: } \phi_{nm}(x, y) = \sin \frac{n\pi x}{L} \sin \frac{m\pi y}{H}, \quad n, m = 1, 2, \dots$$

Recall

$$h(t) = C, \cos(\sqrt{\lambda} ct) + C_2 \sin(\sqrt{\lambda} ct)$$

By principle of superposition,

$$u(x, y, t) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} A_{nm} \sin \frac{n\pi x}{L} \sin \frac{m\pi y}{H} \cos(\sqrt{\lambda_{nm}} ct) + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} B_{nm} \sin \frac{n\pi x}{L} \sin \frac{m\pi y}{H} \sin(\sqrt{\lambda_{nm}} ct)$$

Next we use I.C.s and orthogonality to find A_{nm} and B_{nm} .

$$\alpha(x, y) = \alpha(x, y, 0) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} A_{nm} \sin \frac{n\pi x}{L} \sin \frac{m\pi y}{H}$$

Orthogonality I:

$$\int_0^H \int_0^L \alpha(x, y) \cdot \sin \frac{m\pi y}{H} \frac{dy}{H} = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} A_{nm} \sin \frac{n\pi x}{L} \int_0^H \frac{\sin \frac{m\pi y}{H} \sin \frac{m\pi y}{H}}{H} dy$$

$$= \begin{cases} 0, & m \neq \tilde{m} \\ \frac{H}{2}, & m = \tilde{m} \end{cases}$$

Change $\tilde{m} \rightarrow m$:

$$\frac{2}{H} \int_0^H \alpha(x, y) \sin \frac{m\pi y}{H} \frac{dy}{H} = \sum_{n=1}^{\infty} A_{nm} \sin \frac{n\pi x}{L} \int_0^H \frac{\sin \frac{m\pi y}{H} \sin \frac{m\pi y}{H}}{H} dy$$

Orthogonality II:

$$\frac{2}{H} \int_0^L \int_0^H \alpha(x,y) \sin \frac{m\pi y}{H} \cdot \sin \frac{n\pi x}{L} dy dx = \sum_{n=1}^{\infty} A_{nm} \underbrace{\int_0^L \sin \frac{n\pi x}{L} \sin \frac{n\pi x}{L} dx}_0$$

$$\tilde{n} \rightarrow n \quad \left\{ \begin{array}{l} 0, \quad n \neq \tilde{n} \\ \frac{L}{2}, \quad n = \tilde{n} \end{array} \right.$$

$$\therefore A_{nm} = \frac{4}{LH} \int_{x=0}^L \int_{y=0}^H \alpha(x,y) \sin \frac{n\pi x}{L} \sin \frac{m\pi y}{H} dy dx, \quad n, m = 1, 2, \dots$$

$$f(x,y) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} B_{nm} \underbrace{\sin \frac{n\pi x}{L} \cdot \sin \frac{m\pi y}{H}}_{\text{like } A_{nm}}$$

Hence,

$$B_{nm} \cdot \sqrt{\lambda_{nm}} \cdot c = \frac{4}{LH} \int_{x=0}^L \int_{y=0}^H \beta(x,y) \sin \frac{n\pi x}{L} \sin \frac{m\pi y}{H} dy dx$$

or


$$B_{nm} = \frac{1}{c\sqrt{\lambda_{nm}}} \cdot \frac{4}{LH} \int_0^L \int_0^H \beta(x,y) \sin \frac{n\pi x}{L} \sin \frac{m\pi y}{H} dy dx$$

where $\lambda_{nm} = \left(\frac{n\pi}{L}\right)^2 + \left(\frac{m\pi}{H}\right)^2$, $n, m = 1, 2, \dots$

Ex Consider the case when $L=1, H=1$. Set $c=1$.
Initially flat membrane $\Rightarrow \alpha(x,y)=0 \Rightarrow A_{nm}=0$

Initial "kick":

$$f(x, y) = \begin{cases} 1, & \frac{1}{4} \leq x, y \leq \frac{3}{4} \\ 0, & \text{otherwise} \end{cases}$$



$$B_{nm} = \frac{4}{\sqrt{2nm}} \int_{x=1/4}^{3/4} \int_{y=1/4}^{3/4} \sin \frac{n\pi x}{1} \sin \frac{m\pi y}{1} dy dx =$$


$$= \frac{4}{\sqrt{2nm}} \int_{x=1/4}^{3/4} \sin n\pi x dx \cdot \int_{y=1/4}^{3/4} \sin m\pi y dy =$$

$$= \frac{4}{\sqrt{2nm}} \cdot \frac{1}{nm\pi^2} \left(\cos \frac{3n\pi}{4} - \cos \frac{n\pi}{4} \right) \left(\cos \frac{3m\pi}{4} - \cos \frac{m\pi}{4} \right)$$

$$A_{nm} = (n\pi)^2 + (m\pi)^2$$

$$u(x, y, t) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} B_{nm} \sin n\pi x \sin m\pi y \cdot \sin(\sqrt{A_{nm}} t)$$

$$u(x, y, t) \approx \sum_{n=1}^N \sum_{m=1}^M B_{nm} \sin n\pi x \sin m\pi y \cdot \sin(\sqrt{A_{nm}} t)$$



m_x points in x-direction

m_y points in y-direction

Let's compute solution at

$m_t = 12$ instances of time


Let $N = M = 50$, $m_x = m_y = 40$, $m_t = 12$

2D $50^2 \cdot 40^2 \cdot 12 = 48$ million


1D $50 \cdot 40 \cdot 12 = 24,000$

See these graphs and a Matlab code on the course webpage


$u(x,y,t)$ at time $t = 0.01$




$u(x,y,t)$ at time $t = 0.096522$




$u(x,y,t)$ at time $t = 0.18304$




$u(x,y,t)$ at time $t = 0.26957$




$u(x,y,t)$ at time $t = 0.35609$




$u(x,y,t)$ at time $t = 0.44261$




$u(x,y,t)$ at time $t = 0.52913$




$u(x,y,t)$ at time $t = 0.61565$




$u(x,y,t)$ at time $t = 1.7404$




$u(x,y,t)$ at time $t = 1.827$




$u(x,y,t)$ at time $t = 1.9135$




$u(x,y,t)$ at time $t = 2$




$u(x,y,t)$ at time $t = 0.70217$




$u(x,y,t)$ at time $t = 0.7887$




$u(x,y,t)$ at time $t = 0.87522$




$u(x,y,t)$ at time $t = 0.96174$




$u(x,y,t)$ at time $t = 1.1348$




$u(x,y,t)$ at time $t = 1.3078$




$u(x,y,t)$ at time $t = 1.0483$




$u(x,y,t)$ at time $t = 1.2213$




$u(x,y,t)$ at time $t = 1.3943$




$u(x,y,t)$ at time $t = 1.4809$



$u(x,y,t)$ at time $t = 1.5674$



$u(x,y,t)$ at time $t = 1.6539$



```

clear;
mp = 50; % number of terms for each infinite series
x = linspace(0,1,40); % x-points in plot in x = 0..1
y = linspace(0,1,40); % y-points in plot in x = 0..1
t = linspace(1/12,2,24); % time points
% First compute e-values and coefficients
for n=1:mp
    lambda(n,m) = (n*pi)^2 + (m*pi)^2;
    B(n,m) = (4/(pi*sqrt(lambda(n,m))))*(1/n)*(cos(3*pi*n/4) - cos(pi*n/4))..
        *(1/m)*(cos(3*pi*m/4) - cos(pi*m/4));
end
% Compute full solution at all points in space and time
for k=1:length(t)
    for l=1:length(x)
        for j=1:length(y)
            u(l,j,k) = 0;
            for n=1:mp
                for m=1:mp
                    u(l,j,k) = u(l,j,k) + B(n,m)*sin(n*pi*x(l))*sin(m*pi*y(j))*..
                        sin(sqrt(lambda(n,m))*t(k));
                end
            end
        end
    end
end
% Create the plots
mcount = 0;
for k=1:6
    figure(k)
    clf
    for m=1:4
        mcount = mcount + 1;
        subplot(2,2,m)
        mesh(x,y,u(:,:,mcount));
        colormap('jet');
        set(gca,'fontsize',10);
        title(['u(x,y,t) at time t = ',num2str(t(mcount))]);
        xlabel('x');
        ylabel('y');
        set(gca,'fontsize',10);
        axis([0 1 0 1 -0.75 0.75]);
    end
end
% Print the plots to JPG files
figure(1); print-djpeg95 memb1.jpg
figure(2); print-djpeg95 memb2.jpg

```

figure(3): print -djpeg95 memb3.jpg
figure(4): print -djpeg95 memb4.jpg
figure(5): print -djpeg95 memb5.jpg
figure(6): print -djpeg95 memb6.jpg

Multidimensional Eigenvalue Problems

After applying separation of variables of variables to heat or wave equations, we obtained the following multidimensional eigenvalue problem:

$$\nabla^2 \phi + \lambda \phi = 0 \quad (x, y) \in \Omega \quad (1)$$

$$\beta_1 \phi + \beta_2 \nabla \phi \cdot \hat{n} = 0 \quad (x, y) \in \partial\Omega$$

Eqⁿ (1) is called Helmholtz equation. It is a partial case of regular Sturm-Liouville problem:

$$\nabla(p \nabla \phi) + q \phi + \lambda \sigma \phi = 0$$

Helmholtz eqⁿ: $p=1, q=0, \sigma=1.$

Thm (Helmholtz equation)

1. All e 'values are real.
2. There is an infinite sequence of e 'values. There is the smallest e 'value and no larger e 'value.
3. Corresponding to an e 'value, there may be more than one e 'function (recall, in 1D, for a regular S.-d. problem, every e 'value had only one e 'function).