

## Splines

Let  $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$ . A spline of degree  $m$  is a function  $S(x)$  which satisfies the following conditions:

1. For  $x \in [x_i, x_{i+1}]$ ,  $S(x) = S_i(x)$ : polynomial of degree  $\leq m$
2.  $S^{(m-1)}$  exists and continuous at the interior points  $x_1, \dots, x_{n-1}$ , i.e.  
$$\lim_{x \rightarrow x_i^-} S_{i-1}^{(m-1)}(x) = \lim_{x \rightarrow x_i^+} S_i^{(m-1)}(x)$$

EXAMPLE

$$S(x) = \begin{cases} 0, & -1 \leq x \leq 0 \\ x^2, & 0 \leq x \leq 1 \end{cases}$$

$$x_0 = -1, \quad x_1 = 0, \quad x_2 = 1$$

$S(x)$  is a spline of degree  $m = 2$  (quadratic spline).

EXAMPLE

$$S(x) = \begin{cases} 0, & -1 \leq x \leq 0 \\ 1 - (x - 1)^2, & 0 \leq x \leq 1 \end{cases}$$

$S(x)$  is not a quadratic spline because  $\lim_{x \rightarrow 0^-} S'(x) \neq \lim_{x \rightarrow 0^+} S'(x)$ .

## Cubic Spline Interpolation

Given  $f, x_0, \dots, x_n$  as above, find a cubic spline  $S$  such that  $S(x_i) = f(x_i)$ ,  $i = 0, \dots, n$ .

$n + 1$  points  $\Rightarrow n$  intervals  $\Rightarrow 4n$  coefficients

$2n$  conditions needed to interpolate  $f(x_0), \dots, f(x_n)$

$2(n - 1)$  conditions needed to ensure  $S', S''$  are continuous at the interior points

This leaves 2 free conditions. Popular choices are:

- (i)  $S''(x_0) = S''(x_n) = 0$ : the natural cubic spline interpolant
- (ii)  $S'(x_0) = f'(x_0), S'(x_n) = f'(x_n)$ : the clamped cubic spline interpolant

Given  $f(x)$ ;  $x_0, \dots, x_n$ , how to find  $S(x)$ ?

EXAMPLE

$$x_i = ih, \quad h = 1/n, \quad x_0 = 0, \quad x_n = 1$$

STEP 1 (2nd derivative conditions)

Since  $S_i(x)$  is a cubic polynomial,  $S_i''(x)$  is linear for  $x$

$$S_i''(x) = a_i \left( \frac{x_{i+1} - x}{h} \right) + a_{i+1} \left( \frac{x - x_i}{h} \right), \quad i = 0, \dots, n-1$$

(using Lagrange 1st order interpolating polynomial).

Then

$$\left. \begin{array}{l} S_i''(x_i) = a_i \\ S_i''(x_{i+1}) = a_{i+1} \end{array} \right\} \Rightarrow S_{i-1}''(x_i) = a_i = S_i''(x_i)$$

Thus,  $S''(x)$  is continuous at the interior nodes.

STEP 2 (function values)

Integrate twice

$$S_i(x) = \frac{a_i(x_{i+1} - x)^3}{6h} + \frac{a_{i+1}(x - x_i)^3}{6h} + b_i(x_{i+1} - x) + c_i(x - x_i)$$

the last two terms are written in this special form just for convenience

$$\left. \begin{array}{l} S_i(x_i) = a_i \frac{h^2}{6} + b_i h = f_i \\ S_i(x_{i+1}) = a_{i+1} \frac{h^2}{6} + c_i h = f_{i+1} \end{array} \right\} \begin{array}{l} b_i h = f_i - a_i \frac{h^2}{6} \\ c_i h = f_{i+1} - a_{i+1} \frac{h^2}{6} \end{array}$$

Substitute

$$\begin{aligned} S_i(x) &= \frac{a_i(x_{i+1} - x)^3}{6h} + \frac{a_{i+1}(x - x_i)^3}{6h} \\ &+ \left( f_i - a_i \frac{h^2}{6} \right) \left( \frac{x_{i+1} - x}{h} \right) + \left( f_{i+1} - a_{i+1} \frac{h^2}{6} \right) \left( \frac{x - x_i}{h} \right) \end{aligned}$$

STEP 3 (1st derivative conditions).

$$S_i'(x) = -\frac{a_i(x_{i+1} - x)^2}{2h} + \frac{a_{i+1}(x - x_i)^2}{2h} - \left( \frac{f_i}{h} - a_i \frac{h}{6} \right) + \left( \frac{f_{i+1}}{h} - a_{i+1} \frac{h}{6} \right) \quad (1)$$

$$S'_i(x_i) = -a_i \frac{h}{2} - \left( \frac{f_i}{h} - a_i \frac{h}{6} \right) + \left( \frac{f_{i+1}}{h} - a_{i+1} \frac{h}{6} \right)$$

$$S'_i(x_{i+1}) = a_{i+1} \frac{h}{2} - \left( \frac{f_i}{h} - a_i \frac{h}{6} \right) + \left( \frac{f_{i+1}}{h} - a_{i+1} \frac{h}{6} \right)$$

We require,  $S'_{i-1}(x_i) = S'_i(x_i)$ , so (shifting  $i \rightarrow i - 1$  in  $S'_i(x_{i+1})$ )

$$\begin{aligned} a_{i-1} \frac{h}{2} - \left( \frac{f_{i-1}}{h} - a_{i-1} \frac{h}{6} \right) + \left( \frac{f_i}{h} - a_i \frac{h}{6} \right) \\ = -a_i \frac{h}{2} - \left( \frac{f_i}{h} - a_i \frac{h}{6} \right) + \left( \frac{f_{i+1}}{h} - a_{i+1} \frac{h}{6} \right) \end{aligned}$$

Collect like terms

$$a_{i-1} \frac{h}{6} + a_i \left( \frac{h}{2} - \frac{h}{6} + \frac{h}{2} - \frac{h}{6} \right) + a_{i+1} \frac{h}{6} = (f_{i-1} - 2f_i + f_{i+1}) / h$$

Multiply both sides by  $\frac{6}{h}$

$$a_{i-1} + 4a_i + a_{i+1} = \frac{6}{h^2} (f_{i-1} - 2f_i + f_{i+1}) \quad (*)$$

This holds for  $i = 1, \dots, n - 1$ .

**STEP 4** (boundary conditions)

$$\begin{aligned} S''_0(x_0) = 0 &\Rightarrow a_0 = 0 \\ S''_{n-1}(x_n) = 0 &\Rightarrow a_n = 0 \end{aligned} \quad (*)$$

$$\begin{pmatrix} 4 & 1 & 0 & 0 & \dots & 0 \\ 1 & 4 & 1 & 0 & \dots & 0 \\ & \ddots & \ddots & \ddots & & \\ & & \ddots & \ddots & \ddots & \\ & & & \ddots & \ddots & 1 \\ 0 & & & & 1 & 4 \end{pmatrix} \begin{pmatrix} a_1 \\ \vdots \\ \vdots \\ a_{n-1} \end{pmatrix} = \frac{6}{h^2} \begin{pmatrix} f_0 - 2f_1 + f_2 \\ \vdots \\ \vdots \\ f_{n-2} - 2f_{n-1} + f_n \end{pmatrix} \quad (3)$$

The coefficient matrix is

- tridiagonal
- symmetric

- positive definite
- strictly diagonally dominant

Notes:

1. For clamped splines, there are two additional equations that involve  $a_0$  and  $a_n$ .
2. Condition (\*) explains why the spline has additional inflection points. See the handout about natural cubic spline interpolation.

Recall

$A = (a_{ij})$  is strictly diagonally dominant if

$$|a_{ii}| > \sum_{\substack{j=1 \\ j \neq i}}^n |a_{ij}| \quad \text{for all } i = 1, \dots, n$$

**Theorem** *If  $A$  is strictly diagonally dominant, then  $A$  is invertible.*

**Proof.** Suppose  $A$  is not invertible. Then there exists a vector  $x \neq 0$  such that  $Ax = 0$ . Choose index  $i$  such that  $|x_i| = \|x\|_\infty = \max_j |x_j|$ . Then

$$0 = (Ax)_i = \sum_{j=1}^n a_{ij}x_j = a_{ii}x_i + \sum_{\substack{j=1 \\ j \neq i}}^n a_{ij}x_j$$

$$\Rightarrow |a_{ii}x_i| = \left| - \sum_{\substack{j=1 \\ j \neq i}}^n a_{ij}x_j \right| \leq \sum_{\substack{j=1 \\ j \neq i}}^n |a_{ij}||x_j|$$

$$\Rightarrow |a_{ii}| \leq \sum_{\substack{j=1 \\ j \neq i}}^n |a_{ij}| \frac{|x_j|}{|x_i|} \leq \sum_{\substack{j=1 \\ j \neq i}}^n |a_{ij}|$$

This contradicts the assumption that  $A$  is strictly diagonally dominant.

Note

In practice, the values  $a_i = S''(x_i)$  are found by Gaussian elimination for a tridiagonal system.

**Theorem** Let  $f$  be defined on  $[a, b]$ ,  $a = x_0 < x_1 < \dots < x_{n-1} < x_n = b$  and let  $S$  be the natural cubic spline interpolant of  $f$  (or clamped cubic spline).

$$1. \quad |f(x) - S(x)| \leq \frac{5}{384} \max_{a \leq x \leq b} |f^{(4)}(x)| \cdot h^4$$

where  $h = \max_i |x_{i+1} - x_i|$

$$\int_a^b [S''(x)]^2 dx \leq \int_a^b [f''(x)]^2 dx \quad \text{Minimum curvature property of cubic splines}$$

### Note

1. Cubic spline interpolation is 4<sup>th</sup> order accurate.
2. Condition (2) is optimality property: spline  $S(x)$  "oscillates least" of all smooth functions satisfying interpolation condition + BC (natural BC or clamped spline  $S'(x_0) = y'_0$ ,  $S'(x_n) = y'_n$ ).

Curvature

$$\kappa(x) = \frac{|f''(x)|}{(1 + [f'(x)]^2)^{3/2}} \approx |f''(x)|$$

$\Rightarrow \int_a^b [f''(x)]^2 dx$  is a crude measure of the total curvature over an interval.

$$\int_a^b [S''(x)]^2 dx \leq \int_a^b [f''(x)]^2 dx :$$

any smooth interpolating function must have a total curvature at least as large as that of the cubic spline (natural or clamped)

### Proof of (2) for natural splines

algebraic identity:  $F^2 - S^2 = (F - S)^2 - 2S(S - F)$

Let  $F = f''(x)$ ,  $S = S''(x)$ , integrate over  $[a, b]$ .

$$\begin{aligned} \int_a^b [f''(x)]^2 dx - \int_a^b [S''(x)]^2 dx &= \int_a^b [f''(x) - S''(x)]^2 dx \\ &\quad - 2 \int_a^b S''(x)(S''(x) - f''(x)) dx \end{aligned}$$

The first term is  $\geq 0$ . We will show that the second term is 0.

Integration by parts:

$$\begin{aligned} \int_a^b u(x)v'(x)dx &= u(x)v(x)|_a^b - \int_a^b u'(x)v(x)dx \\ \int_a^b S''(x)(S''(x) - f''(x))dx &= \sum_{i=0}^{n-1} \int_{x_i}^{x_{i+1}} S''(x)(S''(x) - f''(x))dx \\ &= \sum_{i=0}^{n-1} \left( S''(x)(S'(x) - f'(x))|_{x_i}^{x_{i+1}} - \int_{x_i}^{x_{i+1}} S'''(x)(S'(x) - f'(x))dx \right) \\ &= - \sum_{i=0}^{n-1} S'''(x_i) \int_{x_i}^{x_{i+1}} (S'(x) - f'(x))dx \end{aligned}$$

using  $S''(x_0) = S''(x_n) = 0$ ; telescoping sum and  $S'''_i(x)$  is constant

$$- \sum_{i=0}^{n-1} S'''(x_i)(S(x) - f(x))|_{x_i}^{x_{i+1}} = 0$$

using  $S(x_i) = f(x_i)$ ,  $i = 0, \dots, n$ .