

The FREQ Procedure

Frequency Percent Row Pct Col Pct	Table of judgeA by judgeB							
	judgeA	judgeB						Total
		4	5	6	7	8	9	
	5	0 0.00 0.00 0.00	0 0.00 0.00 0.00	1 10.00 100.00 50.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	1 10.00
	6	1 10.00 50.00 100.00	1 10.00 50.00 50.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	2 20.00
	7	0 0.00 0.00 0.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	1 10.00 50.00 33.33	1 10.00 50.00 100.00	2 20.00
	8	0 0.00 0.00 0.00	1 10.00 25.00 50.00	1 10.00 25.00 50.00	1 10.00 25.00 100.00	1 10.00 25.00 33.33	0 0.00 0.00 0.00	4 40.00
	9	0 0.00 0.00 0.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	0 0.00 0.00 0.00	1 10.00 100.00 33.33	0 0.00 0.00 0.00	1 10.00
	Total	1 10.00	2 20.00	2 20.00	1 10.00	3 30.00	1 10.00	10 100.00

The FREQ Procedure

Statistics for Table of judgeA by judgeB

Statistic	Value	ASE
Gamma	0.3125	0.2411
Kendall's Tau-b	0.2599	0.2040
Stuart's Tau-c	0.2500	0.2000
Somers' D C R	0.2703	0.2078
Somers' D R C	0.2500	0.2014
Pearson Correlation	0.4282	0.1924
Spearman Correlation	0.3751	0.2579
Lambda Asymmetric C R	0.2857	0.2415
Lambda Asymmetric R C	0.3333	0.3333
Lambda Symmetric	0.3077	0.2577
Uncertainty Coefficient C R	0.5095	0.0984
Uncertainty Coefficient R C	0.5874	0.0966
Uncertainty Coefficient Symmetric	0.5457	0.0899

Sample Size = 10

The CORR Procedure

2 Variables: judgeA judgeB

Simple Statistics						
Variable	N	Mean	Std Dev	Median	Minimum	Maximum
judgeA	10	7.20000	1.22927	7.50000	5.00000	9.00000
judgeB	10	6.60000	1.64655	6.50000	4.00000	9.00000

Kendall Tau b Correlation Coefficients, N = 10 Prob > tau under H0: Tau=0		
	judgeA	judgeB
judgeA	1.00000	0.25994 0.3394
judgeB	0.25994 0.3394	1.00000