

## Disjoint Cycles with Prescribed Lengths and Independent Edges in Graphs

Hong Wang

Department of Mathematics  
The University of Idaho  
Moscow, Idaho, USA 83844

We conjecture that if  $k \geq 2$  is an integer and  $G$  is a graph of order  $n$  with minimum degree at least  $(n + 2k)/2$ , then for any  $k$  independent edges  $e_1, \dots, e_k$  in  $G$  and for any integer partition  $n = n_1 + \dots + n_k$  with  $n_i \geq 4(1 \leq i \leq k)$ ,  $G$  has  $k$  disjoint cycles  $C_1, \dots, C_k$  of orders  $n_1, \dots, n_k$ , respectively such that  $C_i$  passes through  $e_i$  for all  $1 \leq i \leq k$ . We show that this conjecture is true for the case  $k = 2$ . The minimum degree condition is sharp in general.

Keywords: cycles, disjoint cycles, cycle coverings

MSC(2010): 05C38, 05C70, 05C75

### 1 Introduction

It is well known [8] that if a graph  $G$  of order  $n$  with minimum degree at least  $(n + 2)/2$ , then for each edge  $e$ ,  $G$  has a cycle of order  $l$  passing through  $e$  for each  $3 \leq l \leq n$ . A set of graphs are said to be disjoint if no two of them have any vertex in common. We ask this question: Given a graph  $G$  of order  $n = n_1 + \dots + n_k$  with  $n_i \geq 3(1 \leq i \leq k)$  and  $k$  independent edges  $e_1, \dots, e_k$  in  $G$ , when does  $G$  have  $k$  disjoint cycles of orders  $n_1, \dots, n_k$ , respectively such that  $C_i$  passes through  $e_i$  for each  $1 \leq i \leq k$ ? If the orders of the  $k$  cycles are not restricted, a similar problem was proposed in [7]. It was conjectured that for each integer  $k \geq 2$ , there exists  $n_0(k)$  such that if  $G$  is a graph of order  $n \geq n_0(k)$  and  $d(x) + d(y) \geq n + 2k - 2$ , then for any  $k$  independent edges  $e_1, \dots, e_k$  of  $G$ ,  $G$  has  $k$  disjoint cycles  $C_1, \dots, C_k$  covering all the vertices of  $G$  such that  $C_i$  passes through  $e_i$  for all  $1 \leq i \leq k$ . This conjecture was confirmed and completely solved by Egawa, Faudree, Györi, Ishigami, Schelp and Wang in [4]. Here we propose the following conjecture:

**Conjecture A** *Let  $k \geq 2$  be an integer and let  $G$  be a graph of order  $n$  with minimum degree at least  $(n + 2k)/2$ . Then for any  $k$  independent edges  $e_1, \dots, e_k$  in  $G$  and for any integer partition  $n = n_1 + \dots + n_k$  with  $n_i \geq 4(1 \leq i \leq k)$ ,  $G$  has  $k$  disjoint cycles  $C_1, \dots, C_k$  of orders  $n_1, \dots, n_k$ , respectively such that  $C_i$  contains  $e_i$  for all  $1 \leq i \leq k$ .*

To see the sharpness in general, we observe  $K_{(n-2(k-1))/2, (n-2(k-1))/2} + K_{2(k-1)}$ . This graph has minimum degree  $(n + 2k)/2 - 1$ . Let  $e_1, \dots, e_k$  be  $k$  independent edges such that  $e_1, \dots, e_{k-1}$  are taken from the clique  $K_{2(k-1)}$ . Let  $n = n_1 + \dots + n_k$  be such that  $n_k$  is odd. Then the graph does not contain  $k$  required cycles.

In Conjecture A, the condition  $n_i \geq 4(1 \leq i \leq k)$  is necessary in general. This can be demonstrated in the following example with  $n_i = 3(1 \leq i \leq k)$ . Choose positive integers  $a, b$  and  $k$  such that  $a \geq k/2 + 1$ ,  $b \geq 2$ ,  $k > a + b$  and  $k - b$  is even. Let  $K$  be the complete graph on  $V = \{x_1, y_1, \dots, x_k, y_k, z_1, \dots, z_k\}$ . Let  $(V, E)$  be a graph of order  $3k$  with  $V = \{x_1, y_1, \dots, x_k, y_k, z_1, \dots, z_k\}$  such that  $E = E(K) - \{y_i z_j | a + 1 \leq i \leq k, 1 \leq j \leq (k - b)/2\} - \{x_i z_j | a + 1 \leq i \leq k, (k - b)/2 + 1 \leq j \leq k - b\}$ . This graph does not contain  $k$  disjoint triangles containing  $k$  independent edges  $x_i y_i (1 \leq i \leq k)$  since  $k - b > a$  and a triangle containing a vertex of  $\{z_1, \dots, z_{k-b}\}$  and an edge of  $\{x_i y_i | 1 \leq i \leq k\}$  must contain an edge of  $\{x_i y_i | 1 \leq i \leq a\}$ . Its minimum degree is  $\min\{2k - 1 + (k + b)/2, 2k - 1 + a\} \geq 5k/2$ .

If the  $k$  disjoint cycles are not required to pass through given edges, we have El-Zahar's conjecture [5]. The conjecture says that if  $G$  is a graph of order  $n = n_1 + \dots + n_k$  with  $n_i \geq 3(1 \leq i \leq k)$  and minimum degree at

least  $\lceil n_1/2 \rceil + \dots + \lceil n_k/2 \rceil$  then  $G$  contains  $k$  disjoint cycles of order  $n_1, \dots, n_k$ , respectively. It was confirmed for the case  $k = 2$  in [5]. Abbasi[1] announced a solution of this conjecture for large  $n$  using regularity lemma.

In this paper, we prove Conjecture *A* for the case  $k = 2$ :

**Theorem B** *Let  $G$  be a graph of order  $n$  with minimum degree at least  $(n+4)/2$ . Then for any two independent edges  $e_1$  and  $e_2$  in  $G$  and for any integer partition  $n = n_1 + n_2$  with  $n_1 \geq 3$  and  $n_2 \geq 3$ ,  $G$  has two disjoint cycles  $C_1$  and  $C_2$  of orders  $n_1$  and  $n_2$ , respectively such that  $e_1 \in E(C_1)$  and  $e_2 \in E(C_2)$ .*

We shall use terminology and notation from [2] except as indicated. Let  $G = (V, E)$  be a graph. Let  $x \in V(G)$ . Let  $H$  be a subset of  $V(G)$  or a subgraph of  $G$ . We define  $N(x, H) = \{u \in N(x) | u \text{ belongs to } H\}$ . Let  $d(x, H) = |N(x, H)|$ . If  $X$  is a subset of  $V(G)$  or a subgraph of  $G$ , define  $N(X, H) = \cup_x N(x, H)$  and  $d(X, H) = \sum_x d(x, H)$  where  $x$  runs over  $X$ . Clearly, if  $X$  and  $H$  do not have any common vertex, then  $d(X, H)$  is the number of edges of  $G$  between  $X$  and  $H$ . We also use  $[H]$  to denote the induced subgraph of  $G$  by the vertices in  $H$ . For  $x, y \in V(G)$ , define  $I(xy, H) = N(x, H) \cap N(y, H)$  and let  $i(xy, H) = |I(xy, H)|$ . We use  $e(G)$  to denote  $|E(G)|$ . The order of  $G$  is denoted by  $|G|$ .

A path from  $u$  to  $v$  is called a  $u$ - $v$  path. If  $P$  is a path of  $G$  and  $v$  is an endvertex of  $P$ , we use  $\alpha(P, v)$  to denote the order of the longest  $u$ - $v$  subpath of  $P$  with  $uv \in E(G)$ . Clearly, if  $\alpha(P, v) \geq 3$  then  $P + uv$  has a cycle of order  $\alpha(P, v)$ . Let  $w \in V(G)$  ( $e \in E(G)$ , respectively). Let  $P = w_1 w_2 \dots w_t$  be a longest path starting at  $w = w_1$  ( $e = w_1 w_2$ , respectively). We say that  $P$  is an optimal path at  $w$  ( $e$ , respectively) in  $G$  if  $\alpha(P', x_t) \leq \alpha(P, w_t)$  for any longest path  $P' = x_1 x_2 \dots x_t$  starting at  $w = x_1$  ( $e = x_1 x_2$ , respectively) in  $G$ . If  $e \in E(P)$ , we define  $\sigma(P, e) = \min\{|E(P_1)|, |E(P_2)|\}$  where  $P_1$  and  $P_2$  are the two components of  $P - e$ . Thus if  $\sigma(P, e) = 0$  then  $e$  is an end edge of  $P$ . For an edge  $e \in E(G)$ , an  $e$ -path or  $e$ -cycle is a path or a cycle that passes through  $e$ . If  $P$  is a  $u$ - $v$  path, we define  $d^*(P, H) = d(uv, H)$ .

A cycle  $C$  of  $G$  is called an *end-cycle* at  $u \in V(C)$  if  $N(x, G) \subseteq V(C)$  and  $[C]$  has a  $u$ - $x$  hamiltonian path for each  $x \in V(C - u)$ .

If  $C = x_1 \dots x_t x_1$  is a cycle of  $G$ , we assume an orientation of  $C$  is given by default such that  $x_2$  is the successor of  $x_1$ . Then  $C[x_i, x_j]$  is the  $x_i$ - $x_j$  path on  $C$  along the orientation of  $C$  and  $C^-[x_i, x_j]$  is the  $x_i$ - $x_j$  path on  $C$  in the direction against the orientation of  $C$ . Define  $C[x_i, x_j] = C[x_i, x_j] - x_j$  and  $C(x_i, x_j) = C[x_i, x_j] - x_i$ . The predecessor and successor of  $x_i$  on  $C$  are denoted by  $x_i^-$  and  $x_i^+$ . We will use similar definitions for a path.

Let  $P = x_1 \dots x_t$  be a path of  $G$ . If  $\{x_1 x_{i+1}, x_t x_i\} \subseteq E$  with  $1 \leq i \leq t - 1$ , we say that  $x_i x_{i+1}$  is an *accessible edge* of  $P$ . Let  $C = u_1 u_2 \dots u_m u_1$  be a cycle of  $G$ . Let  $u_i$  and  $u_j$  be two distinct vertices of  $C$ . For each  $e \in E(C)$ , if  $e$  is an *accessible edge* of either  $C[u_i, u_j]$  or  $C[u_j, u_i]$ , then we say that  $e$  is an *accessible edge* of  $C$  w.r.t.  $\{u_i, u_j\}$ .

## 2 Proof of Theorem B

In this section, we list Lemmas 2.1-2.7 and use them to prove the theorem. The proofs of these lemmas are in Section 4. Let  $G = (V, E)$  be a graph order  $n$  with  $\delta(G) \geq (n+4)/2$ . Suppose, for a contradiction, that theorem fails for  $G$ . Let  $G$  be a counter example with  $n$  minimal. Let  $n = n_1 + n_2$  be an integer partition with  $n_1 \geq 3$  and  $n_2 \geq 3$  and let  $e_1$  and  $e_2$  be two independent edges such that  $G$  does not contain two disjoint cycles of orders  $n_1$  and  $n_2$  passing through  $e_1$  and  $e_2$ , respectively. For each  $X \subseteq V$  with  $|X| \leq 3$ ,  $\delta(G - X) \geq (n+4)/2 - |X| \geq ((n - |X|) + 1)/2$  and by Lemma 3.4,  $G - X$  is hamiltonian connected. Consequently, it is easy to see that if  $n_1 = 3$  or  $n_2 = 3$ , then  $G$  has the two required cycles, a contradiction. Therefore  $n_1 \geq 4$ ,  $n_2 \geq 4$  and so  $n \geq 8$ .

For the sake of convenience, for each  $i \in \{1, 2\}$ , let  $\mathcal{P}_i$  be the set of all the subgraphs of  $G$  which have  $e_i$ -hamiltonian paths and  $\mathcal{H}_i$  the set of all the subgraphs of  $G$  which have  $e_i$ -hamiltonian cycles. Furthermore, for each  $i \in \{1, 2\}$  and  $J \in \mathcal{P}_i$ , let  $\mathcal{P}_i(J)$  denote the set of all the  $e_i$ -hamiltonian paths of  $J$  and let  $\mathcal{P}_i^*(J)$  denote the subset of  $\mathcal{P}_i(J)$  such that a path  $P \in \mathcal{P}_i(J)$  belongs to  $\mathcal{P}_i^*(J)$  if and only if  $\sigma(P, e_i) \geq 1$ .

For each  $i \in \{1, 2\}$  and  $J \in \mathcal{P}_i$ , let  $S_i(J)$  be the set of all the vertices  $x$  of  $J - V(e_i)$  such that  $x$  is an end vertex of some  $P \in \mathcal{P}_i(J)$  and let  $\delta_i(J) = \min\{d(x, J) | x \in S_i(J)\}$ .

As  $\delta(G) \geq (n+4)/2$ ,  $G$  has a hamiltonian cycle containing both  $e_1$  and  $e_2$ . Thus  $G$  has two disjoint subgraphs  $G_1$  and  $G_2$  such that for each  $i \in \{1, 2\}$ ,  $|G_i| = n_i$  and  $G_i \in \mathcal{P}_i$ . We choose  $G_1$  and  $G_2$  such that

$$e(G_1) + e(G_2) \text{ is maximum.} \quad (1)$$

Let  $P_1 = x_1 \dots x_{n_1}$  and  $P_2 = y_1 \dots y_{n_2}$  be two paths such that,  $P_1 \in \mathcal{P}_1(G_1)$ ,  $P_2 \in \mathcal{P}_2(G_2)$ ,  $x_1 \in S_1(G_1)$ ,  $y_1 \in S_2(G_2)$ ,  $d(x_1, G_1) = \delta_1(G_1)$  and  $d(y_1, G_2) = \delta_2(G_2)$ . For any  $x \in V(G_1)$  and  $y \in V(G_2)$ , we use  $\xi(x, y)$  to denote  $d(x, G_2) - d(x, G_1) + d(y, G_1) - d(y, G_2) - 2d(x, y)$ . Thus  $e(G_1 - x + y) + e(G_2 - y + x) = e(G_1) + e(G_2) + \xi(x, y)$ . By (1), we readily obtain the following Property A and Property B. The first one is evident.

*Property A* Let  $x \in V(G_1)$  and  $y \in V(G_2)$ . If  $G_1 - x + y \in \mathcal{P}_1$  and  $G_2 - y + x \in \mathcal{P}_2$  then  $\xi(x, y) \leq 0$ . ■

*Property B* Either  $\mathcal{P}_1^*(G_1) \neq \emptyset$  or  $\mathcal{P}_2^*(G_2) \neq \emptyset$ .

*Proof of Property B.* Say  $\mathcal{P}_1^*(G_1) = \emptyset$  and  $\mathcal{P}_2^*(G_2) = \emptyset$ . Then  $e_1 = x_{n_1-1}x_{n_1}$  and  $N(x_{n_1}, G_1) \subseteq \{x_{n_1-1}, x_{n_1-2}\}$ . Thus  $n_2 \geq d(x_{n_1}, G_2) \geq (n_1 + n_2 + 4)/2 - 2 = (n_1 + n_2)/2$  and so  $n_2 \geq n_1$ . Similarly,  $n_1 \geq (n_1 + n_2)/2$ . It follows that  $n_1 = n_2$ ,  $N(x_{n_1}, G_1) = \{x_{n_1-2}, x_{n_1-1}\}$  and  $N(y_{n_2}, G_2) = \{y_{n_2-2}, y_{n_2-1}\}$ . Thus  $N(e_1, G_1) = \{x_{n_1-2}, x_{n_1-1}, x_{n_1}\}$  and  $N(e_2, G_2) = \{y_{n_2-2}, y_{n_2-1}, y_{n_2}\}$ . Consequently,  $G_1 - V(e_1) + V(e_2) \in \mathcal{P}_2$ ,  $G_2 - V(e_2) + V(e_1) \in \mathcal{P}_1$ ,  $e(G_1 - V(e_1) + V(e_2)) + e(G_2 - V(e_2) + V(e_1)) > e(G_1) + e(G_2)$ . This contradicts (1). ■

To reach a contradiction, we will investigate the structure of  $G_1$  and  $G_2$  which lead us to construct a sequence  $(G_1, G_2), (G_3, G_4), \dots, (G_{2k-1}, G_{2k})$  of pairs of disjoint subgraphs of  $G$ . This will be accomplished by seven lemmas. Lemmas 2.1-2.6 are the steps to Lemma 2.7 and we use Lemma 2.7 to show that the sequence yields a contradiction.

**Lemma 2.1** *Either  $d(x_1, G_1) \leq (n_1 + 1)/2$  or  $d(y_1, G_2) \leq (n_2 + 1)/2$ .*

**Lemma 2.2** *Either  $d(x_1, G_1) \geq (n_1 + 2)/2$  or  $d(y_1, G_2) \geq (n_2 + 2)/2$ .*

By Lemma 2.1 and Lemma 2.2, we may assume w.l.o.g. that  $d(x_1, G_1) \leq (n_1 + 1)/2$  and  $d(y_1, G_2) \geq (n_2 + 2)/2$ , i.e.,  $\delta_1(G_1) \leq (n_1 + 1)/2$  and  $\delta_2(G_2) \geq (n_2 + 2)/2$ . Clearly,  $d(x_1, G_2) \geq (n_2 + 3)/2$ .

**Lemma 2.3**  $G_2 \notin \mathcal{H}_2$ .

By Lemma 2.3,  $G_2 \notin \mathcal{H}_2$ . As  $\delta_2(G_2) \geq (n_2 + 2)/2$  and by Lemma 3.3,  $\mathcal{P}_2^*(G_2) = \emptyset$ . Let  $P = v_{n_2}v_{n_2-1} \dots v_1$  be an optimal path of  $G_2$  at  $e_2 = v_{n_2}v_{n_2-1}$ . Say  $\alpha(P, v_1) = r$ . As  $G_2 \notin \mathcal{H}_2$ ,  $r \leq n_2 - 1$ . As  $\delta_2(G_2) \geq (n_2 + 2)/2$  and by Lemma 3.9,  $J = v_1v_2 \dots v_rv_1$  is an end-cycle at  $v_r$  in  $G_2$  such that  $d(v_i, J) \geq (n_2 + 2)/2$  for all  $i \in \{1, \dots, r-1\}$ . Let  $J^* = \{v_2, v_3, \dots, v_{r-2}\}$ . Clearly,  $r \geq (n_2 + 2)/2 + 1 = (n_2 + 4)/2$ .

**Lemma 2.4** *There exists no  $u \in V(G_1) - V(e_1)$  such that  $G_1 - u \in \mathcal{P}_1$ ,  $G_2 + u \in \mathcal{H}_2$  and  $d(u, J^*) > 0$ .*

**Lemma 2.5**  $\delta_1(G_1) \leq (n_1 - 1)/2$ .

Let  $w_1 \in S_1(G_1)$  with  $d(w_1, G_1) = \delta_1(G_1)$ . Then  $d(w_1, G_2) \geq (n_1 + n_2 + 4)/2 - (n_1 - 1)/2 = (n_2 + 5)/2$ . Clearly,  $d(w_1, J) \geq (n_2 + 5)/2 - (n_2 - r) \geq 9/2$ . Thus  $d(w_1, J^*) > 0$ . By Lemma 2.4,  $G_2 + w_1 \notin \mathcal{H}_2$ . This implies that  $w_1v_{n_2} \notin E$  and if  $v_{n_2}v_{n_2-2} \in E$  then  $w_1v_{n_2-1} \notin E$ . Hence  $\mathcal{P}_2^*(G_2 + w_1) = \emptyset$ . For each  $v \in S_2(G_2 + w_1)$ , if  $d(v, G_2 + w_1) \leq (n_2 + 4)/2$ , then  $d(v, G_1 - w_1) \geq n_1/2$  and so  $G_1 - w_1 + v \in \mathcal{P}_1$  by Lemma 3.2(a). But  $e(G_1 - w_1 + v) + e(G_2 + w_1 - v) > e(G_1) + e(G_2)$ , contradicting (1). Hence  $\delta_2(G_2 + w_1) \geq (n_1 + 5)/2$ . In the meantime, we see that  $n_2 - 1 \geq \lceil (n_2 + 5)/2 \rceil$ . Thus  $n_2 \geq 7$ . With  $G_1 - w_1$  and  $G_2 + w_1$ , this argument also implies the existence of the following two subgraphs  $G_3$  and  $G_4$ .

Let  $G_3$  and  $G_4$  be two disjoint subgraphs of  $G$  with  $e(G_3) + e(G_4)$  maximal such that  $|G_3| = n_1 - 1$ ,  $|G_4| = n_2 + 1$ ,  $G_3 \in \mathcal{P}_1$ ,  $G_4 \in \mathcal{P}_2$  and  $\mathcal{P}_2^*(G_4) = \emptyset$ . By the above argument,  $e(G_3) + e(G_4) \geq e(G_1) + e(G_2) - (n_1 - 1)/2 + (n_2 + 5)/2$ . If  $d(v, G_4) \leq (|G_4| + 3)/2$  for some  $v \in S_2(G_4)$ , then  $d(v, G_3) \geq (|G_3| + 1)/2$  and  $e(G_3 + v) + e(G_4 - v) > e(G_1) + e(G_2)$ . This contradicts (1) since  $G_3 + v \in \mathcal{P}_1$  by Lemma 3.2(a). Thus

$\delta_2(G_4) \geq (n_2 + 5)/2 = (|G_4| + 4)/2$ . This argument is the key for a generalization leading to the following definition and the proofs of Lemma 2.6 and Lemma 2.7.

Let  $k \geq 2$  be the largest integer such that there exist a sequence  $(G_1, G_2), (G_3, G_4), \dots, (G_{2k-1}, G_{2k})$  of disjoint pairs of subgraphs of  $G$  such that for each  $i \in \{1, \dots, k-1\}$ ,  $G_{2i-1} \in \mathcal{P}_1$ ,  $G_{2i} \in \mathcal{P}_2$ ,  $\mathcal{P}_2^*(G_{2i}) = \emptyset$  and there exists  $w_i \in S_1(G_{2i-1})$  such that  $\delta_1(G_{2i-1}) = d(w_i, G_{2i-1}) \leq (|G_{2i-1}| - 1)/2$ ,  $d(w_i, G_{2i}) \geq (|G_{2i}| + 5)/2$  and  $G_{2i} + w_i \notin \mathcal{H}_2$ . Moreover, for each  $i \in \{1, \dots, k-1\}$ ,  $e(G_{2i+1}) + e(G_{2i+2})$  is maximal such that  $|G_{2i+1}| = |G_{2i-1}| - 1$ ,  $|G_{2i+2}| = |G_{2i}| + 1$ ,  $G_{2i+1} \in \mathcal{P}_1$ ,  $G_{2i+2} \in \mathcal{P}_2$  and  $\mathcal{P}_2^*(G_{2i+2}) = \emptyset$ . By the above argument,  $k$  is well defined.

**Lemma 2.6** *The following two statements hold:*

- (a) *For each  $i \in \{1, \dots, k\}$ ,  $|G_{2i-1}| = n_1 - i + 1$  and  $|G_{2i}| = n_2 + i - 1$ .*
- (b) *For each  $i \in \{1, \dots, k\}$ ,  $\delta_2(G_{2i}) \geq (|G_{2i}| + 4)/2$ .*

Say  $s = |G_{2k-1}|$  and  $|G_{2k}| = t$ . As  $n_2 \geq 7$ ,  $t \geq 8$ . By Lemm 2.6,  $\delta_2(G_{2k}) \geq (t+4)/2$ . Let  $L = y_t y_{t-1} \dots y_1$  be an optimal path at  $e_2 = y_t y_{t-1}$  in  $G_{2k}$ . Say  $r = \alpha(L, y_1)$ . Then  $r \geq \delta_2(G_{2k}) + 1 \geq \lceil (t+4)/2 + 1 \rceil = \lceil (t+6)/2 \rceil \geq 7$ . As  $\mathcal{P}_2^*(G_{2k}) = \emptyset$ ,  $r \leq t - 1$ . Let  $R = [y_1, y_2, \dots, y_r]$  and  $R' = R - y_r$ . By Lemma 2.6 and Lemma 3.9,  $y_1 y_2 \dots y_r y_1$  is an end-cycle at  $y_r$  in  $G_{2k}$  and so  $\delta(R') \geq (t+4)/2 - 1 \geq (|R'| + 4)/2$ . By the minimality of  $|G|$ , Theorem B holds for  $R'$ . Note that  $R' - \{x, y\}$  is hamiltonian connected for all  $\{x, y\} \subseteq V(R')$  by Lemma 3.4. Clearly,  $s \geq d(y_t, G_{2k-1}) \geq (s+t+4)/2 - 2 = (s+t)/2$ . This implies that  $s \geq t$  and if equality holds then  $N(y_t, G_{2k}) = \{y_{t-1}, y_{t-2}\}$  and  $r \leq t - 2$ .

**Lemma 2.7** *For no  $x \in V(G_{2k-1})$ ,  $G_{2k-1} - x \in \mathcal{P}_1$ ,  $G_{2k} + x \in \mathcal{H}_2$  and  $d(x, R' - \{y_1, y_{r-1}\}) > 0$ .*

To prove Theorem B, let  $y_c \in V(R' - \{y_1, y_{r-1}\})$ . Then  $d(y_c y_t, G_{2k-1}) \geq s + t + 4 - (t - 1) = s + 5$  and so  $i(y_c y_t, G_{2k-1}) \geq 5$ . By Lemma 2.7,  $G_{2k-1} - x \notin \mathcal{P}_1$  for all  $x \in I(y_c y_t, G_{2k-1})$  and so  $G_{2k-1} \notin \mathcal{H}_1$ . If  $\delta_1(G_{2k-1}) \leq (s-1)/2$ , let  $w_k \in S_1(G_{2k-1})$  with  $d(w_k, G_{2k-1}) = \delta_1(G_{2k-1})$ . As  $d(w_k, G_{2k}) \geq (t+5)/2$ ,  $d(w_k, R' - \{y_1, y_{r-1}\}) \geq 1$ . By Lemma 2.7,  $G_{2k} + w_k \notin \mathcal{H}_2$ . Thus  $w_k y_t \notin E$  and if  $y_t y_{t-2} \in E$  then  $w_k y_{t-1} \notin E$ . Therefore  $\mathcal{P}_2^*(G_{2k} + w_k) = \emptyset$ . This allows us to define  $(G_{2k+1}, G_{2k+2})$  to lengthen the sequence  $(G_1, G_2), \dots, (G_{2k-1}, G_{2k})$ . This contradicts the maximality of  $k$ . Therefore  $\delta_1(G_{2k-1}) \geq s/2$ . Recall that  $d(y_t, G_{2k-1}) \geq (s+t)/2$ . If  $\mathcal{P}_1^*(G_{2k-1}) \neq \emptyset$ , then by Lemma 3.5(c), we see that  $G_{2k-1}$  has a  $u$ - $v$   $e_1$ -hamiltonian path such that  $v \notin V(e_1)$ ,  $d(v, G_{2k-1}) = s/2$  and  $v y_t \in E$ . As  $d(v, G_{2k}) \geq (t+4)/2$ ,  $d(v, R' - \{y_1, y_{r-1}\}) > 0$  and so  $G_{2k} + v \in \mathcal{H}_2$ , contradicting Lemma 2.7. Therefore  $\mathcal{P}_1^*(G_{2k-1}) = \emptyset$ . Let  $P = z_s z_{s-1} \dots z_1$  be an optimal path at  $e_1 = z_s z_{s-1}$  in  $G_{2k-1}$ . Say  $\alpha(P, z_1) = q$ . As  $d(z_s, G_{2k-1}) \leq 2$ ,  $t \geq d(z_s, G_{2k}) \geq (s+t+4)/2 - 2$  and so  $t \geq s$ . Since  $s \geq t$ , it follows that  $s = t$  and  $d(z_s, G_{2k}) = t = d(y_t, G_{2k-1})$ . By Lemma 2.7, we see that  $d(z_i, R' - \{y_1, y_{r-1}\}) = 0$  for all  $i \in \{1, \dots, q-1\}$ . Then  $t + 2 \leq d(y_c, G) \leq r - 1 + d(y_c, G_{2k-1}) \leq r - 1 + t - q + 1 = t + r - q$ . Thus  $r - q \geq 2$ . Then  $t + 2 \leq d(z_1, G) \leq q - 1 + d(z_1, G_{2k}) \leq q - 1 + t - r + 3 \leq t$ , a contradiction. This proves the theorem.

### 3 Auxiliary Lemmas

In the following,  $G = (V, E)$  is a graph. We will use the following lemmas. Lemma 3.1 is an easy observation.

**Lemma 3.1** *Let  $P = x_1 \dots x_r$  be a path of order  $r$  in  $G$ . Let  $u$  and  $v$  be two vertices of  $G - V(P)$ . Suppose that  $d(uv, P) \geq r + 1$  and  $\{u x_{i+1}, v x_i\} \not\subseteq E$  for all  $i \in \{1, \dots, r-1\}$ . Then  $d(uv, P) = r + 1$  and  $\{u x_1, v x_r\} \subseteq E$ . Moreover, either  $N(u, P) = \{x_1, \dots, x_a\}$  and  $N(v, P) = \{x_a, \dots, x_r\}$  for some  $a \in \{1, \dots, r\}$ , or  $d(x_i, uv) = 0$  for some  $1 < i < r$ . ■*

**Lemma 3.2** *Let  $P$  be a  $u$ - $v$  path of order  $r$  in  $G$ ,  $e \in E(P)$  and  $x \in V(G) - V(P)$ . The following five statements hold:*

- (a) *If  $d(x, P) > r/2$ , then  $P + x$  has an  $e$ -hamiltonian path.*

- (b) If  $d(x, P) > (r + 1)/2$ ,  $P + x$  has an  $e$ -hamiltonian path ending at  $v$ .
- (c) If  $d(x, P) > (r + 2)/2$  then  $P + x$  has a  $u$ - $v$   $e$ -hamiltonian path.
- (d) If  $d(xv, P) \geq r + 2$  then  $[P + x]$  has a  $u$ - $x$   $e$ -hamiltonian path.
- (e) If  $d(xv, P) \geq r + 1$  then  $[P + x]$  has an  $e$ -hamiltonian path.
- (f) If  $d(x, P) > (r + 1)/2$  and  $uv \in E$  then  $P + uv + x$  has an  $e$ -hamiltonian cycle.

**Proof.** Let  $P_1$  and  $P_2$  be the two components of  $P - e$  with  $v$  in  $P_2$ . If  $d(x, f) = 2$  for some  $f \in E(P - e)$  then (a), (b) and (c) hold. So if one of (a), (b) and (c) fails then  $d(x, f) \leq 1$  for all  $f \in E(P) - \{e\}$ . This implies that  $d(x, P_i) \leq (|P_i| + 1)/2$  for  $i \in \{1, 2\}$  and so  $d(x, P) \leq (r + 2)/2$ . Furthermore, for each  $i \in \{1, 2\}$ , if  $d(x, P_i) = (|P_i| + 1)/2$ , then  $|P_i|$  is odd and  $x$  is adjacent to the two endvertices of  $P_i$  and so the first three statements follow.

If one of (d) and (e) fails, then  $\{vz, xz^+\} \not\subseteq E$  for each  $z \in V(P)$  with  $zz^+ \neq e$ . This implies that  $d(xv, P) \leq r + 1$ . So (d) holds. Obviously, (e) would hold if  $uv \in E$  or  $d(x, uv) > 0$ . To see (e), say  $uv \notin E$  and  $d(x, uv) = 0$ . Then apply (d) to  $P - u$  and  $x$ .

To obtain (f), we see that there exists an edge  $e'$  on  $P + uv$  with  $e' \neq e$  such that  $d(x, e') = 2$ . ■

**Lemma 3.3** *Let  $P$  be a  $u$ - $v$  path of order  $r \geq 3$  in  $G$ . Let  $e \in E(P)$ . Suppose that  $d(uv, P) \geq r + \epsilon$  where  $\epsilon = 0$  if  $\sigma(P, e) = 0$  and  $\epsilon = 1$  if  $\sigma(P, e) > 0$ . Then  $[P]$  has an  $e$ -hamiltonian cycle.*

**Proof.** If  $uv \in E$ , nothing to prove. So assume  $uv \notin E$ . Then the condition implies that some edge  $f \in E(P) - \{e\}$  is an accessible edge and this yields a required cycle. ■

**Lemma 3.4** [3] *If  $H$  is a graph of order  $r \geq 3$  and  $d(xy, H) \geq r + 1$  for each pair  $x$  and  $y$  of nonadjacent vertices of  $H$ , then  $H$  is hamiltonian connected and so for each  $e \in E(H)$ ,  $H$  has an  $e$ -hamiltonian cycle.*

**Lemma 3.5** *Let  $P = x_1 \dots x_r$  be a path of order  $r \geq 3$  in  $G$ . Let  $e \in E(P)$ . Suppose that  $[P]$  does not have an  $e$ -hamiltonian cycle and  $d(x_1x_r, P) \geq r$ . Let  $R = \{x_i | d(x_i, x_1x_r) = 0, 1 < i < r\}$  and  $\mathcal{P}$  be the set of all the components of  $P - R \cup \{x_1, x_r\} - e$ . Then  $\sigma(e, P) > 0$ ,  $d(x_1x_r, P) = r$  and the following three statements hold:*

- (a)  $R \cup \{x_1, x_r\}$  is an independent set;
- (b)  $d(x_l, P') \leq 1$  for all  $x_l \in R$  and  $P' \in \mathcal{P}$ ;
- (c) If  $d^*(L, P) \geq r$  for every  $e$ -hamiltonian path  $L$  of  $[P]$  with  $\sigma(L, e) > 0$ , then either  $V(P)$  has a partition  $X \cup Y$  such that  $|X| = r/2$ ,  $V(e) \subseteq X$ ,  $Y = R \cup \{x_1, x_r\}$  and  $N(y, P) = X$  for all  $y \in Y$ , or  $[P] - V(e)$  has two complete components  $H_1$  and  $H_2$  such that  $|H_1| + |H_2| = r - 2$  and  $V(H_1 \cup H_2) \subseteq N(x)$  for each  $x \in V(e)$ .

**Proof.** By Lemma 3.3,  $\sigma(e, P) > 0$  and  $d(x_1x_r, P) = r$ . Clearly,  $|\mathcal{P}| \leq |R| + 2$  and  $|\mathcal{P}| + |R| \leq \sum_{P' \in \mathcal{P}} |P'| + |R| \leq r - 2$ . Say  $e = x_a x_{a+1}$ . Since  $[P]$  does not have an  $e$ -hamiltonian cycle, each  $x_i x_{i+1}$  with  $i \neq a$  is not an accessible edge of  $P$ . By Lemma 3.1,  $d(x_1x_r, P') \leq |P'| + 1$  for each  $P' \in \mathcal{P}$ . Thus  $d(x_1x_r, P) \leq (r - 2) - |R| + |\mathcal{P}| \leq r$ . It follows that  $|\mathcal{P}| = |R| + 2$  and  $d(x_1x_r, P') = |P'| + 1$  for each  $P' \in \mathcal{P}$ . Consequently,  $\{x_2, x_a, x_{a+1}, x_{r-1}\} \cap R = \emptyset$ ,  $R$  does not contain two consecutive vertices of  $P$ , and for each  $P' = P[x_i, x_j] \in \mathcal{P}$  there exists  $i \leq k \leq j$  such that  $N(x_1, P') = \{x_i, \dots, x_k\}$  and  $N(x_r, P') = \{x_k, \dots, x_j\}$ . In particular,  $\{x_1x_{a+1}, x_r x_a\} \subseteq E$ . It is easy to see that  $R$  is an independent set for otherwise  $[P]$  has an  $e$ -hamiltonian cycle. So (a) holds.

To see (b), say  $d(x_l, P') \geq 2$  for some  $x_l \in R$  and  $P' = P[x_i, x_j] \in \mathcal{P}$ . Let  $x_k \in V(P')$  be such that  $N(x_1, P') = \{x_i, \dots, x_k\}$  and  $N(x_r, P') = \{x_k, \dots, x_j\}$ . Say w.l.o.g. that  $l < i$ . Let  $x_p \in V(P')$  be such that  $x_l x_p \in E$  and  $p \neq i$ . If  $p \leq k$ , then  $x_1 P[x_1, x_{l-1}] P^- [x_r, x_p] x_l P[x_{l+1}, x_{p-1}] x_1$  is an  $e$ -hamiltonian cycle of  $[P]$  and if  $p > k$  then  $x_1 P[x_1, x_l] P[x_p, x_r] P^- [x_{p-1}, x_{l+1}] x_1$  is an  $e$ -hamiltonian cycle of  $[P]$ , a contradiction. Hence (b) holds.

To see (c), it is easy to observe that for each  $x_l \in R$ ,  $[P]$  has an  $x_1$ - $x_l$   $e$ -hamiltonian path and an  $x_r$ - $x_l$   $e$ -hamiltonian path. If  $R \neq \emptyset$ , then  $d(x_l x_1, P) \geq r$ ,  $d(x_l x_r, P) \geq r$  and so  $d(x_l, P) \geq r/2$  for each  $x_l \in R$ . Since  $|\mathcal{P}| = |R| + 2$  and  $|\mathcal{P}| + |R| \leq r - 2$ , it follows that  $|\mathcal{P}| = r/2$  and  $|P'| = 1$  for all  $P' \in \mathcal{P}$ . Thus  $X \cup Y$  with  $Y = R \cup \{x_1, x_r\}$  and  $X = V(P) - Y$  is a partition of  $V(P)$  satisfying (c). Next, assume that  $R = \emptyset$ . Let  $2 \leq b \leq a$  and  $a + 1 \leq c \leq r - 1$  be such that  $N(x_1, P) = \{x_2, \dots, x_b\} \cup \{x_{a+1}, \dots, x_c\}$  and

$N(x_r, P) = \{x_b, \dots, x_a\} \cup \{x_c, \dots, x_{r-1}\}$ . Then we readily see that for each  $x_i \in N(x_1, P) - \{x_b, x_a, x_{a+1}, x_c\}$  and  $x_j \in N(x_r, P) - \{x_b, x_a, x_{a+1}, x_c\}$ ,  $[P]$  has an  $x_i$ - $x_r$   $e$ -hamiltonian path, an  $x_1$ - $x_j$   $e$ -hamiltonian path, an  $x_i$ - $x_j$   $e$ -hamiltonian path and so  $x_i x_j \notin E$ . It follows that  $N(x_i, P) \cup \{x_i\} = N(x_1, P) \cup \{x_1\}$  and  $N(x_j, P) \cup \{x_j\} = N(x_r, P) \cup \{x_r\}$  for all  $x_i \in N(x_1, P) - \{x_b, x_a, x_{a+1}, x_c\}$  and  $x_j \in N(x_r, P) - \{x_b, x_a, x_{a+1}, x_c\}$ . Thus if  $b < a$  then  $x_1 P^- [x_c, x_{b+1}] P^- [x_r, x_{c+1}] P^- [x_b, x_1]$  is an  $e$ -hamiltonian cycle of  $[P]$ , a contradiction. Hence  $b = a$ . Similarly,  $c = a + 1$ . This proves (c).  $\blacksquare$

**Lemma 3.6** *Let  $C$  be a cycle of order  $r$  in  $G$ . Let  $u$  and  $v$  be two distinct vertices on  $C$  and  $e$  an edge of  $C$  with  $e \notin \{uu^+, vv^+\} = \emptyset$ . Set  $R = \{x | d(x, uv) = 0, x \in V(C) - \{u, v\}\}$ . Let  $\mathcal{P}$  be the set of all the components of  $C - (R \cup \{u, v\}) - e$ . Suppose that  $d(uv, C) \geq r + 1$  and  $[C]$  does not have a  $u^+v^+$   $e$ -hamiltonian path. Then  $d(uv, C) = r + 1$  and the following four statements hold:*

- (a) *Each edge of  $C - e$  is inaccessible on  $C$  w.r.t.  $\{u, v\}$ ;*
- (b)  *$V(e) \cap (R \cup \{u, v\}) = \emptyset$ ,  $d(uv, P) = |P| + 1$  for all  $P \in \mathcal{P}$  and  $|\mathcal{P}| = |R| + 3$ .*
- (c)  *$R$  is an independent set and  $d(x, P) \leq 1$  for all  $x \in R$  and  $P \in \mathcal{P}$ .*
- (d) *If  $d(z, C) \geq (r + 1)/2$  for all  $z \in V(C) - V(e)$  then  $r$  is odd. Moreover, either  $[C]$  has a vertex-cut  $X$  with  $V(e) \subseteq X$  and  $|X| = 3$  such that  $[C]$  has exactly two components isomorphic to  $K_{(r-3)/2}$  and  $X \subseteq N(y)$  for all  $y \in V(C) - X$ , or  $V(C)$  has a partition  $X \cup Y$  such that  $|X| = (r + 1)/2$ ,  $|Y| = (r - 1)/2$ ,  $Y = R \cup \{u, v\}$ ,  $V(e) \subseteq X$  and  $N(y, C) = X$  for all  $y \in Y$ .*

**Proof.** It is easy to check that (a) holds since  $[C]$  does not have a  $u^+v^+$   $e$ -hamiltonian path. In particular,  $uv \notin E$ . Clearly,  $|\mathcal{P}| \leq |R| + 3$  and  $|\mathcal{P}| + |R| \leq \sum_{P \in \mathcal{P}} |P| + |R| = r - 2$ . By (a) and Lemma 3.1,  $d(uv, P) \leq |P| + 1$  for each  $P \in \mathcal{P}$  and so  $d(uv, C) \leq r + 1$ . Since  $d(uv, C) \geq r + 1$ , it follows that  $d(uv, C) = r + 1$ ,  $|\mathcal{P}| = |R| + 3$ ,  $V(e) \cap (R \cup \{u, v\}) = \emptyset$ , and  $d(uv, P) = |P| + 1$  for all  $P \in \mathcal{P}$ . So (b) holds.

As  $|\mathcal{P}| = |R| + 3$ ,  $R$  does not contain two consecutive vertices of  $C$ . To prove (c), Let  $C = x_1 \dots x_r x_1$  be such that  $x_1 = u$ ,  $x_2 = u^+$ ,  $x_p = v$  and  $x_{p+1} = v^+$ . W.l.o.g., say  $e = x_q x_{q+1}$  for some  $q \in \{p + 1, \dots, r - 1\}$ . We first check that  $R$  is an independent set. Let  $L_1 = C(x_1, x_p)$ ,  $L_2 = C(x_p, x_q]$  and  $L_3 = C[x_{q+1}, x_r]$ . Let  $R_i = R \cap V(L_i)$  for  $i \in \{1, 2, 3\}$ . Say  $x_i x_j \in E$  for some  $\{x_i, x_j\} \subseteq R$  with  $i < j$ . We shall obtain a contradiction by showing that  $[C]$  has an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path. According to the locations of  $x_i$  and  $x_j$  in  $R = R_1 \cup R_2 \cup R_3$ , there are six cases to check, which are very similar in the verification. So we just show one example with  $x_i \in R_1$  and  $x_j \in R_3$ . In this case,  $\{x_1 x_{i+1}, x_p x_{i-1}\} \subseteq E$  and  $\{x_1 x_{j-1}, x_p x_{j+1}\} \subseteq E$  by (a), (b) and Lemma 3.1. Then

$$x_2 C[x_2, x_{i-1}] x_p C^- [x_p, x_i] x_i x_j C[x_j, x_1] x_{j-1} C^- [x_{j-1}, x_{p+1}]$$

is an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path of  $[C]$ , a contradiction.

Next, we show that  $d(x, P) \leq 1$  for all  $x \in R$  and  $P \in \mathcal{P}$ . On the contrary, say  $d(x, P) \geq 2$  for some  $x \in R$  and  $P \in \mathcal{P}$ . We shall obtain a contradiction by showing that  $[C]$  has an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path. According to the locations of  $x$  in  $R_1 \cup R_2 \cup R_3$  and  $P$  on  $L_1 \cup L_2 \cup L_3$ , there are nine cases to check, which are also very similar in the verification. So we just show one example with  $x \in R_1$  and  $P$  on  $L_3$ . Say  $P = C[x_i, x_j]$ . By (a), (b) and Lemma 3.1,  $N(x_1, P) = \{x_a, \dots, x_j\}$  and  $N(x_p, P) = \{x_i, \dots, x_a\}$  for some  $i \leq a \leq j$ . Since  $d(x, P) \geq 2$ ,  $x x_t \in E$  for some  $x_t \in V(P)$  with  $t \neq x_i$ . If  $t > a$ , then

$$x_2 C[x_2, x^-] x_p C^- [x_p, x] x x_t C[x_t, x_1] x_{t-1} C^- [x_{t-1}, x_{p+1}]$$

is an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path of  $[C]$ , a contradiction. Thus  $t \leq a$ . Then

$$x_2 C[x_2, x] x x_t C[x_t, x_1] x^+ C[x^+, x_p] x_{t-1} C^- [x_{t-1}, x_{p+1}]$$

is an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path of  $[C]$ , a contradiction.

To prove (d), we have  $d(x, C) \leq |\mathcal{P}|$  for all  $x \in R$  by (c). Since  $|\mathcal{P}| \leq r - |R| - 2$  and  $|\mathcal{P}| = |R| + 3$ , we obtain  $d(x, C) \leq (r + 1)/2$  for all  $x \in R$ . It follows that if  $R \neq \emptyset$  then  $r$  is odd and  $|\mathcal{P}| = 1$  for all  $P \in \mathcal{P}$ .

Consequently, if  $Y = R \cup \{u, v\}$  and  $X = V(C) - Y$  then  $N(y, C) = X$  for all  $y \in Y$  and so (d) holds. So assume that  $R = \emptyset$ . By (a), (b) and Lemma 3.1, there exists  $x_{a_i} \in V(L_i)$  for  $i \in \{1, 2, 3\}$  such that

$$\begin{aligned} N(x_1, C) &= V(L_1[x_2, x_{a_1}]) \cup V(L_2[x_{a_2}, x_q]) \cup V(L_3[x_{a_3}, x_r]) \\ N(x_p, C) &= V(L_1[x_{a_1}, x_{p-1}]) \cup V(L_2[x_{p+1}, x_{a_2}]) \cup V(L_3[x_{q+1}, x_{a_3}]). \end{aligned}$$

We claim that for each vertex  $x$  of  $L_1[x_2, x_{a_1}] \cup L_2(x_{a_2}, x_q) \cup L_3(x_{a_3}, x_r)$ ,  $N(x, C) \subseteq N(x_1, C) \cup \{x_1\}$ . If this is false, say  $xy \in E(G)$  for some vertex  $x$  of  $L_1[x_2, x_{a_1}] \cup L_2(x_{a_2}, x_q) \cup L_3(x_{a_3}, x_r)$  and  $y \in V(C) - N(x_1, C) - \{x_1\}$ . We shall obtain a contradiction by showing that  $[C]$  has an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path. According to the locations of  $x$  in  $L_1[x_2, x_{a_1}] \cup L_2(x_{a_2}, x_q) \cup L_3(x_{a_3}, x_r)$  and  $y$  on  $L_1 \cup L_2 \cup L_3$ , there are nine cases to check, which are very similar in the verification. So we just show one example with  $x$  in  $L_3(x_{a_3}, x_r]$  and  $y$  on  $L_1(x_{a_1}, x_{p-1}]$ . In this case,

$$x_2 C[x_2, y^-] x_p C^- [x_p, y] x C[x, x_1] x^- C^- [x^-, x_{p+1}]$$

is an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path of  $[C]$ , a contradiction.

Similarly,  $N(y, C) \subseteq N(x_p, C) \cup \{x_p\}$  for each vertex  $y$  of  $L_1(x_{a_1}, x_{p-1}) \cup L_2[x_{p+1}, x_{a_2}] \cup L_3(x_{q+1}, x_{a_3})$ . As  $d(x, C) \geq (r+1)/2$  for all  $x \in V(C) - V(e)$ , we see that  $r$  is odd and  $d(x_1, C) = d(x_p, C) = (r+1)/2$ . Furthermore, if  $\{x_{a_2}, x_{a_3}\} = \{x_q, x_{q+1}\}$ , then  $\{x_{a_1}, x_q, x_{q+1}\}$  is a vertex-cut of  $[C]$  and each component of  $[C] - \{x_{a_1}, x_q, x_{q+1}\}$  is isomorphic to  $K_{(r-3)/2}$ . Consequently, (d) holds. So assume that  $\{x_{a_2}, x_{a_3}\} \neq \{x_q, x_{q+1}\}$ . We shall obtain a contradiction by showing that  $[C]$  has an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path. If  $x_{q+1} \neq x_{a_3}$ . Then

$$x_2 C[x_2, x_{a_1}] x_1 C^- [x_1, x_{q+2}] x_{a_1+1} C[x_{a_1+1}, x_p] x_{q+1} C^- [x_{q+1}, x_{p+1}]$$

is an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path of  $[C]$ , a contradiction. Therefore  $x_{q+1} = x_{a_3}$  and  $x_q \neq x_{a_2}$ . Then

$$x_2 C[x_2, x_p] x_{q+1} x_q x_1 C^- [x_1, x_{q+1}^+] x_{q-1} C^- [x_{q-1}, x_{p+1}]$$

is an  $x_2$ - $x_{p+1}$   $e$ -hamiltonian path of  $[C]$ , a contradiction. This proves the lemma.  $\blacksquare$

**Lemma 3.7** *Let  $C$  be a cycle of order  $r$  in  $G$ . Let  $\lambda$  be a positive integer. Let  $e \in E(C)$ . Suppose that  $d^*(P, C) \geq r + \lambda$  for every  $e$ -hamiltonian path  $P$  of  $[C]$ . Then  $d(xy, C) \geq r + \lambda$  for every pair  $x$  and  $y$  of distinct vertices of  $C$  with  $V(e) \neq \{x, y\}$ .*

**Proof.** On the contrary, say that there are two distinct vertices  $x$  and  $y$  on  $C$  with  $V(e) \neq \{x, y\}$  such that  $d(xy, C) \leq r + \lambda - 1$ . Clearly, either  $e \notin \{xx^-, yy^-\}$  or  $e \notin \{xx^+, yy^+\}$ . Say w.l.o.g. the former holds. Then  $d(xx^-, C) \geq r + \lambda$  and  $d(yy^-, C) \geq r + \lambda$ . Thus  $d(x^-y^-, C) \geq 2(r + \lambda) - (r + \lambda - 1) \geq r + 2$ . By Lemma 3.6,  $[C]$  has an  $x$ - $y$   $e$ -hamiltonian path and therefore  $d(xy, C) \geq r + \lambda$ , a contradiction.  $\blacksquare$

**Lemma 3.8** *Let  $C = x_1 \dots x_r x_1$  be a cycle in  $G$ . Let  $e = x_1 x_2$ . Suppose that  $d^*(P, C) \geq r + 1$  for each  $e$ -hamiltonian path  $P$  of  $[C]$  with  $\sigma(P, e) > 0$ . If there exists  $x_j \in V(C) - V(e)$  such that  $d(x_j, C) \leq r/2$  then one of the following two statement holds:*

- (a) *If  $4 \leq j \leq r - 1$  then  $d(x_i, C) \geq (r + 2)/2$  for all  $3 \leq i \leq r$  with  $i \neq j$ ;*
- (b) *If  $j \in \{3, r\}$  then  $d(x_i, C) \geq (r + 2)/2$  for all  $4 \leq i \leq r - 1$ .*

**Proof.** To prove (a), say  $4 \leq j \leq r - 1$ . Then  $d(x_{j-1}, C) \geq r + 1 - d(x_j, C) \geq (r + 2)/2$ . Similarly,  $d(x_{j+1}, C) \geq (r + 2)/2$ . If  $d(x_i, C) \leq (r + 1)/2$  for some  $3 \leq i \leq r$  with  $i \neq j$ , let  $x_i$  be the one closest to  $x_j$  on  $C - e$ . Say w.l.o.g.  $i > j$ . Then  $d(x_{i-1}, C) \geq (r + 2)/2$ . Thus  $d(x_{j-1}x_{i-1}, C) \geq r + 2$ . By Lemma 3.6,  $[C]$  has an  $x_j$ - $x_i$   $e$ -hamiltonian path and so  $d(x_i x_j, C) \geq r + 1$ . Thus  $d(x_i, C) \geq r + 1 - r/2 = (r + 2)/2$ , a contradiction.

To prove (b), say w.l.o.g. that  $d(x_3, C) \leq r/2$ , i.e.,  $d(x_3, C) \leq \lfloor r/2 \rfloor$ . If  $r \leq 4$ , nothing to prove. So assume  $r \geq 5$ . Then  $d(x_4, C) \geq r + 1 - \lfloor r/2 \rfloor = \lceil (r + 2)/2 \rceil$ . Similarly, if  $d(x_r, C) \leq r/2$  then  $d(x_{r-1}, C) \geq \lceil (r + 2)/2 \rceil$  and so  $d(x_4 x_{r-1}, C) \geq r + 2$ . If  $d(x_r, C) \not\leq r/2$ , i.e.,  $d(x_r, C) \geq \lceil (r + 1)/2 \rceil$ , then  $d(x_4 x_r, C) \geq \lceil (r + 2)/2 \rceil + \lceil (r + 1)/2 \rceil = r + 2$ . Let  $s \in \{r - 1, r\}$  be maximal such that  $d(x_4 x_s, C) \geq r + 2$ . If  $d(x_i, C) \leq (r + 1)/2$  for some  $i \in \{5, \dots, r - 1\}$ , let  $x_i$  be the one closest to  $x_s$  on  $C - e$ . Then  $d(x_4 x_{i+1}, C) \geq r + 2$ . By Lemma 3.6,  $[C]$  has an  $x_3$ - $x_i$   $e$ -hamiltonian path and so  $d(x_i, C) \geq r + 1 - r/2 = (r + 2)/2$ , a contradiction.  $\blacksquare$

**Lemma 3.9** [6] *Let  $P = x_t x_{t-1} \dots x_1$  be an optimal path at  $x_t$  in  $G$ . Let  $r = \alpha(P, x_1)$  and  $c > r/2$ . Suppose that for each  $v \in V(G)$ , if there exists a longest path starting at  $x_t$  in  $G$  such that the path ends at  $v$  then  $d(v) \geq c$ . Then  $N(x_i) \subseteq \{x_1, x_2, \dots, x_r\}$ ,  $[P]$  has an  $x_t$ - $x_i$  hamiltonian path and  $d(x_i) \geq c$  for all  $i \in \{1, 2, \dots, r-1\}$ . Moreover, if  $t > r$  then  $x_r$  is a cut-vertex of  $G$ .*

**Lemma 3.10** *Let  $P = x_t x_{t-1} \dots x_1$  be an optimal path at  $x_t$  in  $G$ . Let  $r = \alpha(P, x_1)$ . Suppose that  $r \geq 3$  and for each  $v \in V(G)$ , if there exists a longest path starting at  $x_t$  in  $G$  such that the path ends at  $v$  then  $d(v) \geq (r+2)/2$ . Then for each pair  $x_i$  and  $x_j$  of distinct vertices in  $\{x_1, x_2, \dots, x_{r-1}\}$ , the following three statements hold:*

- (a) *If  $d(x_r, \{x_1, x_2, \dots, x_{r-1}\}) \geq 3$  then  $[P] - x_i$  has an  $x_t$ - $x_j$  hamiltonian path;*
- (b) *If  $N(x_r, \{x_1, x_2, \dots, x_{r-1}\}) = \{x_1, x_{r-1}\}$  but  $i \notin \{1, r-1\}$  then  $[P] - x_i$  has an  $x_t$ - $x_j$  hamiltonian path;*
- (c) *If  $N(x_r, \{x_1, x_2, \dots, x_{r-1}\}) = \{x_1, x_{r-1}\}$  and  $i \in \{1, r-1\}$  but  $j \notin \{1, r-1\}$  then  $[P] - x_i$  has an  $x_t$ - $x_j$  hamiltonian path.*

**Proof.** Obviously, the lemma is true if  $r \leq 4$ . So assume  $r \geq 5$ . Let  $H = [\{x_1, \dots, x_r\} - \{x_i\}]$ . By Lemma 3.9, for each  $x_l \in \{x_1, \dots, x_{r-1}\}$ ,  $[P]$  has an  $x_t$ - $x_l$  hamiltonian path,  $N(x_l, G) \subseteq V(H) \cup \{x_i\}$  and  $d(x_l, H + x_i) \geq (r+2)/2$ . Moreover,  $x_r$  is a cut-vertex of  $[P]$  if  $t > r$ , and consequently,  $H + x_i$  has an  $x_r$ - $x_i$  hamiltonian path and so  $H$  has a hamiltonian path starting at  $x_r$ . Obviously, for each  $v \in V(H - x_r)$ ,  $d(v, H) \geq (r+2)/2 - 1 = ((r-1) + 1)/2$ . Let  $L$  be an optimal path at  $x_r$  in  $H$ . Say  $L$  is an  $x_r$ - $y$  path. Then  $\alpha(L, y) \leq r-1$ . As  $\delta(H - x_r) \geq (r+2)/2 - 2 = (r-2)/2$ ,  $H - x_r$  is hamiltonian. If  $d(x_r, H) \geq 2$ , then  $H$  is 2-connected and by applying Lemma 3.9 to  $L$  in  $H$ , we see that  $\alpha(L, y) = r-1$ . Consequently,  $H$  has an  $x_r$ - $x_j$  hamiltonian path and so  $[P - x_i]$  has an  $x_t$ - $x_j$  hamiltonian path. Therefore (a) and (b) hold. If  $d(x_r, H) = 1$ , then  $x_i \in \{x_1, x_{r-1}\}$  and so  $\alpha(L, y) = r-2$ . Moreover, the vertex  $z$  with  $\{x_i, z\} = \{x_1, x_{r-1}\}$  is a cut-vertex of  $H$ . To see (c), we have  $x_j \notin \{x_1, x_{r-1}\}$  and  $H$  has an  $x_r$ - $x_j$  hamiltonian path. ■

## 4 Proof of Lemmas 2.1-2.7

**Proof of Lemma 2.1.** On the contrary, say  $d(x_1, G_1) \geq (n_1 + 2)/2$  and  $d(y_1, G_2) \geq (n_2 + 2)/2$ , i.e.,  $\delta_1(G_1) \geq (n_1 + 2)/2$  and  $\delta_2(G_2) \geq (n_2 + 2)/2$ . Say w.l.o.g.  $G_1 \notin \mathcal{H}_1$ . By Lemma 3.3, we see that  $\mathcal{P}_1^*(G_1) = \emptyset$ . Let  $P = u_{n_1} u_{n_1-1} \dots u_1$  be an optimal path at  $e_1 = u_{n_1} u_{n_1-1}$  in  $G_1$ . Then  $N(u_{n_1}, G_1) \subseteq \{u_{n_1-1}, u_{n_1-2}\}$ . Say  $\alpha(P, u_1) = r$ . As  $\delta_1(G_1) \geq (n_1 + 2)/2$  and by Lemma 3.9,  $u_1 \dots u_r u_1$  is an end-cycle at  $u_r$  in  $G_1$  and for each  $j \in \{1, \dots, r-1\}$ ,  $G_1$  has a  $u_{n_1}$ - $u_j$   $e_1$ -hamiltonian path and  $d(u_j, G_1) \geq (n_1 + 2)/2$ . Since  $n_2 \geq d(u_{n_1}, G_2) \geq (n+4)/2 - d(u_{n_1}, G_1) \geq n/2$ , we obtain  $n_2 \geq n_1$ . Note that  $r-1 \geq (n_1 + 2)/2$  and so  $n_1 \geq 6$ .

By Property B,  $\mathcal{P}_2^*(G_2) \neq \emptyset$ . As  $\delta_2(G_2) \geq (n_2 + 2)/2$  and by Lemma 3.3,  $G_2 \in \mathcal{H}_2$ . Thus  $d(y, G_2) \geq (n_2 + 2)/2$  for all  $y \in V(G_2) - V(e_2)$ . Let  $v_1 \dots v_{n_2} v_1$  be a hamiltonian cycle of  $G_2$  with  $e_2 = v_1 v_2$ . Let  $i, j \in \{1, \dots, r-1\}$  with  $i \notin \{1, r-1\}$ . By Lemma 3.10,  $G_1 - u_i$  has an  $u_{n_1}$ - $u_j$   $e_1$ -hamiltonian path. Clearly,  $d(u_{n_1} u_j, G_2) \geq n + 4 - (n_1 - 1) = n_2 + 5$ . Thus for some  $s \in \{4, \dots, n_2 - 1\}$ ,  $d(v_s, u_{n_1} u_j) = 2$  and so  $G_1 - u_i + v_s \in \mathcal{H}_1$ . Thus  $G_2 - v_s + u_i \notin \mathcal{H}_2$ . As  $d(v_{s-1} v_{s+1}, G_2 - v_s) \geq n_2 + 2 - 2 = n_2$  and by Lemma 3.3,  $G_2 - v_s \in \mathcal{H}_2$ . Let  $C = w_1 \dots w_t w_1$  be an  $e_2$ -hamiltonian cycle of  $G_2 - v_s$  with  $t = n_2 - 1$ . As  $d(u_i, G_1) \leq n_1 - 2$ ,  $d(u_i, C) \geq (n+4)/2 - (n_1 - 2) - 1 \geq 3$ . As  $C + u_i \notin \mathcal{H}_2$ , we see that there are two distinct vertices  $u$  and  $v$  in  $C$  such that  $\{u, v\} \cap V(e_2) = \emptyset$  and either  $\{u^+, v^+\} \subseteq N(u_i)$  or  $\{u^-, v^-\} \subseteq N(u_i)$ . Say w.l.o.g.  $\{u^+, v^+\} \subseteq N(u_i)$ . As  $G_2 - v_s + u_i \notin \mathcal{H}_2$ ,  $[C]$  does not have a  $u^+ v^+$   $e_2$ -hamiltonian path. Clearly,  $d(x, C) \geq (n_2 + 2)/2 - 1 = (t+1)/2$  for all  $x \in V(C) - V(e_2)$ . Thus we may apply Lemma 3.6(d) to  $[C]$ . First, assume that  $[C]$  has a vertex-cut  $X$  with  $|X| = 3$  and  $V(e_2) \subseteq X$  such that each of the two components  $[C] - X$  is isomorphic to  $K_{(t-3)/2}$ . As  $G_2 - v_s + u_i \notin \mathcal{H}_2$ , we see that  $N(u_i, C) = X$ . Thus  $v_s x \in E$  for all  $x \in V(C) - X$  as  $\delta_2(G_2) \geq (n_2 + 2)/2$ . Let  $v' \in I(u_{n_1} u_j, C - X)$ . Then  $G_1 - u_i + v' \in \mathcal{H}_1$  and  $G_2 - v' + u_i \in \mathcal{H}_2$  by Lemma 3.10, a contradiction. Therefore  $V(C)$  has a partition  $X \cup Y$  such that  $|X| = (t+1)/2$ ,  $V(e_2) \subseteq X$ ,  $|Y| = (t-1)/2$ ,  $\{u, v\} \subseteq Y$  and  $N(y, C) = X$  for all  $y \in Y$ . As  $\delta_2(G_2) \geq (n_2 + 2)/2$ , we obtain  $Y \subseteq N(v_s)$ . As  $G_2 - v_s + u_i \notin \mathcal{H}_2$ , we see that  $N(u_i, C) \subseteq X$ . As  $d(u_{n_1}, G_1) \leq 2$ , we readily see that  $d(u_{n_1}, Y) > 0$ . Let  $v' \in N(u_{n_1}, Y)$ . Clearly,

$d(v', G_1) \geq (n+4)/2 - (n_2+2)/2 = (n_1+2)/2$ . Thus  $v'u_p \in E$  for some  $p \in \{1, \dots, r-1\}$  with  $p \neq i$ . By Lemma 3.10,  $G_1 - u_i$  has a  $u_{n_1}-u_p$   $e_1$ -hamiltonian path. With  $v'$  and  $u_p$  in place of  $v_s$  and  $u_j$  in the above argument, we see that  $V(G_2 - v')$  has a partition  $X' \cup Y'$  such that  $|X'| = (t+1)/2$ ,  $V(e_2) \subseteq X'$ ,  $|Y'| = (t-1)/2$ ,  $N(y, G_2 - v') = X'$  for all  $y \in Y'$ ,  $Y' \subseteq N(v')$  and  $N(u_i, G_2 - v') \subseteq X'$ . Since  $Y' \neq Y$  and  $Y$  is an independent set, we see that  $Y \subseteq X' \cup \{v'\}$ . Thus  $N(u_i, Y) \neq \emptyset$ , a contradiction.  $\blacksquare$

**Proof of Lemma 2.2.** On the contrary, say  $d(x_1, G_1) \leq (n_1+1)/2$  and  $d(y_1, G_2) \leq (n_2+1)/2$ . Then  $d(x_1, G_2) \geq (n_2+3)/2$  and  $d(y_1, G_1) \geq (n_1+3)/2$ . By Lemma 3.2(a),  $G_1 - x_1 + y_1 \in \mathcal{P}_1$  and  $G_2 - y_1 + x_1 \in \mathcal{P}_2$ . By Property A,  $\xi(x_1, y_1) \leq 0$ . This implies that  $d(x_1, G_1) = (n_1+1)/2$ ,  $d(x_1, G_2) = (n_2+3)/2$ ,  $d(y_1, G_2) = (n_2+1)/2$ ,  $d(y_1, G_1) = (n_1+3)/2$  and  $x_1 y_1 \in E$ . Since either  $G_1 \notin \mathcal{H}_1$  or  $G_2 \notin \mathcal{H}_2$ , say w.l.o.g.  $G_1 \notin \mathcal{H}_1$ . As  $\delta_1(G_1) = (n_1+1)/2$  and by Lemma 3.3,  $\mathcal{P}_1^*(G_1) = \emptyset$ . Therefore  $e_1 = x_{n_1} x_{n_1-1}$  and  $N(x_{n_1}, G_1) \subseteq \{x_{n_1-1}, x_{n_1-2}\}$ . Thus  $n_2 \geq d(x_{n_1}, G_2) \geq (n+4)/2 - 2$ . This implies  $n_2 \geq n_1$ .

By Property B,  $\mathcal{P}_2^*(G_2) \neq \emptyset$ . As  $\delta_2(G_2) = (n_2+1)/2$  and by Lemma 3.3,  $G_2 \in \mathcal{H}_2$ . Then  $d(y, G_2) \geq (n_2+1)/2$  for all  $y \in V(G_2) - V(e_2)$ . Let  $H_1 = G_1 - x_1$  and  $H_2 = G_2 + x_1$ . By Property A and Lemma 3.2(a) as above, we readily see that  $H_2 - y \in \mathcal{P}_2$ , if  $d(y, G_2) = (n_2+1)/2$  then  $yx_1 \in E$ , and so  $d(y, H_2) \geq (n_2+3)/2$  for all  $y \in V(H_2) - V(e_2)$ . Let  $C = v_1 v_2 \dots v_t v_1$  be a hamiltonian cycle of  $H_2$  with  $t = n_2 + 1$  and  $e_2 = v_1 v_2$ . Let  $Y$  be the set of those vertices  $y \in V(H_2) - V(e_2)$  such that  $H_2 - y \in \mathcal{H}_2$ . Then  $H_1 + y \notin \mathcal{H}_1$  for all  $y \in Y$ . For each  $v_s \in V(C) - \{v_1, v_2, v_3, v_t\}$ ,  $d(v_{s-1} v_{s+1}, C - v_s) \geq n_2 + 3 - 2 = n_2 + 1$  and so  $H_2 - v_s \in \mathcal{H}_2$  by Lemma 3.3. Thus  $V(C) - \{v_1, v_2, v_3, v_t\} \subseteq Y$ . Since  $\mathcal{P}_1^*(G_1) = \emptyset$  and  $N(x_{n_1}, G_1) \subseteq \{x_{n_1-1}, x_{n_1-2}\}$ , we see that  $d(x_2 x_{n_1}, H_1) \leq n_1 - 2$ . It follows that  $d(x_2 x_{n_1}, H_2) \geq n + 4 - (n_1 - 2) = t + 5$ . Consequently,  $v_s \in I(x_2 x_{n_1}, H_2)$  for some  $v_s \in V(C) - \{v_1, v_2, v_3, v_t\}$  and so  $H_1 + v_s \in \mathcal{H}_1$ , a contradiction.  $\blacksquare$

**Proof of Lemma 2.3.** On the contrary, say that  $G_2 \in \mathcal{H}_2$ . Then  $y \in S_2(G_2)$  and so  $d(y, G_2) \geq (n_2+2)/2$  for all  $y \in V(G_2) - V(e_2)$  and  $G_1 \notin \mathcal{H}_1$ . As  $d(x_1, G_2) \geq (n_2+3)/2$ ,  $G_2 + x_1 \in \mathcal{H}_2$  by Lemma 3.2(f) and so  $S_2(G_2 + x_1) = V(G_2 + x_1) - V(e_2)$ . By Property A and Lemma 3.2(a), we readily see that  $d(y, G_2 + x_1) \geq (n_2+3)/2$  for all  $y \in V(G_2) - V(e_1)$ . Set  $H_1 = G_1 - x_1$  and  $H_2 = G_2 + x_1$ . Let  $A = \{v \in V(H_2) - V(e_2) \mid H_2 - v \in \mathcal{H}_2\}$ . Then  $H_1 + v \notin \mathcal{H}_1$  for each  $v \in A$ . Let  $C = v_1 v_2 \dots v_{n_2} v_1$  be a hamiltonian cycle of  $G_2$  with  $e_1 = v_1 v_2$ . Say  $X_0 = \{v_{n_2}, v_1, v_2, v_3\}$ . We claim

*Claim 1* The following two statements hold:

- (a)  $V(H_2) - X_0 \subseteq A$ ;
- (b) If  $d(v_1, H_2 - X_0) \geq 1$  then  $v_{n_2} \in A$  and if  $d(v_2, H_2 - X_0) \geq 1$  then  $v_3 \in A$ .

*Proof of Claim 1.* Clearly,  $x_1 \in A$ . Let  $v_i \in V(G_2) - X_0$ . Then  $d(v_{i-1} v_{i+1}, G_2 - v_i) \geq (n_2+2) - 2 = (n_2-1) + 1$  and by Lemma 3.3,  $G_2 - v_i \in \mathcal{H}_2$ . Since  $d(x_1, G_2 - v_i) \geq (n_2+3)/2 - 1 = ((n_2-1) + 2)/2$ ,  $H_2 - v_i \in \mathcal{H}_2$ . Hence (a) holds.

To see (b), we just need show the first assertion by the symmetry. If  $x_1 v_1 \in E$  then  $x_1 v_1 \dots v_{n_2-1} \in \mathcal{P}_2(H_2 - v_{n_2})$  and  $d(x_1 v_{n_2-1}, H_2 - v_{n_2}) \geq n_2 + 3 - 2 = n_2 + 1$ . By Lemma 3.3,  $H_2 - v_{n_2} \in \mathcal{H}_2$ . If  $v_1 v_i \in E$  for some  $v_i \in V(G_2) - X_0$ , then  $v_{i-1} v_{i-2} \dots v_2 v_1 v_i v_{i+1} \dots v_{n_2-1} \in \mathcal{P}_2(G_2 - v_{n_2})$  and  $d(v_{i-1} v_{n_2-1}, G_2 - v_{n_2}) \geq n_2$ . As above, we see  $H_2 - v_{n_2} \in \mathcal{H}_2$ . Hence (b) holds.  $\square$

We now divide the proof of the lemma into the following two cases. Say  $l = n_1 - 1$ .

Case 1.  $H_1 \notin \mathcal{H}_1$ .

Let  $P = z_1 \dots z_l$  be an arbitrary path in  $\mathcal{P}_1(H_1)$ . Then  $I(z_1 z_l, A) = \emptyset$ . Thus  $d(z_1 z_l, H_2) \leq n_2 + 5$  and so  $d(z_1 z_l, H_1) \geq l$ . By Lemma 3.3,  $d(z_1 z_l, H_1) = l$  and  $\sigma(P, e_1) > 0$ . Thus  $d(z_1 z_l, H_2) = n_2 + 5$ ,  $X_0 = I(z_1 z_l, H_2)$ ,  $A = V(H_2) - X_0$  and  $d(x, z_1 z_l) = 1$  for all  $x \in A$ . By Claim 1,  $N(v_1 v_2, H_2) \subseteq X_0$ . Then  $n_1 - 1 = l \geq d(v_1, H_1) \geq (n_1 + n_2 + 4)/2 - d(v_1, G_2) \geq (n_1 + n_2 + 4)/2 - 3$  and  $d(x_1, G_2) \leq (n_2 - 2)$ . As  $d(x_1, G_2) \geq (n_2 + 3)/2$ , we see that  $n_2 \geq 7$ . As  $n_2 - 3 \geq d(v_5, G_2) \geq (n_2 + 2)/2$ , it follows that  $n_1 \geq n_2 \geq 8$  and  $d(x_1, H_1) \geq 4$ .

We apply Lemma 3.5 to  $H_1$ . First, assume that  $V(H_1)$  has a partition  $X \cup Y$  such that  $|X| = l/2$ ,  $V(e_1) \subseteq X$  and  $N(y, H_1) = X$  for all  $y \in Y$ . Then every two distinct vertices in  $Y$  can play the role of  $z_1$  and  $z_l$ . Hence  $d(x_1, Y) \geq l/2 - 1 \geq 2$  and so  $G_1 \in \mathcal{H}_1$ , a contradiction. Therefore  $H_1 - V(e_1)$  has two components  $J_1$  and  $J_2$  such that  $H_1 - V(e_1) = J_1 \cup J_2$ , each of  $J_1$  and  $J_2$  is complete and  $d(x, H_1) = l - 1$  for each  $x \in V(e_1)$ .

Say w.l.o.g.  $z_1 \in V(J_1)$  and  $d(z_1, H_1) \leq d(z_l, H_1)$ . Then  $d(z_1, G_1) \leq (n_1 + 1)/2$  and so  $d(z_1, G_2) \geq (n_2 + 3)/2$ . Clearly,  $G_1 - z_1 \in \mathcal{P}_1$  and  $G_1 - z_1$  has an  $x_1 - z_l$  hamiltonian  $e_1$ -path. Switching the roles of  $z_1$  and  $x_1$  in the above argument, we also obtain  $X_0 = I(x_1 u, G_2 + z_1)$ . By Claim 1,  $\{v_3, v_{n_2}\} \subseteq A$ , a contradiction.

Case 2.  $H_1 \in \mathcal{H}_1$ .

Let  $L = u_1 u_2 \dots u_l u_1$  be a hamiltonian cycle of  $H_1$  with  $e_1 = u_1 u_2$ ,  $B = V(L - u_1)$  and  $a = n_2 + 1 - |A|$ . If  $a \geq 3$  then  $N(v_1, H_2) \subseteq X_0$  or  $N(v_2, H_2) \subseteq X_0$  by Claim 1. As  $\delta_2(G_2) \geq (n_2 + 2)/2$ , it follows that  $n_2 \geq 6$  if  $a \geq 3$ . We divide this case into the following three subcases.

Subcase 2.1.  $d^*(P, H_1) \geq l + 2$  for all  $P \in \mathcal{P}_1(H_1)$ .

By Lemma 3.7,  $d(xy, H_1) \geq l + 2$  for all  $x, y \in V(H_1)$  with  $x \neq y$  and  $xy \neq e_1$ . By Lemma 3.6, for all  $x, y \in V(H_1)$  with  $x \neq y$  and  $xy \neq e_1$ ,  $H_1$  has an  $x$ - $y$   $e_1$ -hamiltonian path. Since  $H_1 + v_i \notin \mathcal{H}_1$  for all  $v_i \in A$ , we see that the following Claim 2 holds:

*Claim 2* For each  $v_i \in A$ , if  $d(v_i, H_1) \geq 2$  then  $N(v_i, H_1) = V(e_1)$ . □

By Claim 2,  $n_2 \geq (n_1 + n_2 + 4)/2 - d(v_i, H_1) \geq (n_1 + n_2 + 4)/2 - 2$  for all  $v_i \in A$ . Thus  $n_2 \geq n_1$ . By Claim 2,  $d(v_i, B) \leq 1$  for all  $v_i \in A$  and so  $d(A, B) \leq |A| = n_2 + 1 - a$ . On the other hand,  $d(A, B) \geq \sum_{u \in B} d(u, A) \geq \sum_{u \in B} ((n_1 + n_2 + 4)/2 - d(u, H_1) - a) \geq (n_1 - 2)((n_1 + n_2 + 4)/2 - (n_1 - 2) - a)$ . Therefore  $(n_1 - 2)((n_1 + n_2 + 4)/2 - (n_1 - 2) - a) - (n_2 + 1 - a) \leq 0$ . Denote the left side of this inequality by  $f(n_1)/2$  with  $n_2 = n - n_1$ . Then  $f(n_1) = -2n_1^2 + (n + 14 - 2a)n_1 + (-4n - 18 + 6a) \leq 0$  for  $4 \leq n_1 \leq n/2$ . As  $f''(n_1) < 0$ ,  $f(n_1) \geq \min\{f(4), f(n/2)\} = \min\{6 - 2a, 3n - an - 18 + 6a\}$ . Thus  $a \geq 3$  for otherwise  $f(n_1) > 0$ . Thus  $N(v_1, H_2) \subseteq X_0$  or  $N(v_2, H_2) \subseteq X_0$ . Say w.l.o.g.  $N(v_1, H_2) \subseteq X_0$ . Then  $n_1 - 1 \geq d(v_1, H_1) \geq (n_1 + n_2 + 4)/2 - 3$  which implies that  $n_1 \geq n_2$ . Let  $v_i \in A - X_0$ . Then  $n_2 - 1 \geq d(v_i, H_2) \geq (n_1 + n_2 + 4)/2 - 2$  which implies that  $n_2 \geq n_1 + 2$ , a contradiction.

Subcase 2.2.  $d^*(P, H_1) \geq l + 1$  for all  $P \in \mathcal{P}_1(H_1)$ .

By the above subcase,  $d^*(P, H_1) = l + 1$  for some  $P \in \mathcal{P}_1(H_1)$ . Thus  $d^*(P, H_2) \geq n_1 + n_2 + 4 - l - 1 = n_2 + 4$ . As  $d^*(P, v_i) \leq 1$  for all  $v_i \in A$ . Thus  $d^*(P, v') = 2$  and so  $v' \notin A$  for some  $v' \in \{v_3, v_{n_2}\}$ . It follows that  $a \geq 3$  and so  $n_2 \geq 6$ . By Claim 1,  $N(v_1, H_2) \subseteq X_0$  and we may assume that  $v_{n_2} \notin A$ . As in the above paragraph, this implies that  $n_1 \geq n_2$ . Let  $z$  be an arbitrary vertex in  $A - X_0$ . Then  $n_1 - 1 \geq d(z, H_1) \geq (n_1 + n_2 + 4)/2 - (n_2 - 1) \geq 3$ . It is easy to see that there exist two distinct vertices  $u$  and  $w$  on  $L$  such that either  $\{u^-, w^-\} \subseteq N(z)$  and  $e_1 \notin \{uu^-, ww^-\}$  or  $\{u^+, w^+\} \subseteq N(z)$  and  $e_1 \notin \{uu^+, ww^+\}$ . Say w.l.o.g.  $\{u^+, w^+\} \subseteq N(z)$  and  $e_1 \notin \{uu^+, ww^+\}$ . By Lemma 3.7,  $d(xy, H_1) \geq l + 1$  for all  $\{x, y\} \subseteq V(H_1)$  with  $x \neq y$  and  $xy \neq e_1$ . We claim that  $d(x, H_1) \geq (l + 1)/2$  for all  $x \in V(H_1)$ . If this is false, say  $d(x_0, H_1) \leq l/2$  for some  $x_0 \in V(H_1)$ . Then  $d(x, H_1) \geq (l + 2)/2$  for all  $x \in V(H_1 - x_0)$  with  $x_0 x \neq e_1$  and  $d(x_0, H_2) \geq (n_1 + n_2 + 4)/2 - l/2 \geq (n_2 + 5)/2 \geq 5$ . Thus  $d(x_0, A - X_0) > 0$ . It is easy to see that in the choices of the vertices  $u$ ,  $w$  and  $z$  in the above, we can choose  $u$ ,  $w$  and  $z$  such that  $x_0 \notin \{u, w\}$ . Thus  $d(uw, H_1) \geq l + 2$  and by Lemma 3.6,  $H_1$  has a  $u^+ - w^+$   $e_1$ -hamiltonian path and so  $H_1 + z \in \mathcal{H}_1$ , a contradiction. Hence  $d(x, H_1) \geq (l + 1)/2$  for all  $x \in V(H_1)$ .

We now apply Lemma 3.6(d) to  $H_1$  since  $H_1$  does not have a  $u^+ - w^+$   $e_1$ -hamiltonian path. First, assume that  $H_1$  has a vertex-cut  $X$  with  $|X| = 3$  and  $V(e_1) \subseteq X$  such that  $H_1 - X = H'_1 \cup H''_1$  where  $H'_1$  and  $H''_1$  are isomorphic to  $K_{(l-3)/2}$ . Then  $N(z, H_1) = X$  as  $H_1 + z \notin \mathcal{H}_1$ . As  $z$  is arbitrary in  $A - X_0$ ,  $N(A - X_0, H_1) = X$ . It follows that  $d(x, G) \leq (l + 1)/2 + 4 < (n_1 + n_2 + 4)/2$  for  $x \in V(H_1 - X)$ , a contradiction. Therefore  $V(H_1)$  has a partition  $X \cup Y$  such that  $|X| = (l + 1)/2$ ,  $V(e_1) \subseteq X$ ,  $\{u, w\} \subseteq Y$ , and  $N(y, H_1) = X$  for all  $y \in Y$ . Clearly,  $\{u^+, w^+\} \subseteq X$ . Thus  $N(z, H_1) \subseteq X$  as  $H_1 + z \notin \mathcal{H}_1$ . Let  $y \in Y$ . As  $d(y, A - X_0) \geq (n_1 + n_2 + 4)/2 - (l + 1)/2 - 4 > 0$ , let  $z' \in N(y, A - X_0)$ . With  $z'$  in place of  $z$  in this argument, we see that  $V(H_1)$  has a partition  $X' \cup Y'$  such that  $|X'| = (l + 1)/2$ ,  $V(e_1) \subseteq X'$ ,  $N(y', H_1) = X'$  for all  $y' \in Y'$  and  $N(z', H_1) \subseteq X'$ . It follows that  $Y' \cap X \neq \emptyset$  and so  $Y' \subseteq X$ . Thus  $|X| \geq (l + 1)/2 + 1 = (l + 3)/2$ , a contradiction.

Subcase 2.3. For some  $P \in \mathcal{P}_1(H_1)$ ,  $d^*(P, H_1) \leq l$ .

For each  $P \in \mathcal{P}_1(H_1)$ , as  $d^*(P, A) \leq |A|$ ,  $d^*(P, H_1) \geq n_1 + n_2 + 4 - (n_2 + 1 + a) = l + 4 - a \geq l$ . Thus  $a = 4$  and by Claim 1,  $N(v, H_2) \subseteq X_0$  for  $v \in \{v_1, v_2\}$ . As before, it follows that  $n_1 \geq n_2 \geq 6$ . Let  $z$  be an arbitrary

vertex in  $A$ . Then  $d(z, H_1) \geq (n_1 + n_2 + 4)/2 - (n_2 - 2) \geq 4$ .

First, assume that there exists  $P \in \mathcal{P}_1^*(H_1)$  such that  $d^*(P, H_1) = l$ . As  $H_1 + v \notin \mathcal{H}_1$  for all  $v \in A$ , it follows that  $d^*(P, v) = 1$  for all  $v \in A$  and  $d^*(P, X_0) = 8$ . Say  $P = z_1 z_2 \dots z_l$  with  $d(z_1, P) \leq d(z_l, P)$ . Then  $d(z_1, P) \leq l/2$ ,  $d(z_1, H_2) \geq \lceil (n_1 + n_2 + 4)/2 \rceil - \lfloor l/2 \rfloor \geq 5$ . Let  $z_c \in \{z_1, z_l\}$  and  $v_b \in A$  be such that  $v_b z_c \in E$ . We claim that  $G_2 + z_c - v_j \in \mathcal{H}_2$  for all  $v_j \in V(G_2) - V(e_2)$ . To see this, say  $G_2 + z_c - v_j \notin \mathcal{H}_2$  for some  $v_j \in V(G_2) - V(e_2)$ . Clearly,  $v_{j-1} v_{j+1} \notin E$  otherwise  $G_2 + z_l - v_j \in \mathcal{H}_2$ . First assume that  $v_j \notin \{v_3, v_{n_2}\}$ . Then  $d(v_{j-1} v_{j+1}, G_2 - v_j) \geq n_2 + 2 - 2 = (n_2 - 1) + 1$ . This implies that  $C - v_j$  has an accessible edge  $e'$  with  $e' \neq e_2$ . Since  $N(v_1 v_2, G_2) \subseteq X_0$  and  $d(z_c, X_0) = 4$ , it follows that  $G_2 - v_j + z_c \in \mathcal{H}_2$ , a contradiction. Hence  $v_j \in \{v_3, v_{n_2}\}$ . Say w.l.o.g.  $v_j = v_3$ . Then  $P' = v_4 \dots v_b z_c v_2 v_1 v_{n_2} v_{n_2-1} \dots v_{b+1}$  is an  $e_2$ -hamiltonian path of  $G_2 - v_3 + z_c$  with  $d(v_4 v_{b+1}, G_2 - v_3 + z_c) \geq n_2$ . As  $d(v_4, e_2) = 0$ , this implies that  $P'$  has an accessible edge  $e''$  with  $e'' \neq e_2$  and so  $G_2 - v_j + z_c \in \mathcal{H}_2$ , a contradiction. Hence this claim holds. Let  $H'_1 = G_1 - z_c$  and  $H'_2 = G_2 + z_c$ . We claim that  $H'_1 \notin \mathcal{P}_1$ . To see this, say  $H'_1 \in \mathcal{P}_1$ . Then for any  $Q \in \mathcal{P}_1(H'_1)$  and  $v \in V(H'_2) - V(e_2)$ ,  $H'_1 + v \notin \mathcal{H}_1$  and so  $d^*(Q, v) \leq 1$ . Thus for any  $Q \in \mathcal{P}_1(H'_1)$ ,  $d^*(Q, H'_2) \leq n_2 + 3$  and so  $d^*(Q, H'_1) \geq l + 2$ . Let  $v_j \in A - \{x_1\}$ . Then  $d(v_j, H'_1) \geq (n_1 + n_2 + 4)/2 - d(v_j, G_2) - d(v_j, z_c) \geq (n_1 + n_2 + 4)/2 - (n_2 - 3) - 1 \geq 4$ . By Lemma 3.6 and Lemma 3.7, we see that  $H'_1 + v_j \in \mathcal{H}_1$ , a contradiction.

Therefore  $H'_1 \notin \mathcal{P}_1$ . As  $d(z_1, H_1) \leq \lfloor l/2 \rfloor$ ,  $d(z_1, G_2) \geq \lceil (n_1 + n_2 + 4)/2 \rceil - \lfloor l/2 \rfloor - 1 \geq 5$ . The above argument implies that  $H_1 - z_1 + x_1 \notin \mathcal{P}_1$  and so  $x_1 z_l \notin E$ . Thus  $z_1 x_1 \in E$  and so  $H_1 - z_l + x_1 \in \mathcal{P}_1$ . Consequently, the above argument implies that  $d(z_l, G_2) = d(z_l, X_0) = 4$ . Thus  $d(x_1 z_l, H_1 - z_1) \geq n_1 + n_2 + 4 - (n_2 - 2) - 4 - 2 = (l - 1) + 2$ . By Lemma 3.2,  $H_1 - z_1 + x_1 \in \mathcal{P}_1$ , a contradiction.

Therefore for each  $P \in \mathcal{P}_1^*(H_1)$ ,  $d^*(P, H_1) \geq l + 1$ . Recall that  $L = u_1 u_2 \dots u_l u_1$  is a hamiltonian cycle of  $H_1$  with  $e_1 = u_1 u_2$ . To apply Lemma 3.8, let us first assume that  $d(u_t, H_1) \leq l/2$  for some  $u_t \in V(L) - V(e_1)$ . If  $4 \leq t \leq l - 1$ , then  $d(u_j, H_1) \geq (l + 2)/2$  for all  $3 \leq j \leq l$  with  $j \neq t$ . As  $d(z, H_1) \geq 4$ , it is easy to see that there exist two distinct vertices  $u$  and  $w$  on  $L$  with  $u_t \notin \{u, w\}$  such that either  $\{u^-, w^-\} \subseteq N(z)$  and  $e_1 \notin \{u^- u, w^- w\}$  or  $\{u^+, w^+\} \subseteq N(z)$  and  $e_1 \notin \{u^+ u, w^+ w\}$ . By Lemma 3.6, we see that  $H_1 + z \in \mathcal{H}_1$ , a contradiction. Hence  $u_t \in \{u_3, u_l\}$ . By Lemma 3.8,  $d(u_j, H_1) \geq (l + 2)/2$  for all  $5 \leq j \leq l - 1$ . To avoid the existence of  $u$  and  $w$  as above such that  $H_1 + z \in \mathcal{H}_1$ , we see that  $N(z, H_1) = \{u_1, u_2, u_4, u_{l-1}\}$ . As  $z$  is an arbitrary vertex in  $A$ , we see that  $d(u_t, A) = 0$  and so  $d(u_t, H_2) \leq 5$ . Thus  $d(u_t, H_1) \geq (n_1 + n_2 + 4)/2 - 5 \geq (l + 1)/2$ , a contradiction.

Therefore  $d(u_j, H_1) \geq (l + 1)/2$  for all  $u_j \in V(H_1) - V(e_1)$ . As  $d(z, H_1) \geq 4$ , there exist two distinct vertices  $u$  and  $w$  on  $C$  such that either  $\{u^-, w^-\} \subseteq N(z)$  and  $e_1 \notin \{uu^-, ww^-\}$  or  $\{u^+, w^+\} \subseteq N(z)$  and  $e_1 \notin \{uu^+, ww^+\}$ . Say w.l.o.g.  $\{u^+, w^+\} \subseteq N(z)$  and  $e_1 \notin \{uu^+, ww^+\}$ . We now apply word by word the argument in the last paragraph of Subcase 2.2 to  $H_1$  and  $H_2$  and a contradiction follows.  $\blacksquare$

**Proof of Lemma 2.4.** As  $\mathcal{P}_2^*(G_2) = \emptyset$ ,  $N(v_{n_2}, G_2) \subseteq \{v_{n_2-1}, v_{n_2-2}\}$  and so  $n_1 \geq d(v_{n_2}, G_1) \geq (n_1 + n_2 + 4)/2 - d(v_{n_2}, G_2)$ . Thus  $n_1 \geq n_2$  and if  $n_1 = n_2$  then  $N(v_{n_2}, G_2) = \{v_{n_2-2}, v_{n_2-1}\}$  and so  $r \leq n_2 - 2$ . Since  $n_2 - 2 \geq r - 1 \geq (n_2 + 2)/2$ , we see that  $n_2 \geq 6$  and if  $r \leq n_2 - 2$  then  $n_2 \geq 8$ .

On the contrary, say that the lemma fails. Let  $u_0 \in V(G_1) - V(e_1)$  with  $d(u_0, G_1)$  minimal be such that  $G_1 - u_0 \in \mathcal{P}_1$ ,  $G_2 + u_0 \in \mathcal{H}_2$  and  $d(u_0, J^*) > 0$ . Let  $v_c \in J^*$  with  $u_0 v_c \in E$ . As  $G_2 + u_0 \in \mathcal{H}_2$ , we see that  $u_0 v_{n_2} \in E$  if  $v_{n_2} v_{n_2-2} \notin E$  and  $d(u_0, v_{n_2} v_{n_2-1}) \geq 1$  if  $v_{n_2} v_{n_2-2} \in E$ . Thus we may assume w.l.o.g. that  $u_0 v_{n_2} \in E$ . Let  $B$  be the set of all the vertices  $v_i$  in  $G_2$  such that  $G_2 - v_i + u_0 \in \mathcal{H}_2$ . By Lemma 3.10,  $V(J) - \{v_c, v_r\} \subseteq B$ , and if  $d(u_0, J - v_r) \geq 2$  then  $v_c \in B$ . Set  $H = G_1 - u_0$  and  $l = |H| = n_1 - 1$ . We claim the following:

*Claim A* If  $d(u_0, G_1) \leq (n_1 + 1)/2$  then  $r \in \{n_2 - 2, n_2 - 1\}$  and  $B = \{v_1, \dots, v_{r-1}\}$ . Moreover, for each  $P \in \mathcal{P}_1(H)$  we have that  $d^*(P, v_i) \leq 1$  for all  $1 \leq i \leq r - 1$ ,  $d^*(P, H) \geq l$  and if  $d^*(P, H) = l$  then  $r = n_2 - 2$ ,  $d^*(P, v_{n_2-2} v_{n_2-1} v_{n_2}) = 6$ ,  $d^*(P, u_0) = 2$  and  $d^*(P, v_i) = 1$  for all  $1 \leq i \leq r - 1$ .

*Proof of Claim A.* Say  $d(u_0, G_1) \leq (n_1 + 1)/2$ . Then  $d(u_0, G_2) \geq (n_2 + 3)/2$ . As  $G_2 + u_0 \in \mathcal{H}_2$ , for each  $y \in V(G_2) - V(e_2)$ ,  $G_2 - y + u_0 \in \mathcal{P}_2$  and so if  $G_1 - u_0 + y \in \mathcal{P}_1$  then  $\xi(u_0, y) \leq 0$  by Property A. Let  $y$  be an arbitrary vertex of  $G_2 - V(e_2)$ . If  $d(y, G_2) \leq (n_2 + 1)/2$  then  $d(y, G_1) \geq (n_1 + 3)/2$  and so  $G_1 - u_0 + y \in \mathcal{P}_1$  by Lemma 3.2(a). Consequently,  $\xi(u_0, y) \leq 0$ . This implies that  $d(y, G_2) = (n_2 + 1)/2$  and  $u_0 y \in E$ . Therefore  $d(y, G_2) \geq (n_2 + 1)/2$  for all  $y \in V(G_2) - V(e_2)$ . Consequently,  $r \in \{n_2 - 2, n_2 - 1\}$ . As  $d(u_0, G_2) \geq (n_2 + 3)/2$

and  $r - 1 \geq \lceil (n_2 + 2)/2 \rceil$ , we see that  $d(u_0, J - v_r) \geq \lceil (n_2 + 3)/2 \rceil - (n_2 - r) - 1 \geq 3$ . By Lemma 3.10,  $B = \{v_1, \dots, v_{r-1}\}$ . Let  $P$  be an arbitrary path in  $\mathcal{P}_1(H)$ . Say  $u$  and  $w$  are the two endvertices of  $P$ . Then  $I(uw, G_2) \cap B = \emptyset$ , i.e.,  $d^*(P, v_i) \leq 1$  for all  $i \in \{1, \dots, r-1\}$ . It follows that  $d(uw, G_2) \leq n_2 + 3$  and if equality holds then  $r = n_2 - 2$  and  $\{v_{n_2-2}, v_{n_2-1}, v_{n_2}\} = I(uw, G_2)$ . Clearly,  $d(uw, G_1) \geq n_1 + n_2 + 4 - (n_2 + 3) = l + 2$  and so  $d(uw, H) \geq l$ . Claim *A* follows.  $\square$

We now break into two cases here.

Case 1.  $H \notin \mathcal{H}_1$ .

Then  $d^*(P, H) \leq l$  by Lemma 3.3 and so  $d^*(P, G_2) \geq n_1 + n_2 + 4 - l - d^*(P, u_0) \geq n_2 + 3$  for all  $P \in \mathcal{P}_1(H)$ . First, assume that  $d(u_0, H) \leq (n_1 + 1)/2$ . By Claim *A* and Lemma 3.3,  $r = n_2 - 2$  and for each  $P \in \mathcal{P}_1(H)$ ,  $d^*(P, H) = l$ ,  $\sigma(e_1, P) \neq 0$ ,  $d^*(P, \{u_0, v_{n_2-2}, v_{n_2-1}, v_{n_2}\}) = 8$ , and  $d^*(P, v_i) = 1$  for all  $1 \leq i \leq r - 1$ . We apply Lemma 3.5(c) to  $H$ . First, assume that  $V(H)$  has a partition  $X \cup Y$  such that  $|X| = l/2$ ,  $V(e_1) \subseteq X$  and  $N(y, H) = X$  for all  $y \in Y$ . Then any two distinct vertices in  $Y$  can play the role of the two endvertices of  $P$ . Hence  $d(v_1, Y) \geq l/2 - 1 \geq 2$  and so  $H + v_1 \in \mathcal{H}_1$ , a contradiction. Therefore  $H - V(e_1)$  has exactly two components  $H_1$  and  $H_2$  such that both  $H_1$  and  $H_2$  are complete and  $d(x, H_1 \cup H_2) = l - 2$  for each  $x \in V(e_1)$ . It follows that  $V(H_1 \cup H_2) \subseteq N(u_0)$ . Thus  $n_1 - 3 \leq d(u_0, G_1) \leq (n_1 + 1)/2$ . This implies that  $n_1 \leq 7$ . As mentioned in the beginning paragraph, we have  $r = n_2 - 2$  and  $n_1 \geq n_2 \geq 8$ , a contradiction.

Therefore  $d(u_0, H) \geq (n_1 + 2)/2$ . Let  $P = z_1 \dots z_l$  be arbitrary in  $\mathcal{P}_1(H)$  with  $z_1 \in S_1(H)$ . We claim  $d(z_1, G_1) \geq (n_1 + 2)/2$ . If this is not true, say  $d(z_1, G_1) \leq (n_1 + 1)/2$ . Then  $d(z_1, G_2) \geq (n_2 + 3)/2$ . Clearly,  $d(z_1, J) \geq \lceil (n_2 + 3)/2 - (n_2 - r) \rceil \geq 4$  as  $r - 1 \geq (n_2 + 2)/2$  and so  $d(z_1, J^*) > 0$ . By Lemma 3.2(a),  $G_1 - z_1 = H - z_1 + u_0 \in \mathcal{P}_1$ . As  $d(z_1, G_1) < d(u_0, G_1)$  and by the minimality of  $d(u_0, G_1)$ ,  $G_2 + z_1 \notin \mathcal{H}_2$ , i.e.,  $z_1 v_{n_2} \notin E$  and if  $v_{n_2} v_{n_2-2} \in E$  then  $z_1 v_{n_2-1} \notin E$ . By Lemma 3.2(b),  $G_2 + z_1 \in \mathcal{P}_2$ . If  $v \in S_2(G_2 + z_1)$  then  $d(v, G_2 + z_1) \geq (n_2 + 3)/2$  for otherwise  $\xi(z_1, v) > 0$ ,  $d(v, G_1) \geq (n_1 + 2)/2$  and  $G_1 - z_1 + v \in \mathcal{P}_1$  by Lemma 3.2(a), contradicting (1). Let  $s$  be the maximal index such that  $z_1 v_s \in E$ . Set  $r' = \max\{r, s\}$ . By Lemma 3.9, for all  $v \in \{z_1, v_1, \dots, v_{r'-1}\}$ ,  $d(v, G_2 + z_1) \geq (n_2 + 3)/2$ ,  $N(v, G_2 + z_1) \subseteq \{z_1, v_1, \dots, v_{r'}\}$  and  $G_2 + z_1$  has a  $v_{n_2-v}$   $e_2$ -hamiltonian path. Therefore  $d(v, G_2) \geq (n_2 + 1)/2$  for all  $v \in \{v_1, \dots, v_{r'-1}\}$ . It follows that  $r' = r$  or  $r' = r + 1$ . As  $d(z_1 z_l, G_2) \geq n_2 + 3$ ,  $i(z_1 z_l, J + v_{r'}) \geq 3$ . As  $I(z_1 z_l, B) = \emptyset$ , we see that  $I(z_1 z_l, G_2) = \{v_c, v_r, v_{r+1}\}$ . It follows that  $d(z_1 z_l, H) = l$ ,  $N(z_1 z_l, G_2) = V(G_2)$ ,  $d(u_0, z_1 z_l) = 2$ ,  $B = V(J) - \{v_c, v_r\}$  and  $d(v_i, z_1 z_l) = 1$  for all  $v_i \in B$ . This argument implies that for any  $u$ - $v$  path in  $\mathcal{P}_1^*(H)$ ,  $d(uv, H) = l$  because  $\min\{d(u, H), d(v, H)\} \leq l/2$  and so  $\min\{d(u, G_1), d(v, G_1)\} \leq (n_1 + 1)/2$ .

We now apply Lemma 3.5(c) to  $H$ . First, assume that  $V(H)$  has a partition  $X \cup Y$  such that  $|X| = l/2$ ,  $V(e_1) \subseteq X$  and  $N(y, H) = X$  for all  $y \in Y$ . Then any two distinct vertices in  $Y$  can play the role of the two endvertices of  $P$ . Hence  $d(v_i, Y) \geq l/2 - 1 \geq 2$  and so  $H + v_i \in \mathcal{H}_1$  for each  $v_i \in B$ , a contradiction. Therefore  $H - V(e_1)$  has exactly two components  $H_1$  and  $H_2$ . Say  $z_1 \in V(H_1)$  and  $z_l \in V(H_2)$ . Then  $z_1$  can be any vertex in  $H_1$  and  $z_l$  can be any vertex in  $H_2$  for the above argument. Consequently,  $V(H_2) \subseteq N(v_{n_2})$  and  $V(H_1 \cup H_2) \subseteq N(v_c) \cap N(u_0)$ . Clearly,  $G_1 - x + v_c \in \mathcal{H}_1$  for any  $x \in V(H_2)$ . Let  $v_d \in B - \{v_c\}$ . If  $xv_d \in E$  for some  $x \in V(H_2)$  then  $G_2 - v_c + x \in \mathcal{H}_2$ , a contradiction. Therefore  $d(v_d, H_2) = 0$  and so  $N(v_d, H_1) = V(H_1)$ . As  $d(v_d, G_1) \geq (n_1 + n_2 + 4)/2 - (n_2 - 2) \geq 4$ ,  $|H_1| \geq 2$ . As  $V(H_1 \cup H_2) \subseteq N(u_0)$ , we see  $G_1 - z_l + v_d \in \mathcal{H}_1$ . As  $G_2 - v_d$  has a  $v_{n_2-v_c}$   $e_2$ -hamiltonian path,  $G_2 - v_d + z_l \in \mathcal{H}_2$ , a contradiction.

Therefore  $d(z_1, G_1) \geq (n_1 + 2)/2$  and so  $d(z_1, H) \geq (l + 1)/2$ . Thus  $\delta_1(H) \geq (l + 1)/2$ . As  $H \notin \mathcal{H}_1$  and by Lemma 3.3,  $\mathcal{P}_1^*(H) = \emptyset$  and so  $e_1 = z_l z_{l-1}$ . As  $d(u_0, H) \geq (n_1 + 2)/2$  and by Lemma 3.2(a),  $H - z_1 + u_0 \in \mathcal{P}_1$ . As  $H \notin \mathcal{H}_1$ ,  $d(z_1 z_l, H) \leq l - 1$  by Lemma 3.3. Choose  $P$  to be an optimal path at  $e_1$  in  $H$ . Say  $t = \alpha(P, z_1)$ . By Lemma 3.9,  $C = z_1 z_2 \dots z_t z_l$  is an end-cycle at  $z_t$  in  $H$  such that  $d(z_i, C) \geq (l + 1)/2$  for all  $i \in \{1, 2, \dots, t - 1\}$ . Thus for all  $i \in \{1, 2, \dots, t - 1\}$ , each  $z_i$  can play the role of  $z_1$  in the above and so  $d(z_i, G_1) \geq (n_1 + 2)/2$ . Clearly,  $d(u_0, C - z_t) > 0$ . Say w.l.o.g.  $u_0 z_1 \in E$ . As  $\mathcal{P}_1^*(H) = \emptyset$ ,  $N(z_l, G_1) \subseteq \{z_{l-1}, z_{l-2}, u_0\}$ . Clearly,  $d(z_l, J - v_r) \geq (n_1 + n_2 + 4)/2 - 3 - (n_2 - r + 1) > 0$ . Recall that  $u_0 v_{n_2} \in E$ . Therefore if we set  $G' = G_1 - V(e_1) + V(e_2)$  and  $G'' = G_2 - V(e_2) + V(e_1)$ , then  $G' \in \mathcal{H}_2$  and  $G'' \in \mathcal{H}_1$ . Recall that if  $z_l z_{l-2} \in E$  then  $N(z_{l-1}, G_1) \subseteq \{z_l, z_{l-2}, u_0\}$  as  $\mathcal{P}_1^*(H) = \emptyset$ . We readily see that  $d(e_1, G_1 - V(e_1)) \leq l$ ,  $d(e_1, G_2) \geq (n_1 + n_2 + 4) - l - 2 = n_2 + 3$ ,  $d(e_2, G_2 - V(e_2)) \leq n_2 - 2$  and  $d(e_2, G_1) \geq n_1 + n_2 + 4 - (n_2 - 2) - 2 = n_1 + 4$ . Thus

$$\begin{aligned}
e(G') + e(G'') &= e(G_1) - d(e_1, G_1 - V(e_1)) + d(e_1, G_2) \\
&+ e(G_2) - d(e_2, G_2 - V(e_1)) + d(e_2, G_1) - 2d(e_1, e_2) \\
&\geq e(G_1) + e(G_2) + 10 - 2d(e_1, e_2).
\end{aligned} \tag{2}$$

As  $10 - 2d(e_1, e_2) \geq 2$  and by (1), we see that  $n_1 = |G'| \neq n_2$ . As  $n_1 \geq n_2$ ,  $n_1 > n_2$ . As  $N(z_l, G_1) \subseteq \{z_{l-2}, z_{l-1}, u_0\}$ , we obtain that  $n_2 \geq d(z_l, G_2) \geq \lceil (n_1 + n_2 + 4)/2 - d(z_l, G_1) \rceil = n_2$ . It follows that  $N(z_l, G_1) = \{z_{l-2}, z_{l-1}, u_0\}$  and  $d(z_l, G_2) = n_2$ . As  $H + v_j \notin \mathcal{H}_1$  for all  $v_j \in B$ , it follows that  $z_i v_j \notin E$  for all  $i \in \{1, \dots, t-1\}$  and  $v_j \in J - \{v_c, v_r\} \subseteq B$ . Thus  $d(z_1, G_1) + d(z_1, G_2) \leq t + n_2 - r + 2$ . Let  $v \in J - \{v_c, v_r\}$ . Then  $d(v, G_1) + d(v, G_2) \leq l - t + 2 + r - 1$ . Consequently,  $d(z_1) + d(v) \leq n_1 + n_2 + 2$ . But  $d(z_1) + d(z_2) \geq n_1 + n_2 + 4$  as  $\delta(G) \geq (n_1 + n_2 + 4)/2$ , a contradiction.

Case 2.  $H \in \mathcal{H}_1$ .

Let  $C = z_1 z_2 \dots z_l z_1$  be an  $e_1$ -hamiltonian cycle of  $H$  with  $e_1 = z_1 z_2$ . Let  $v_i \in B$ . With the details stated in the beginning paragraph, we see that  $d(v_i, H) \geq \lceil (n_1 + n_2 + 4)/2 \rceil - (r - 1) - d(v_i, u_0) \geq 4$  and if equality holds then  $v_i u_0 \in E$ ,  $r \in \{n_2 - 2, n_2 - 1\}$  and  $d(v_i, G_2) = r - 1$ . We divide this case into the following two subcases.

Subcase 2.1. For each path  $P \in \mathcal{P}_1^*(H)$ ,  $d^*(P, H) \geq l + 1$ .

First, assume that  $d(w, C) \leq l/2$  for some  $w \in V(C) - V(e_1)$ . If  $w \notin \{z_3, z_l\}$  then  $d(x, C) \geq (l + 2)/2$  for all  $x \in V(C - w) - V(e_1)$  by Lemma 3.8. As  $d(v_i, C) \geq 4$  and  $H + v_i \notin \mathcal{H}_1$ , we readily see that there exist two distinct vertices  $z_j$  and  $z_h$  in  $N(v_i, C)$  such that either  $\{z_j^-, z_h^-\} \subseteq V(C) - \{z_1, z_2, w\}$  or  $\{z_j^+, z_h^+\} \subseteq V(C) - \{z_1, z_2, w\}$ . Consequently, by Lemma 3.6,  $H$  has a  $z_j$ - $z_h$   $e_1$ -hamiltonian path and so  $H + v_i$  is hamiltonian, a contradiction. Therefore  $d(z_j, C) \geq (l + 2)/2$  for all  $4 \leq j \leq l - 1$ . As above,  $N(v_i, C)$  does not contain two distinct vertices  $z_j$  and  $z_h$  such that either  $\{z_j^-, z_h^-\} \subseteq V(C) - \{z_1, z_2, z_3, z_l\}$  or  $\{z_j^+, z_h^+\} \subseteq V(C) - \{z_1, z_2, z_3, z_l\}$ . It follows that  $N(v_i, C) = \{z_1, z_2, z_4, z_{l-1}\}$ . The above argument allows us to conclude that  $r \in \{n_2 - 2, n_2 - 1\}$ , and for all  $v \in B$ ,  $N(v, H) = \{z_1, z_2, z_4, z_{l-1}\}$ . As  $V(J) - \{v_c, v_r\} \subseteq B$ ,  $d(w, G_2) \leq 4$  and so  $d(w, C) \geq (n_1 + n_2 + 4)/2 - 5 \geq (l + 1)/2$ , a contradiction.

Therefore  $d(z_i, H) \geq (l + 1)/2$  for all  $i \in \{3, \dots, l\}$ . As  $d(v_i, H) \geq 4$  and  $H + v_i \notin \mathcal{H}_1$ , there exist two distinct vertices  $u$  and  $v$  in  $C - V(e_1)$  such that either  $\{u^+, v^+\} \subseteq N(v_i)$  or  $\{u^-, v^-\} \subseteq N(v_i)$ . Say w.l.o.g.  $\{u^+, v^+\} \subseteq N(v_i)$ . Then  $H$  does not have a  $u^+ - v^+$   $e_1$ -hamiltonian path. We apply Lemma 3.6(d) to  $H$ . First, assume that  $H$  has a vertex-cut  $X$  with  $V(e_1) \subseteq X$  and  $|X| = 3$  such that  $H - X$  has exactly two components isomorphic to  $K_{(l-3)/2}$  and  $X \subseteq N(y)$  for all  $y \in V(C) - X$ . Obviously,  $H + v_i \in \mathcal{H}_1$ , a contradiction. Thus  $V(H)$  has a partition  $X \cup Y$  such that  $|X| = (l + 1)/2$ ,  $|Y| = (l - 1)/2$ ,  $\{u^+, v^+\} \cup V(e_1) \subseteq X$  and  $N(y, H) = X$  for all  $y \in Y$ . As  $H + v_i \notin \mathcal{H}_1$ , it follows that  $N(v_i, H) \subseteq X$ . Let  $y \in Y$ . Then  $d(y, G_2) \geq (n_1 + n_2 + 4)/2 - (l + 1)/2 - 1 = (n_2 + 2)/2$ . Thus  $d(y, B) > 0$ . Let  $v_j \in N(y, B)$ . With  $v_j$  in place of  $v_i$  in the above argument, we see that  $V(H)$  has a partition  $X'$  and  $Y'$  such that  $|X'| = (l + 1)/2$ ,  $V(e_1) \subseteq X'$ ,  $N(v_j, H) \subseteq X'$  and  $X' = N(y', H)$  for all  $y' \in Y'$ . As  $Y$  is an independent set, it follows that  $Y \subseteq X'$  and so  $|X'| \geq (l - 1)/2 + 2 = (l + 3)/2$ , a contradiction.

Subcase 2.2. There exists  $P = z_1 z_2 \dots z_l \in \mathcal{P}_1^*(H)$  such that  $d(z_1 z_l, H) \leq l$ .

Then  $d(z_1 z_l, G_2) \geq n_1 + n_2 + 4 - l - d(u_0, z_1 z_l) \geq n_2 + 3$  and so  $i(z_1 z_l, G_2) \geq 3$ . Say  $d(z_1, G_1) \leq d(z_l, G_1)$ . Then  $d(z_1, G_1) \leq l/2 + 1 = (n_1 + 1)/2$ . Thus  $d(z_1, G_2) \geq (n_2 + 3)/2$ . As  $r \geq \delta_2(G_2) + 1 \geq (n_2 + 2)/2 + 1$ ,  $d(z_1, J - v_r) \geq \lceil (n_2 + 3)/2 - (n_2 - r) - 1 \rceil \geq 3$ . Therefore  $G_2 + z_1$  has a hamiltonian path from  $e_1$  to  $z_1$ . We claim that  $G_1 - z_1 \in \mathcal{P}_1$ . If this is not true, then  $d(u_0, P - z_1) \leq (l - 1)/2$  by Lemma 3.2(a) and so  $d(u_0, G_1) \leq (l + 1)/2$ . By Claim A, it follows that

$$d(z_1 z_l, H) = l, r = n_2 - 2, I(z_1 z_l, G_2) = \{v_{n_2-2}, v_{n_2-1}, v_{n_2}\} \text{ and } d(u_0, z_1 z_l) = 2. \tag{3}$$

Therefore  $G_1 - z_1 \in \mathcal{P}_1$ . By the minimality of  $u_0$ ,  $d(u_0, G_1) \leq d(z_1, G_1) \leq (n_1 + 1)/2$ . Therefore (3) still holds and  $d(u_0, G_1) \leq (n_1 + 1)/2$  in any case. Moreover,  $d(u_0, J) \geq \lceil (n_2 + 3)/2 \rceil - (n_2 - r) \geq 4$  and so  $B = V(J) - \{v_r\}$  as mentioned in the paragraph above Claim A. As  $r - 1 \geq (n_2 + 2)/2$ ,  $n_2 \geq 8$ .

We claim that for each  $\{u, v\} \subseteq V(J) - \{v_r\}$  with  $u \neq v$ ,  $G_2 - \{u, v\} + \{u_0, z_l\} \in \mathcal{H}_2$ . To see this, we note that  $u_0v_{n_2}v_{n_2-1}v_{n_2-2}z_lu_0$  is a cycle in  $G$ . Moreover, we have that for all  $x \in V(J - \{u, v, v_r\})$ ,  $d(x, J - \{u, v, v_r\}) \geq (n_2 + 2)/2 - 3 = ((n_2 - 5) + 1)/2$  and so  $J - \{u, v, v_r\}$  is hamiltonian connected. Clearly, for each  $y \in \{u_0, z_l\}$   $d(y, J - \{u, v, v_r\}) \geq \lceil (n_2 + 3)/2 \rceil - 5 \geq 1$  as  $n_2 \geq 8$ . Thus if  $G_2 - \{u, v\} + \{u_0, z_l\} \notin \mathcal{H}_2$ , then  $d(y, J - \{u, v, v_r\}) = 1$  for each  $y \in \{u_0, z_l\}$ . Consequently,  $n_2 \leq 9$ . As  $\delta_2(G_2) \geq \lceil (n_2 + 2)/2 \rceil$ , it follows that  $J$  is complete and obviously  $G_2 - \{u, v\} + \{u_0, z_l\} \in \mathcal{H}_2$ , a contradiction. Hence the claim holds.

Therefore  $H - z_l + u + v \notin \mathcal{H}_1$  for all  $u, v \in V(J - v_r)$  with  $u \neq v$ . For each vertex  $v \in V(J - v_r)$ , it is easy to see that  $uv \in E$  for some  $u \in N(z_1, J - v_r)$  since  $d(z_1, G_2) \geq (n_2 + 3)/2$  and  $d(v, J) \geq (n_2 + 2)/2$ . Therefore  $d(z_{l-1}, J - v_r) = 0$  for otherwise  $H - z_l + u + v \in \mathcal{H}_1$  for some  $v \in N(z_{l-1}, J - v_r)$  and  $u \in N(z_1, J - v_r)$  with  $uv \in E$ . Thus  $d(z_{l-1}, H - z_l) \geq (n_1 + n_2 + 4)/2 - 5 = (n_1 + n_2)/2 - 3$ . Let  $uv \in E(J - v_r)$  with  $uz_1 \in E$ . Clearly,  $d(v, H - z_l) \geq (n_1 + n_2 + 4)/2 - (r - 1) - 2 = (n_1 - n_2)/2 + 3$ . Thus  $d(vz_{l-1}, H - z_l) \geq (l - 1) + 2$ . By Lemma 3.2(d),  $H - z_l + v$  has an  $e_1$ -hamiltonian path from  $z_1$  to  $v$  and so  $H - z_l + u + v \in \mathcal{H}_1$ , a contradiction. This proves the lemma.  $\blacksquare$

**Proof of Lemma 2.5.** Choose  $v' \in J^*$ . Then  $d(v'v_{n_2}, G_1) \geq n_1 + n_2 + 4 - (n_2 - 1) = n_1 + 5$ . Thus  $i(v'v_{n_2}, G_1) \geq 5$ . By Lemma 2.4,  $G_1 - u \notin \mathcal{P}_1$  for all  $u \in I(v'v_{n_2}, G_1) - V(e_1)$ . Therefore  $G_1 \notin \mathcal{H}_1$ . By Property B,  $\mathcal{P}_1^*(G_1) \neq \emptyset$ . We claim  $\delta_1(G_1) \leq (n_1 - 1)/2$ . To see this, say  $\delta_1(G_1) \geq n_1/2$ . Choose any path from  $\mathcal{P}_1^*(G_1)$  and then apply Lemma 3.5(c) with this path in  $G_1$ . As  $d(v_{n_2}, G_1) \geq (n_1 + n_2 + 4)/4 - 2 = (n_1 + n_2)/2$ , we see that  $G_1$  has an  $x$ - $y$   $e_1$ -hamiltonian path such that  $y \notin V(e_1)$ ,  $d(y, G_1) = n_1/2$  and  $yv_{n_2} \in E$ . As  $d(y, G_2) \geq (n_2 + 4)/2$ ,  $d(y, J^*) > 0$  and so  $G_2 + y \in \mathcal{H}_2$ , contradicting Lemma 2.4.  $\blacksquare$

**Proof of Lemma 2.6.** The statement (a) is evident by the definition of  $(G_{2i-1}, G_{2i})$  ( $1 \leq i \leq k$ ). We show (b) by contradiction. Say on the contrary that  $d(v, G_{2i}) \leq (|G_{2i}| + 3)/2$  for some  $v \in S_2(G_{2i})$  and  $i \in \{1, \dots, k\}$ . Let  $i$  be minimal. Then  $d(v, G_{2i-1}) \geq (|G_{2i-1}| + 1)/2$  and so  $G_{2i-1} + v \in \mathcal{P}_1$  by Lemma 3.2(a). As  $\mathcal{P}_2^*(G_{2i}) = \emptyset$ ,  $\mathcal{P}_2^*(G_{2i} - v) = \emptyset$ . By the maximality of  $e(G_{2(i-1)-1}) + e(G_{2(i-1)})$ , we shall have

$$\begin{aligned} e(G_{2(i-1)-1}) + e(G_{2(i-1)}) &\geq e(G_{2i-1} + v) + e(G_{2i} - v) \\ &\geq e(G_{2i-1}) + e(G_{2i}) - (|G_{2i}| + 3)/2 + (|G_{2i-1}| + 1)/2. \end{aligned} \quad (4)$$

Let  $P = v_qv_{q-1} \dots v_1$  be an optimal path at  $e_2 = v_qv_{q-1}$  in  $G_{2(i-1)}$ , where  $q = |G_{2(i-1)}|$ . Say  $\alpha(P, v_1) = r$ . As  $\delta_2(G_{2(i-1)}) \geq (|G_{2(i-1)}| + 4)$  and  $\mathcal{P}_2^*(G_{2(i-1)}) = \emptyset$ , we see that  $v_1v_2 \dots v_rv_1$  is an end-cycle at  $v_r$  in  $G_{2(i-1)}$ . As  $d(w_{i-1}, G_{2(i-1)}) \geq (|G_{2(i-1)}| + 5)/2$  and  $G_{2(i-1)} + w_{i-1} \notin \mathcal{H}_2$ , we see that  $\mathcal{P}_2^*(G_{2(i-1)} + w_{i-1}) = \emptyset$ . By the maximality of  $e(G_{2(i-1)-1}) + e(G_{2i})$ , we shall have

$$\begin{aligned} e(G_{2i-1}) + e(G_{2i}) &\geq e(G_{2(i-1)-1} - w_{i-1}) + e(G_{2(i-1)} + w_{i-1}) \\ &\geq e(G_{2(i-1)-1}) + e(G_{2(i-1)}) - (|G_{2(i-1)-1}| - 1)/2 + (|G_{2(i-1)}| + 5)/2. \end{aligned} \quad (5)$$

By (4) and (5), we see that  $e(G_{2(i-1)-1}) + e(G_{2(i-1)}) > e(G_{2(i-1)-1}) + e(G_{2(i-1)})$ , a contradiction.  $\blacksquare$

**Proof of Lemma 2.7.** On the contrary, say the claim fails. Let  $x_0 \in V(G_{2k-1})$  such that  $G_{2k-1} - x_0 \in \mathcal{P}_1$ ,  $G_{2k} + x_0 \in \mathcal{H}_2$  and  $d(x_0, R' - \{y_1, y_{r-1}\}) > 0$ . Let  $y_c \in V(R') - \{y_1, y_{r-1}\}$  with  $x_0y_c \in E$ . Since  $G_{2k} + x_0 \in \mathcal{H}_2$  and  $\mathcal{P}_2^*(G_{2k}) = \emptyset$ , either  $x_0y_t \in E$  or  $x_0y_{t-1} \in E$  with  $y_t y_{t-2} \in E$ . Say w.l.o.g.  $x_0y_t \in E$ .

Set  $H = G_{2k-1} - x_0$  and  $p = |H| = s - 1$ . As  $s \geq t$  and  $t - 1 \geq r$ , for each  $y \in V(R')$ ,  $d(y, H) \geq \lceil (s + t + 4)/2 - (r - 1) - d(y, x_0) \rceil \geq 3$ .

Assume for the moment that for every  $P \in \mathcal{P}_1(H)$ ,  $d^*(P, H) \geq p + 2$  for each  $P \in \mathcal{P}_1(H)$ . By Lemma 3.3,  $H \in \mathcal{H}_1$ . By Lemma 3.7,  $d(uv, H) \geq p + 2$  for all  $u, v \in V(H)$  with  $u \neq v$  and  $\{u, v\} \neq V(e_1)$ . Let  $y_i$  and  $y_j$  be two distinct vertices of  $R' - y_c$  such that  $\{y_i, y_j\} \neq \{y_1, y_{r-1}\}$  and  $y_i y_j \in E$ . Let  $C$  be an  $e_1$ -hamiltonian cycle of  $H$ . Then there is an orientation of  $C$  such that for some  $u, v \in V(C)$  with  $u \neq v$  and  $V(e_1) \neq \{u, v\}$ , we have  $e_1 \notin \{uu^+, vv^+\}$  and  $\{y_i u^+, y_j v^+\} \subseteq E$ . Let  $y' \in N(y_r, R' - y_c)$  be such that  $y' \notin \{y_i, y_j\}$ . By Lemma 3.6,  $H$  has a  $u^+ - v^+$   $e_1$ -hamiltonian path. Since Theorem B holds for  $R'$ ,  $R'$  has two disjoint paths  $P''$  and  $P'$  such that  $|P''| = n_1 - p$ ,  $|P'| = r - 1 - |P''|$ ,  $P''$  is from  $y_i$  to  $y_j$  and  $P'$  is from  $y'$  to  $y_c$ . Thus  $[H, P''] \in \mathcal{H}_1$  and  $G_{2k} - V(P'') + x_0 \in \mathcal{H}_2$ , i.e.,  $G$  contains two required cycles, a contradiction.

Therefore  $d^*(P, H) \leq p + 1$  for some  $P \in \mathcal{P}_1(H)$ . Say  $P = z_1 \dots z_p$ . First, assume that  $d(y_i, z_1 z_p) > 0$  for some  $y_i \in V(R') - \{y_1, y_{r-1}, y_c\}$ . Say w.l.o.g.  $z_1 y_i \in E$ . Then  $y_i z_p \notin E$ . If there exists  $z_p y_j \in E$  for some  $y_j \in N(y_i, R') - \{y_c\}$  then we obtain the two required cycles as above. Therefore  $z_p y_j \notin E$  for all  $y_j \in N(y_i, R') - \{y_c\}$ . Thus  $d(z_p, R) \leq r - (d(y_i, R) - 2)$  and so  $d(z_p, G_{2k}) \leq t - r + r - (d(y_i, R) - 2) = t - d(y_i, R) + 2$ . As  $d(y_i, R) \geq (t+4)/2$ ,  $d(z_p, G_{2k}) \leq t/2$ . Therefore  $d(z_p, H) \geq (s+t+4)/2 - t/2 - d(z_p, x_0) \geq (s+2)/2$ . Similarly, if  $z_p y_1 \in E$  then  $z_1 y_a \notin E$  for each  $y_a \in N(y_1, R' - \{y_{r-1}, y_c\})$ . Consequently,  $d(z_1, R) \leq r - (d(y_1, R) - 3)$  and  $d(z_1, G_{2k}) \leq t - d(y_1, R) + 3 \leq (t+2)/2$ . It follows that  $d(z_1, H) \geq s/2$  and so  $d(z_1 z_p, H) \geq s + 1 = p + 2$ , a contradiction. Therefore  $z_p y_1 \notin E$ . Similarly,  $z_p y_{r-1} \notin E$ . Thus  $N(z_p, R) \subseteq \{y_r, y_c\}$  and so  $d(z_p, G_{2k}) \leq t - r + 2$ . Let  $y_j \in N(y_i, R') - \{y_c\}$ . Then  $d(z_p y_j, G_{2k}) \leq t - r + 2 + r - 1 = t + 1$ . Thus  $d(z_p y_j, H) \geq s + t + 4 - (t + 1) - d(x_0, z_p y_j) \geq p + 2$ . By Lemma 3.2(d),  $H + y_j$  has a  $z_1$ - $y_j$   $e_1$ -hamiltonian path and so  $H + y_i + y_j$  has a  $y_i$ - $y_j$   $e_1$ -hamiltonian path. As above, we see that  $G$  contains two required cycles, a contradiction.

Therefore  $N(z_1, R) \cup N(z_p, R) \subseteq \{y_1, y_{r-1}, y_r, y_c\}$  and so  $d(z_1 z_p, G_{2k}) \leq 2(t - r) + 8$ . As  $r \geq \delta_2(G_{2k}) + 1 \geq (t + 6)/2$ , we get  $d(z_1 z_p, G_{2k}) \leq t + 2$ . Therefore  $p + 1 \geq d(z_1 z_p, H) \geq s + t + 4 - (t + 2) - d(x_0, z_1 z_p) \geq p + 1$ . This implies that  $N(z_1, R) = N(z_p, R) = \{y_1, y_{r-1}, y_r, y_c\}$ ,  $r = (t + 6)/2$  and  $R \cong K_{(t+6)/2}$ . It follows that  $G$  contains two required cycles as above. ■

**Acknowledgements** This work is supported by NSA Grant H98230-08-1-0098.

## 5 References

- [1] S. Abbasi, Spanning cycles in dense graphs, Ph.D. thesis, Rutgers University 1999.
- [2] B. Bollobá, Extremal Graph Theory, Academic Press, London (1978).
- [3] J. Bondy and V. Chvátal, a method in graph theory, Discrete Mathematics, 15(1976), 111–135.
- [4] Y. Egawa, R. Faudree, E. Gyori, Y. Ishigami, R. Schelp, and H. Wang, Vertex-disjoint cycles containing specified edges, Graphs and Combinatorics, 16(2000), 81–92.
- [5] M.H. El-Zahar, On circuits in graphs, Discrete Math. 50(1984), 227–230.
- [6] P. Erdős and T. Gallai, On maximal paths and circuits of graphs, Acta Math. Acad. Sci. Hungar. 10(1959), 337–356.
- [7] H. Wang, Covering a graph with cycles passing through given edges, Journal of Graph Theory, 26(1997), 105–109.
- [8] J.E. Williamson, Panconnected graph II, period. Math. Hungar., 8(1977), 105–116.