

College of Science  
P.O. Box 443025, University of Idaho  
Moscow, ID 83844-3025

---

---

SCIENTIFIC AREA: **Earth Sciences (Global Change)**

PROJECT TITLE:

**PALEO-CLIMATIC and ENVIRONMENTAL ICE CORE RESEARCH,  
DATA ANALYSIS and INTERPRETATION**  
(mid- low- latitudes, high altitudes glaciers at the Northern Hemisphere)

PI: **Vladimir B. Aizen**  
CONTACT: Tel: 208-885-5888, fax: 208-885-5724, e-mail: [aizen@uidaho.edu](mailto:aizen@uidaho.edu)

**ANNUAL REPORT**

**DOE/INEEL Contract Number: BBWI 0000073 R10**  
**Budget Period: 01/10/2002 to 30/09/2003**

---

---

*20 February, 2004*

RESEARCH PERSONELL ' FY 2002/2003:

Professor, Dr. **Vladimir Aizen**, glaciologist  
Department of Geography, College of Science, P.O. Box  
443025, University of Idaho  
Moscow, ID 83844-3025, **U.S.A.**  
Tel. (208) 885-5888, fax. (208) 885-5724  
E-mail: [aizen@uidaho.edu](mailto:aizen@uidaho.edu)

Associate Professor, Dr. **Elena Aizen**, climatologist  
Department of Geography, College Science, P.O. Box  
443025, University of Idaho  
Moscow, ID 83844-3025, **U.S.A.**  
Tel. (208) 885-5888, fax. (208) 885-5724  
E-mail: [eaizen@bren.ucsb.edu](mailto:eaizen@bren.ucsb.edu)

Professor, Dr. **John D. Marshall**, Director: Idaho Stable  
Isotopes Laboratory, College of Natural Resources, P.O.  
Box 441133, Moscow, ID 83844-1133, **U.S.A.**  
Tel. (208) 885-6695, fax. (208) 885-6226  
E-mail: [jdm@uidaho.edu](mailto:jdm@uidaho.edu)

Professor, Dr. **Scott Wood**, Department of Geology,  
College of Science, P.O. Box 443025, University of Idaho  
Moscow, ID 83844-3025, **U.S.A.**  
Tel. (208) 885-5966, fax. (208) 885-5724  
E-mail: [swood@uidaho.edu](mailto:swood@uidaho.edu)

Professor, Dr. **Stanislav Nikitin**, geophysicist  
Glacio-Climatological Laboratory, Tomsk State University  
36 Lenina Str., Tomsk 634050, **Russian Federation**  
Tel. +7 (3822) 55-47-03, fax. +7 (3822) 41-58-92  
E-mail: [nikitin@ic.tsu.ru](mailto:nikitin@ic.tsu.ru)

Associate Professor, Dr. **Koji Fujita**, glaciologist  
Division of Earth and Environmental Sciences  
Nagoya University, Furo-cho, Chikusa-ku  
Nagoya 464-8605, **Japan**  
E-mail: [bri@ihas.nagoya-u.ac.jp](mailto:bri@ihas.nagoya-u.ac.jp)

Assistant Professor, Dr. **Nazomu Takeuchi**, microbiologist  
Kyoto Institute of Humanity and Nature, 335  
Takashimacho Kamigyo-ku, Kyoto 602-0878, **Japan**  
E-mail: [takeuchi@chikyu.ac.jp](mailto:takeuchi@chikyu.ac.jp)

Associate Professor, Dr. **Shiyin Liu**, glaciologist  
Cold and Arid Regions Environmental Engineering Research  
Institute, Chinese Academy of Sciences, Lanzhou, 260 West  
Donggang Rd, Lanzhou 730000, **China**  
E-mail: [liusy@ns.lzb.ac.cn](mailto:liusy@ns.lzb.ac.cn)

Assistant Professor, **Dr. Guangjian Wu**, isotope geochemist, Cold and Arid Regions Environmental Engineering Research Institute, Chinese Academy of Sciences, Lanzhou, 260 West Donggang Rd, Lanzhou 730000, **China**  
E-mail: [wugj@ns.lzb.ac.cn](mailto:wugj@ns.lzb.ac.cn)

Manager and cook Altai expedition Tatyana Prokopinskaya, MSc students Aleksei Lushnikov and Alexei Chebotarev, Tomsk State University, Russia; drill engineer Akiyoshi Takahashi, National Polar Research Institute, PhD students Jun Uetake and Tetsuhide Yamasaki, Kyoto Institute of Humanity and Nature, Japan; PhD student Arzhan Surazakov and MSc student Daniel Joswiak, University of Idaho, USA; MSc students Yong Han, Lihong You, Shi Li, Lanzhou Cold and Arid Areas Environmental Engineering Institute, China; support personnel from the University of Idaho, Kyoto Institute, Tomsk State University and Lanzhou Institute were also involved in the Siberian and Tibetan field and analytical research during FY2002/2003.

Table of Contents:

	Pages
RESEARCH PERSONNEL ' 2002/2003.....	2-3
Table of contents.....	4
<b>PREFACE</b> .....	5
<b>PART I. GLACIO-MONITORING 2001, 2002 and 2003 DEEP ICE-CORING EXPEDITION in SIBERIAN ALTAI</b>	
<b>Overview</b> .....	5-6
<b>Our research goal</b> .....	6
<b>Field Research</b> .....	6-9
<b>2003 ice-coring operation</b> .....	9-11
<b>Ice-core properties</b> .....	11
<b>The laboratory analysis and first results</b> .....	12
Stable isotopes from firn core.....	12-13
Establishing a depth–age relationship.....	13-14
Seasonal snow accumulation.....	14-15
Stable isotopes from firn cores and snow pits.....	15-16
Relationship between stable hydrogen- and oxygen-isotope ratios.....	16
Clustering precipitation transferred from external (oceanic) and internal (continental) moisture sources.....	17-18
Internal moisture source.....	18-19
Atlantic moisture source.....	19
Oceanic moisture source.....	19
Relationship between $\delta^{18}\text{O}$ and air temperature.....	19-20
Transfer functions and large-scale atmospheric patterns.....	21
<b>Findings</b> .....	21-22
<b>Further proposed laboratory and analytical research (Altai)</b> .....	22
<b>PART II. SOUTH-EASTERN TIBET AND HIMALAYA</b>	
<b>Overview</b> .....	23
<b>Field itinerary</b> .....	24
<b>Field results</b> .....	25
Zuoqiupu Glacier radio-echo sounding measurements.....	25-26
Meteorological observations.....	26
Shallow firn core and snow sampling.....	26-27
<b>Laboratory analyses</b> .....	27
<b>Discussion</b> .....	27-28
<b>PART III. The Uofl new ICE-CORE LABORATORY and FREEZER</b> .....28-29	
<b>References</b> .....	29-31
<b>Publications</b> .....	32-33
<b>APPENDIX</b>	
<b>Agreement with Lanzhou Cold and Arid Region Environmental Engineering Research Institute, China</b> .....	34-38

## 1. PREFACE

One of the most valuable information about past climate and environmental changes lies in glacial ice, which records preserve precipitated snow for hundreds to thousands of years. These records from pre-industrial or even prehistoric time can be examined through snow-ice stratigraphy and geochemical analysis for stable and radioactive isotopes, major ions, trace elements and green house gases (GHG). We can link this information with changes in atmospheric circulation, air temperature, snow accumulation, atmospheric composition, marine and continental biogenic activity, aerosol loading/volcanic eruptions, continental dust source regions, forest fire activity, anthropogenic emissions, solar variability, radionuclide deposition and the GHG chemical composition of the atmosphere. Supplementary long-term (50 to 150 years) information including meteorological, hydrological, and atmospheric chemistry observational data is used for our statistical calibration, validation and interpretation analyses.

Our project is a multi-disciplinary, multi-institutional, international effort in ice-coring paleo-climatic environmental research and we working to help better understand the impacts of global and environmental changes on the natural ecosystems. In this report, we present some results of our research performed in FY2002/2003.

The presented report is composed of three parts: 'Part I' is the results of two reconnaissance in summers 2001 and 2002, and the summer 2003 deep ice-coring expedition in the Siberian Altai, 'Part II' is a result from the first field reconnaissance in South-Eastern Tibet and Himalaya in the fall of 2002, 'Part III' describes our efforts in establishing a new ice-core processing laboratory and ice-core storage facilities at the University of Idaho. All the above research was conducted under the Project: 'Paleo-Climatic and Environmental Ice Core Research, Data Analysis and Interpretation in Asia Mountains'.

**Our research goal** is to recover ice-core isotope-geochemical records containing information on large-scale atmospheric dynamics, the precipitation-origin, the natural and anthropogenic impact on climatic variability during industrial and pre-industrial time to understand past and forecast future Global Changes

## PART I. GLACIO-MONITORING 2001, 2002 and 2003 DEEP ICE-CORING EXPEDITION in SIBERIAN ALTAI

### Overview

After successfully recovering two deep ice-cores from central Tien Shan in the summer 2000 we focused our research on Siberian Altai and south-eastern Tibet-Himalaya (**Figure 1.1**).




- 3000 m a.s.l. topographic line outlined the central Asia Mountain System
- ice-cores recovered with DOE support
- new developing ice-coring sites

The spatial coverage of available snow, firn and ice records is inadequate to document climatic and environmental change over the vast Asian continent. However, environmental records for the northwestern periphery of the central Asia mountain system (CAMS), obtained from Tien Shan firn/ice cores (*Kreutz and others*, 2001, 2003; *Aizen and others*, in press a, b) and firn/ice-core records from alpine areas in Siberia (*Aizen and others*, in press b; *Olivier and others*, 2003),

are extending the area of climatic and environmental analyses in Asia.

The research presented in this chapter focuses on the Siberian Altai, the most continental northern periphery of the CAMS and the southern periphery of the Asian Arctic Basin (**Figure 1.1**). It is an ideal area for the analysis of climatic records relating to the major Eurasian circulation systems (i.e. the westerly jet stream and the Siberian High). Altai glaciers, located at the center of Eurasian continent, store unique information on inter-hemispheric climate dynamics and on the internal and external hydrological cycles of northern Eurasia. They provide records on the advection of fresh water transported from the Atlantic, Pacific and Arctic Oceans. Firn-ice records from the Altai glaciers can also be associated directly with the large Aral–Caspian internal water system, because moisture is transferred from this closed Asian drainage basin to the great Siberian river basins. Furthermore, the Siberian Altai is only mountain system in the Asian Arctic Basin (**Figure 1.1**) with alpine glaciers that nourish the Siberian rivers. Altai glaciers, particularly their cold snow-firn accumulation zone are appropriate for studying air pollution dynamics at the center of Eurasia, eastward and northward from major air pollutants in Russia, Kazakhstan and China. During the 20<sup>th</sup> century the Altai Mountains became extremely contaminated from industrial activities including mining, metallurgy and chemical production, as well as from the nuclear test areas in Semipalatinsk (Kazakhstan), Lobnor (China), and the Baikonur rocket site (Russia).




**Figure 1.2.** West Belukha Plateau location, Siberian Altai

The major components reflecting physical processes in the atmosphere over the studied region are  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  stable-isotopes, major ions, and trace elements transported and deposited on the glaciers. Data on the stable-isotope and geochemical content in snow and ice cores have contributed to our understanding of environmental signals from different regions of the globe (*Aizen and others*, 1996, in press a, b; *Dansgaard and others*, 1993; *Froehlich and others*, 2002; *Johnsen*, 1977; *Jouzel and others*, 1997; *Naftz and others*, 2002). Information on the spatial distribution of stable isotopes and geochemical components in precipitation in the Siberian Altai is also required to evaluate the contribution of water vapor, associated with external and internal water cycles, to snow accumulation over the northern periphery of the CAMS.

### **Field research 2001, 2002**

The West Belukha plateau (**Figures 1.2 and 1.3**) is the only Siberian location where the Altai glaciers have cold enough temperatures and sufficient snow accumulation to preserve climatic and environmental records that are unaffected by meltwater percolation. The field site selected by *Oliver and others* (2003), on a saddle (4062 m a.s.l.) between the Belukha (4499 m a.s.l.) and West Belukha (4406 m a.s.l.) peaks (**Figures 1.2, 1.3 and 1.4**), was a little too low and, because of very strong snow and wind redistribution between the two peaks, unsuitable for ice-core research.

In the summers 2001, 2002 reconnaissance and 2003 deep ice-coring expedition, snow samples were collected every 3–5 cm from five 2–3 m snow pits that dug in the same location each year on the Belukha snow/firn plateau (4110–4120 m a.s.l.). In 2001, a 21 m shallow snow/firn core




**Figure 1.3.** Belukha glacial massif, Siberian Altai. View from Akkem Lake (northern side). Photo by V. Aizen, August 2003.

A Grant Instruments Co. automatic weather station was installed at 4100 m a.s.l., near a rocky cliff on the northern side of Plateau, about 0.5 km north of the drilling site (**Figure 1.4 and 1.5**). Wind speed, wind direction, air temperature, air humidity and barometric pressure sensors measured meteorological events every 3 hours from July 2002 to August 2003 (**Figure 1.5 and 1.6**).

Temperatures in the snow pits were measured every 10 cm from surface to bottom using electric sensors. The West Belukha snow/firn plateau is in the cold recrystallization zone where melt does not occur; positive temperatures were never observed at the drill site (**Figure 1.6**).

To monitor snow accumulation, we installed 5 wooden stakes, with location determined by GPS (Garmin eTrex summit). One stick was equipped with an automatic snow depth measurement KADEC-SNOW, KONA system. The geo-coordinates of these stakes, snow pits and drill site location are summarized in Table 1. The geodesic coordinate accuracy was  $\pm 0.01$  m in horizontally directions and  $\pm 0.05$  m in vertically.

was recovered at 4115 m a.s.l. ( $49^{\circ}48' N$ ,  $86^{\circ}33' E$ ), where radio-echo sounding indicated the ice thickness is about 180 m thick (**Figure 1.4**). Snow pit, precipitation, and fresh snow samples were collected and placed into pre-cleaned plastic bottles. Density was measured every 5 cm in each snow pit using a  $100 \text{ cm}^3$  stainless steel sampler. A firn core (19 m long and 9.5 cm in diameter) was extracted from the bottom of the 220 cm snow pit (number 4) with a PICO fiberglass auger. The diameter, length and weight of each recovered core section were measured to calculate density. The 2001 firn-core sections, sealed in pre-cleaned polyethylene bags, were packed in insulated shipping containers and delivered to the National Institute of Polar Research in Tokyo (NIPR) for analysis (*at that time we didn't have facilities in UofI for this work*).




**Figure 1.4.** Glacier ice thickness profiles, snow pits and ice-core sites, on the Western Belukha Plateau.

The accumulation gauge sensor detects snow or open air by photo-diodes at an interval of 1 cm. Since strong solar radiation during noontime infiltrates into snow causing an apparent lowering of the surface level, the noontime records were discarded. Two records, morning and evening were chosen and validated to represent the actual surface level.




**Figure 1.5.** Automatic weather station installed at 4100 m a.s.l., 0.5 km north from the drilling site. July 2002, photo by V. Aizen

Unfortunately, accumulation gauge sensors stopped on October 18, 2001, recording only 3 months records but even this data shows that precipitated snow can be drifted out from the plateau by strong winds (**Figure 1.7**).



**Figure 1.7.** Three month accumulation records at 4150 m a.s.l. of the Plateau.



**Figure 1.6.,** Hourly air temperature. July 2002 to August 2003.

### **Long-term meteorological data**

For climatic analysis, we used long-term (50 years) meteorological data from the Akkem (49°54' N, 86°32' E; 2045 m a.s.l.), Kara–Turek (49°57' N, 86°29' E; 2600 m a.s.l.) and Aktru (°50'5 N, 87°46' E; 2110 m a.s.l.) stations located 10, 15 and 30 km north, northwest and northeast of the West Belukha snow/firn plateau. Data from these stations has the highest correlation with air temperature and precipitation in the Altai glacierized area (*Galakhov, 1981*) particularly with the West Belukha

Plateau weather station. All long-term meteorological data was checked for homogeneity, inspected for the presence of random errors, and plotted for periods with missing observations, following recommendations by *Easterling and others (1995)*. The air temperature gradient ( $T_{ds}$ ) at the drill site ( $H_{ds}$ ) was calculated from Equation (1)

$$T_{ds} = T_0 - \gamma(t) \cdot (H_{ds} - H_0), \quad (1)$$

where  $T_0$  (°C) is mean air temperature at the altitude of Akkem station ( $H_0$ );  $\gamma(T)$ , 0.0068°, 0.0072°, 0.0039° and 0.0021°C m<sup>-1</sup>, are the spring, summer, autumn and winter mean altitudinal gradients of air temperature, calculated using long-term data from the Akkem, Kara–Turek, and Aktru meteorological stations, and one year of measurements by the West Belukha Plateau weather station.

To describe atmospheric circulation patterns over southwestern Siberia that might influence regional precipitation regimes at seasonal time scales, we used monthly data of the atmospheric pressure distribution from the North Atlantic Oscillation (NAO), East Atlantic Pattern (EA), East



Atlantic/West Russia Pattern (EA/WR) and Pacific–North American (PNA) indices. The principal component scores of the patterns were obtained from <http://www.cpc.noaa.gov/data/teledoc/telecontents.html>, which were derived from rotated empirical orthogonal function analysis (Richman, 1986).

### 2003 ice-coring operation

In the summer of 2003, after two years reconnaissance and glacio-climatic monitoring in Siberian Altai, Russia two 175 m surface to bottom ice cores have been successfully recovered from the Western Belukha Plateau at elevation 4150 m a.s.l. An electro-mechanical drill with 9.5 cm diameter (inner size) and 135 cm long barrel manufactured by Geo Tecs Co. (Japan) was used for this operation. The maximum core length for one run with this drill is approximately 55 cm. It took 87.5 hours actual working time (7 working days) to drill one core. The total number of the drill runs was 325. Core sections totaled 324, with a mean length of 48.6 cm. Most core sections were not brittle and were recovered in perfect physical condition.

The 2003 expedition started on July 20 from Novosibirsk, a large industrial city in south-western Siberia. From Novosibirsk all expedition gear and personnel were transferred to Gorno



**Figure 1.8.** Snow laboratory at 4150 m a.s.l., West Belukha Plateau. Photo by V. Aizen

(Figure 1.2), we spent sorting our cargo and hiking for acclimatization. Weather was rainy and foggy, which is quite usual for this time of year in the Altai Mountains. Nevertheless, on July 25 the helicopter came and we all moved to the Plateau in several flights. It was the early morning, and a strong blizzard blowing up snow required us to move slowly and carefully windward, dragging all our supplies to the location where we had our basecamp in two previous expeditions. All day until sunset we worked hard to establish our high elevation camp at 4,150 m a.s.l. Walls made from

**Table 1.1.** Geodetic locations by measurement in 2001 (Datum: WGS84).

Location	deg N	min N	deg E	min E	Altitude	Date
Gorno-Altai Airport	51	58.246	85	50.293	339	11-Jul
Akkem Base Camp	49	54.361	86	32.518	2041	11-Jul
Karatjurek Pass	49	56.037	86	30.334	3068	13-Jul
Glacier Camp	49	48.331	86	33.722	4087	16-Jul
Pit 1	49	48.251	86	33.667	4107	19-Jul
Pit 2	49	48.258	86	33.536	4109	19-Jul
Pit 3	49	48.330	86	33.506	4113	19-Jul
Pit 4	49	48.333	86	33.652	4114	19-Jul
Pit 5	49	48.308	86	33.694	4115	19-Jul
Stake 1	49	48.286	86	33.694	4110	19-Jul
Stake 2	49	48.253	86	33.663	4101	18-Jul
Stake 3	49	48.214	86	33.632	4102	19-Jul
Stake 4	49	48.263	86	33.609	4109	19-Jul
Stake 5	49	48.242	86	33.717	4112	19-Jul

Altai, the capital of Altai Republic in the Russian Federation. A bus was rented to transport personnel about 700 km south from Novosibirsk to Gorno Altai. The expedition gear and food were transported with a large commercial truck. After one night in Gorno Altai we flew to Akkem base camp in a Russian MI-MTV commercial helicopter. Two shuttles between Gorno-Altai and Akkem base camp, near the Western Belukha Plateau were required to carry out all personnel (14 people) and about 5 tons of expedition gear, food and fuel. July 21-24 was spent in Akkem cabins preparing for operation on the Plateau. Two days of preparation at Akkem base camp, located at elevation 2,045 m a.s.l.

snow blocks were set up around each tent to prevent them from being buried in drifting snow. A snow trench measuring 2x3x5 m was dug and covered by half inch plywood to store food and kitchen supplies. Two ‘NothFace’ 5m dome tents were putt side-by-side for cooking and dining facilities. A 2.5 kW electric generator was used for lighting and drilling day and night. The next day, July 26 a large, semi-cylindrical drilling tent was set up, measuring 4.5 x 3.5 m x 2.25 m. To secure space for the drill system, the snow surface level was dug out and lowered by 70 cm. This tent is designed to withstand winds in excess of 20 m sec<sup>-1</sup>, and was additionally reinforced with lumbers and ropes in case of even stronger winds which can occur on the plateau. Another snow trench 2x2x5 m was dug in front of the drill shelter to function as a field snow laboratory, where each ice core section was weighted, described, and photographed for preliminary stratigraphic analysis (**Figure 1.8**). Finally, core sections were packed in thermo-insolated boxes to awaite transportation of the plateau. Plywood boards were used as a roof of the trench. To maintain a clean environment, the drill ten and snow laboratory were placed 200 m south of our base camp and electric generator.

Drilling started on July 28 (**Figure 1.9**). At least four people were required for the drilling operation. One person is required to operate the controls while three people maneuver and handle the drill body, which is laid horizontally after each run The drilling operation was usually done from 6 am to 12 pm and from 4 pm to 12 am.


The first ice core was drilled to a depth of 174.3 m in depth on August 4. At this depth, the cutter began to chip from hitting basal rock debris. After obtaining this core, the borehole temperature was measured every 10 m. Drilling for the second core started on



**Figure 1.10.** Cargo terminal, Seattle International Airport. Ice-core arrived. Photo by V. Aizen

August 5. The second core was drilled approximately 4 m east of the first borehole, and finished on the August 10 at the same depth.

All ice core sections were stored in 30 insulated boxes (size: L 129 x W 50 x H 50, Insulation Shipping Containers Co., U.S.A). The boxes were transported by a helicopter from the plateau to Gorno-Altaiisk Airport. From there they were carried by a commercial freezer truck to Novosibirsk International Airport, where they were stored in an airport freezer at –20 degree until transportation to Japan and USA by air cargo (**Figure 1.10**).



**Figure 1.9.** Drilling process. Daniel Joswiak, Arzhan Surazakov, Tetsuhide Yamasaki and Akiyoshi Takahashi. West Belukha Plateau. Photo by V. Aizen



**Figure 1.11.** Alexei Chebotarev with ice-core. West Belukha Plateau. Photo by V. Aizen

### Ice core properties

Most of the cores sections are not brittle and were obtained in a good cylindrical shape (**Figure 1.11**). The cores from surface to about 50 m are mixed of ice and firn layers. Deeper than 50 m, the firn becomes bubbly ice with some clear ice layers.




**Figure 1.12.** Dust layer in core section 111 at 51.4 m depth.

also contained dust layers, which are probably corresponded with the dust layers of our ice cores. These dust layers will be further analyzed discussed, and interpreted in the near future

Core sections deeper than 163 m from the surface were visibly dirty, containing silt and sand, suggesting that this ice is near the basal ice/rock interface. The core sections from 169.6 m depth contained small stones about 1 cm in diameter (**Figure 1.13**).

Borehole temperatures were measured every 10 m, the change with depth is depicted in **Figure 1.14**. Temperatures decreased with depth to 70 m, reaching the minimum of  $-15.7^{\circ}\text{C}$  at 70 m, then increased with depth to  $-14.2^{\circ}\text{C}$ .




**Figure 1.14.** The borehole temperature profile. West Belukha Plateau

Bulk density of the cores increased with depth, reaching  $0.9\text{ g/cm}^3$  at approximately 50 m. The density increased with depth in a roughly linearly fashion from 7 m to 42 m.

Four visible dust layers were observed in the ice core, excluding dust and dirty layers at bottom part the core (**Figure 1.12**). The depths of the dust layers were 51.4, 54.6, 57.8, and 122.4 m. The ice core drilled at the eastern plateau of Mt. Belukha by *Olivier et al.* (2003),



**Figure 1.13.** Loosed stones close to the bottom at 169.6 m depth.



**Figure 1.15.** Density of deep ice-core. West Belukha Plateau.


## The laboratory analysis and first results

### Stable isotope analysis

The Altai firn core 2001 and snow samples of 2001 and 2002 were processed in a dedicated cold room at NIPR using techniques established for ultra-clean sample preparation. Frozen 18 mΩ water blanks were passed through the entire system to ensure there was no contamination and for quality control. The melt index (Koerner and Fisher, 1990), compiled from the measured 5 cm density (Figures 1.15 and 1.16) and analyzed stratigraphy (Figure 1.17) showed < 5% melt.

Each 3–5 cm of the upper 11 m of the snow/firn core, as well as samples from five snow pits, fresh snow and precipitation, were analyzed for  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  at the NIPR. 10 m of the bottom section of the core were analyzed for  $\delta^2\text{H}$  at the University of Maine and for  $\delta^{18}\text{O}$  at the University of Idaho. The technique of snow/firn-core stable-isotope analysis has been described by Kreutz and others (2001).

In the clean room facilities at the Geochemical Laboratory of the National Polar Research Institute in Tokyo each snow and ice sample processed for the major ions and geochemical elements was brought to dryness (using acid-cleaned PTFE vessels) via sub-boiling evaporation inside a HEPA-filtered laminar flow box. Acid digestion of sample particulates was performed by adding a 1:3 HF:HNO<sub>3</sub> solution (Optima brand) and heating (~55°C) overnight. Thus, major and trace element data from these samples represent total sample concentration (soluble plus insoluble). Although we have not quantified the digestion efficiency of this technique, visual inspection suggests that any remaining material was largely organic. Samples were then redissolved in 1N HNO<sub>3</sub> (Optima and Milli-Q) for analysis with a



**Figure 1.16d.** Isotopic composition,  $\delta^{18}\text{O}$  (a);  $\delta^2\text{H}$  (b); snow/firn density,  $\rho$  (c); cumulative snow water equivalent,  $\Sigma\text{Ac}$  (d), in the snow/firn core. (e) Correlation between annual accumulation,  $\text{Ac}$ , records from firn/ice core with precipitation,  $\text{P}$ , at the Akkem station. (f) Distribution of annual accumulation records from firn/ice core and precipitation at the Akkem


Finnigan Element high-resolution magnetic sector inductively coupled plasma mass spectrometer (ICP-MS). The rare earth element suite was measured in low-resolution mode ( $m/\Delta m = 300$ ), while Ca, S, Al, and Fe were measured in medium resolution ( $m/\Delta m = 3000$ ). Detection limits (given as sample blank  $3\sigma$ ) for each element are given in Table 1; total sample blank values were <1% for all elements measured in each sample. The sulfur isotope composition of sample filtrate was determined after filtering through 0.4-mm pore size PTFE filters. Two contrasting firn layers were analyzed: one from a visible dust layer ( $7.61 \pm 7.80$  m WED), and one from a section of core with relatively low  $\text{SO}_4^{2-}$  concentrations representing "background" dust conditions ( $6.87 \pm 7.39$  m WED).




Each sample represents the combination of two adjacent sections of the core. The total amount of S in the samples (measured as  $\text{SO}_4^{2-}$  via ion chromatography) was 190.8 and 15.3 mg, respectively. Both samples were evaporated to a volume of 2 mL, and transferred into tin reaction vessels. Samples were pulse combusted in an elemental analyzer, and  $\text{SO}_2$  gas was separated chromatographically. The  $\text{SO}_2$  was analyzed via gas-source mass spectrometry and is reported in delta notation versus the Canyon Diablo Troilite (CDT) standard. Estimated ratio error for the two samples is  $\pm 0.05\%$ .

### Establishing a depth–age relationship

No significant post-depositional effects were apparent in the 2001 firn-core oxygen and hydrogen records, as the amplitude of the signal near the bottom of the core was similar to that observed near the surface (Figure 1.16a and b). To determine annual snow accumulation layers, we used stable-isotope records with existing, well-preserved seasonal  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  signals (Figure 1.16a and b), visible seasonal accumulation evidence (Figure 1.17) and annual precipitation data from the Akkem station (Figure 1.18b). Identification of annual accumulation layers in the snow/firn core was based on the lowest  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values. Minimum winter air temperatures are a distinctive characteristic of the



**Figure 1.18.** Monthly mean air temperature (a), precipitation (b), and monthly percentages of annual precipitation (c) at Akkem station.



**Figure 1.17.** Stratigraphy of 21 m snow/firn core recovered from the West Belukha plateau with seasonal and annual layer identification: 1 is snow with ice lenses up to 5 mm thick; 2 is fine-grained firn with 1–2 mm ice crusts; 3 is fine-grained compact white firn; 4 is fine-grained firn with aeolian particles; 5 is coarse-grained firn with ice lenses up to 8 mm thick.

Altai meteorological regime (Figure 1.18a). Snow and firn–ice densities (Figure 1.16c) were used to establish cumulative depth/water-equivalent profiles (Figure 1.16d). A significant correlation of 0.71, between stable-isotope values and the snow/firn density, allowed the identified annual layers to be verified. Maximum densities, observed during warm seasons, are associated with high isotopic values, while minimum densities, related to cold seasons, are associated

with low isotopic values. Snow/firn stratigraphy was also used to identify annual layers in the snow/firn core (Figure 1.17). The highest correlation, between annual accumulation layers in the snow/firn core and precipitation at the Akkem station, reached 0.82 (Figure 1.16e and f). The mean

annual accumulation rate, calculated at the drill site, was 690 mm from summer 1984 to summer 2001. To determine the seasonal and monthly oxygen–deuterium distribution in the snow/firn core, a relationship between snow accumulation on the Belukha plateau and precipitation at the Akkem station () was developed. The mean accumulation rate obtained agrees with the calculated accumulation rates presented in the *World Atlas of Snow and Ice Resources* (Kotlyakov, 1997).

### Seasonal snow accumulation

The monthly and seasonal snow accumulation values from summer 1984 to summer 2001 at 4115 m a.s.l. were calculated using the linear relationship obtained from the snow/firn core and

**Table 1.2.** Oxygen and deuterium isotope ratios ( $\delta^{18}\text{O}$ , ‰) in snow pits, firn/ice cores, fresh and old snow, and precipitation from the Belukha plateau, Altai and other glaciers along the northern periphery of central Asia and northern Tibet

	Snow pit 1	Snow pit 2	Snow pit 3	Snow pit 4	Ice core	Fresh snow	Old snow	Precipitation
<i>Altai, Belukha plateau, 4115 m a.s.l. (<math>\delta^{18}\text{O}</math>, ‰)</i>								
<i>n</i>	40	40	40	40	484	3	6	28
Ave	-15.5	-14.6	-15.3	-14.0	-13.6	-15.3	-23.7	-15.3
Max.	-8.8	-6.3	-8.6	-8.1	-9.1	-14.9	-22.4	-6.1
Min.	-27.9	-25.9	-25.4	-24.7	-19.9	-16.6	-25.3	-18.3
St dev.	4.8	4.8	3.7	3.9	1.9		1.3	3.9
$\Delta$	19.2	19.6	16.8	16.6	10.8	1.3	2.9	12.2
<i>Altai, Belukha plateau, 4115 m a.s.l. (<math>\delta^2\text{H}</math>, ‰)</i>								
<i>n</i>	40	40	40	40	484	3	6	28
Ave	-112.1	-105.8	-110.4	-101.7	-97.9	-101.9	-176.2	-114.3
Max.	-59.9	-48.9	-61.9	-63.5	-55.1	-101.9	-161.7	-53.7
Min.	-204.5	-194.3	-191.3	-179.6	-146.7	-115.6	-194.6	-140.8
St dev.	35.9	36.4	28.8	29.1	15.6		11.5	25.7
$\Delta$	144.6	145.4	129.4	116.1	91.6	13.7	32.9	87.1
<i>Altai (<math>\delta^{18}\text{O}</math>, ‰) (Kotlyakov and Gordienko, 1982)</i>								
Tomich glacier, 2300–2750 m a.s.l.			-16.4	-26.1				
<i>Altai (<math>\delta^{18}\text{O}</math>, ‰) (Oliver, 2003)</i>								
Belukha saddle, 4062 m a.s.l.					-12.0			
<i>Tien Shan, Inylchek glacier (<math>\delta^{18}\text{O}</math>, ‰) (Kreutz, and others, 2001; Aizen and others, in press b)</i>								
<i>n</i>	40	20	20	205	25			
Ave	-13.05	-14.18	-12.67	-16.8	-15.24			
Max.	-7.96	-9.58	-10.11	-6.0	-13.92			
Min.	-18.41	-17.74	-14.91	-35.6	-16.00			
St dev.	2.24	2.50	1.35	6.7	0.57			
$\Delta$	10.45	8.16	4.79	29.6	2.08			
<i>Pamir (<math>\delta^{18}\text{O}</math>, ‰) (Kotlyakov and Gordienko, 1982)</i>								
Abramova glacier, 4400 m a.s.l.			-10.5	-16.8				
Glacier, head Lyadjuardara river, 5200 m a.s.l.			-13.3	-27.4				
Krasnoslobodceva glacier, 5040 m a.s.l.			-10.0	-25.5				
Akbaital glacier, 5100 m a.s.l.			-18.0	-21.0				
Bakchigir glacier, 5000 m a.s.l.			-16.4	-26.1				
<i>North Tibet (<math>\delta^{18}\text{O}</math>, ‰) (Thompson and others, 1995)</i>								
Guliya ice cap					-13.1			
Dunde ice cap					-12.5			

Note: *n* is number of samples. Ave, max., min., st dev.,  $\Delta$  are average, maximum, minimum, standard deviation and amplitude.

precipitation at the Akkem station (**Figure 1.16e**), and then normalized according to *Aizen and others* (in press a). Calculated seasonal snow accumulation values were verified using the snow/firn stratigraphic profile (**Figure 1.17**), where winter layers could be differentiated from others by their homogenous bright-white crystal structure. In the spring layers, there are signs of 1–2 mm slender

radiation crusts, while summer layers have ice lenses up to 8 mm thick. The autumn layers could be identified by more compact firn and yellow-brown aeolian particles. Uncertainty in calculating seasonal accumulation using the above relationship (**Figure 1.16e**) was less than  $\pm 10\%$  of the seasonal accumulation rate obtained from the snow/firn core stratigraphic profile. The corresponding annual seasonal  $\delta^{18}\text{O}_{\text{season}}$  and  $\delta^2\text{H}_{\text{season}}$  compositions were averaged as arithmetic means of every 3–5 cm measured stable-isotope ratio in each seasonal accumulation layer of the snow/firn core. The same approaches have been used by Aizen and others (in press a), *Barlow and others* (1993), *Shuman and others* (1995) and *Yao and others* (1999) to evaluate seasonal accumulation in snow/firn/ice cores.

### **Stable isotopes from firn cores and snow pits**

The mean oxygen-isotope ratio from the Altai core of  $-13.6\text{‰}$  is in accordance with oxygen-isotope records obtained from central Asia and the northern Tibetan mountains (**Table 1.2**). The higher average annual air temperature at the Altai drill site, at a lower altitude than the Tien Shan drill site (*Aizen and others*, in press a), corresponds to a slightly higher mean oxygen-isotope ratio than the  $-16.8\text{‰}$  measured in the Inilchek glacier core, in the Tien Shan. For the Altai firn core, the highest  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  seasonal accumulation variability was observed in winter firn layers (**Table 1.3a**). The annual variation in  $\delta^{18}\text{O}$  from the snow pit and firn core (19.6 and 10.8‰) was about the same as that from the Pamir and other Altai glaciers (**Table 1.2**). The Tien Shan glaciers exhibit a greater variation in  $\delta^{18}\text{O}$  (29.6‰) than other Asian sites (**Table 1.2**) which is attributed to the lowest winter


**Table 1.3.** Belukha plateau snow/firn core monthly average  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  composition and their linear relationship

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Nov	Dec
<b>(a) Monthly average <math>\delta^{18}\text{O}</math> and <math>\delta^2\text{H}</math> composition</b>												
	$\delta^{18}\text{O}$											
Ave	-16.9	-16.8	-16.7	-15.5	-13.8	-12.9	-13.1	-13.1	-13.9	-14.2	-15.9	-16.3
Max.	-14	-14	-14	-12.7	-10.8	-9.1	-10.3	-10.1	-9.8	-8.1	-10	-10
Min.	-24.7	-22.8	-21	-19.9	-17.4	-17	-17.9	-19.1	-20.1	-19.5	-19.9	-23.6
St dev.	2.6	2.3	1.9	1.7	1.3	1.9	1.5	1.6	2.1	2.6	2.4	2.9
	$\delta^2\text{H}$											
Ave	-122.7	-122	-120.5	-110.8	-99.6	-92.1	-94.3	-95	-101.1	-102.3	-113.1	-115.7
Max.	-93.5	-93.5	-93.5	-88.6	-74.4	-55.4	-71.6	-72	-68.9	-55.1	-55.1	-55.1
Min.	-179.6	-168	-151.4	-149.9	-128.5	-123.6	-134	-139.7	-149.3	-153.9	-149.9	-176.5
St dev.	21.3	19.5	16.5	14.7	10.9	15.3	12.1	12.8	16.8	21.9	22.2	26.2
<b>(b) Linear relationship of <math>\delta^{18}\text{O}</math> and <math>\delta^2\text{H}</math> composition</b>												
<i>n</i>	17	17	18	31	56	105	101	107	49	32	17	16
<i>a</i>	7.8	8	8	8	7.8	7.9	7.8	7.9	8.3	8.1	8.1	8.3
<i>b</i>	8.6	12.7	13.3	13.7	8.6	10.9	7.9	8.8	7.8	13	13.3	18.1
$R^2$	0.92	0.91	0.88	0.87	0.91	0.94	0.93	0.92	0.97	0.94	0.9	0.95

Note: *n* is number of measurements,  $R^2$  are coefficient of determination.

isotopic means.

Higher winter minimum  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values in the Altai core ( $-19.9\text{‰}$ ) than in the Tien Shan cores ( $-35.6\text{‰}$ ), caused by higher winter air temperatures at the Altai site and more short trajectories of air masses bringing precipitation there, demonstrate the continental effect of meteoric water depletion moving farther from the source of water vapor (*Dansgaard*, 1964; *Friedman and others*, 1964). During the cold season, air masses are moving eastwards across the Eurasian continent along the main route: high latitudes from Iceland to western Siberia and then to central Asia (*Glukh and Kononova*, 1982). Air masses with water vapor precipitated over the Tien Shan have therefore been subject to a stronger depletion of  $\delta^{18}\text{O}$ .



**Figure 1.19.** Deuterium ( $\delta^2\text{H}$ ), oxygen ( $\delta^{18}\text{O}$ ), snow/firn density ( $\rho$ ) and snow temperature  $T$  ( $^{\circ}\text{C}$ ) measurements in the West Belukha plateau snow pits, 2001.

$$\delta^2\text{H} = 8.0 \delta^{18}\text{O} + 10 \quad (2)$$

Similar slopes observed in local relationships (**Figure 1.21a**), and in the GMWL, indicate the same initial relationship of fraction factors and point to the absence of strong melt and percolation in the snow/firn core and snow pits. The  $\delta^2\text{H}$ – $\delta^{18}\text{O}$  relationships related to June accumulation layers in the snow/firn core records (**Table 1.3b**) are closest to the GMWL. The smaller or larger intercept in the local relationship, from the snow/firn core (**Table 3b**), fresh snow, snow pit and precipitation, than in the GMWL ( ) reflects different kinetic evaporation effects on the water vapor transferred to the Altai mountains, e.g. initial water vapor was quickly or slowly evaporated under non-equilibrium conditions (*Kendall and McDonnell, 1998*).

The highest seasonal mean  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  in the Altai (**Table 1.3a**) and Tien Shan (*Aizen and others, 2004a*) snow/firn cores were for the summer accumulation layers. However during the warm season, the air masses are crossing the Eurasian continent eastward along two main routes: high latitudes from Iceland to western Siberia and mid- (even low-) latitudes to middle Asia (*Glukh and Kononova, 1982*). Altai glaciers receive moisture from higher latitudes than do the Tien Shan glaciers, where water vapor is transferred from lower latitudes and altitudes. Hence, the maximum  $\delta^{18}\text{O}$  values in the Altai cores are slightly lower than those observed in the summer layers of the Tien Shan core (**Table 1.2**).

Oxygen- and hydrogen-isotope ratios from Altai snow pits (**Figure 1.19**) varied significantly; the lowest values (**Figure 1.16 top**) corresponding to the lowest minimum air temperature and an abnormally high amount of winter precipitation observed in winter 2000/01 (**Figure 1.18**). The extremely low hydrogen- and oxygen-isotope ratios ( $-28$  and  $-200\text{‰}$ ) in winter 2000/01 were caused by low air temperatures and by a significant amount of direct precipitation not enriched by evaporation.

#### **Relationship between stable hydrogen- and oxygen-isotope ratios**

The relationship in the precipitation, snow pits and snow/firn core from the Belukha plateau (**Figure 1.20**) has the same slope to the covariance (i.e. 8) as that of the global meteoric water line (GMWL, Equation (2)) described by *Craig (1961)*:




**Clustering precipitation transferred from external (oceanic) and internal (continental) moisture sources**

The equations corresponding to evaporation from the ocean under conditions close to equilibrium were obtained by *Dansgaard and Tauber (1969)*. They were based on precipitation data from 38 island and continental stations around the North Atlantic (Equations (3) and 4)):

$$\delta^2\text{H} = 8.1 (\pm 0.1) \delta^{18}\text{O} + 11 (\pm 1) \quad (3)$$

$$\delta^{18}\text{O} = 0.69 T - 13.6 \quad (4)$$

To distinguish between oceanic moisture and water vapor evaporated from internal drainage basins (e.g. local basins or the Aral–Caspian basin) and transferred to the northeastern latitudes of Asia, the  $\delta^2\text{H}$ – $\delta^{18}\text{O}$  isotopic relationships from the Altai snow/firm core were clustered



**Figure 1.20.** Stable deuterium ( $\delta^2\text{H}$ ) and oxygen-isotope ( $\delta^{18}\text{O}$ ) relationship in snow pit, fresh snow, precipitation and snow/firm core on the West Belukha plateau, 2001.

into three distinct datasets (**Figure 1.21a**). The clustering procedure was based on inequalities (5–7) resulting from Equation (3).

$$\text{“Atlantic”}: 8.2 \delta^{18}\text{O} + 10 \leq \delta^2\text{H} \leq 8.0 \delta^{18}\text{O} + 12 \quad (5)$$

$$\text{“Oceanic”}: \delta^2\text{H} < 8.2 \delta^{18}\text{O} + 10 \quad (6)$$


$$\text{“Internal”}: \delta^2\text{H} > 8.0 \delta^{18}\text{O} + 12 \quad (7)$$

where  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  are the isotopic contents in 254 samples from the snow/firm core.

The deuterium excess ( $d$ , ‰) was also defined by  $d = \delta^2\text{H} - 8\delta^{18}\text{O}$  (*Dansgaard, 1964*). Precipitation from the Altai snow/firm-core records, having a hydrogen- and oxygen-isotope relationship based on the conditions of inequality (5) (the same as Equation (3)), was considered to have been transferred from the northern part of the Atlantic Ocean. A smaller intercept in the snow records (inequality (6)) reflects a smaller kinetic effect during evaporation, when conditions were closer to equilibrium than in the records corresponding to Equation (3). Precipitation with this record also originated from oceanic water. Precipitation with a hydrogen- and oxygen-isotope relationship based on the conditions of inequality (7) was more enriched in deuterium than in records corresponding to Equation (3). High deuterium excess indicated strong kinetic effects which were most probably caused by water evaporation from internal moisture sources, e.g. Aral and Caspian basins or convection from a local basin.

Total snow/firm accumulation in the Altai core from June 1984 to July 2001 amounted to 12,591 mm w.e., 69% corresponding to inequalities (5) and (6), i.e. precipitation transferred from external, oceanic moisture sources, and 31% to inequality (7), i.e. precipitation transferred from internal moisture sources (Fig. 10b). There is no significant dependence of deuterium excess on oxygen content (Fig. 10c).

*Internal moisture source (inequality (7))*



**Figure 1.21.** (a) Annual and seasonal clustered stable  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  relationships. (b) Annual and seasonal contributions to accumulation of precipitation originating from external and internal moisture sources: percentage of average oxygen, deuterium and deuterium excess. (c) Seasonal relationship between deuterium excess and oxygen isotopic composition (d) in snow/firn core from the West Belukha plateau, 2001.

The records associated with internal moisture sources have the highest deuterium excess (15.04‰, **Figure 1.21b**) as a result of low air humidity at the center of the continent during water vapor evaporation from local moisture sources.

The smallest share of precipitation transferred from the internal water cycle to the Altai glaciers was observed in winter (27%, **Figure 1.21b**), when the Siberian High is strongest over northern and central Asia, blocking any intrusion of air masses, and conditions for inland evaporation and local convection are weakest because of low continental heating. The mean deuterium excess is highest in winter (up to 16.7‰, **Fig. 1.21b**), when air humidity during

evaporation over inland moisture sources is lowest. Absolute summer deuterium excess reaches 23‰ (**Figure 1.21c**), indicating a strong kinetic effect during evaporation, when air humidity is low over the inland moisture flow located at the center of Eurasia. This is in accordance with results from *Froehlich and others* (2002) on the maximum mean of deuterium excess obtained from stations in the Northern Hemisphere.

In spring, warming of the inner continental regions occurs faster than in the coastal regions, which increases inland evaporation. Therefore, the largest annual proportion of precipitation is brought to the Altai from internal moisture sources in spring (42%, **Figure 1.21b**). Among the seasonal records, the mean intercept in the oxygen-hydrogen relationship and deuterium excess are lowest in spring (**Figure 1.21b**).

In summer and autumn, the contribution of internal continental water vapor to accumulation on central Asian glaciers is also high (up to 31%). Absolute summer deuterium excess reaches 25‰ (Fig. 10c) indicating low air humidity over the inland moisture flow.

#### ***Atlantic moisture source*** (inequality (5))

Moisture with a hydrogen- and oxygen-isotope relationship corresponding to water vapor evaporated over the northern Atlantic Ocean ( $\delta^{18}\text{O} = 8\delta^2\text{H} + 9.83$ , **Figure 1.21a**) comprises more than half the annual accumulation (53%, **Figure 1.21b**). The  $\delta^{18}\text{O}$ – $\delta^2\text{H}$  relationship varied slightly during a year (**Figure 1.21a**). Deuterium excess during a year was almost invariable, ranging from 9.7–10‰ (**Figure 1.21b**). The mean values of the  $\delta^{18}\text{O}$  records related to precipitation origin, from internal and Atlantic Ocean sources, are about the same during all seasons (Fig. 10b). The range of variability in their absolute values is also similar, fluctuating from –9 to –25‰ (**Figure 1.21c**). During the winter,  $\delta^{18}\text{O}$  records associated with inter-land moisture sources are even lower than  $\delta^{18}\text{O}$  records related to North Atlantic water vapor origin. Water vapor evaporated from the southern internal moisture sources in winter, probably moves northward over the cold Asian continent and mixes with cool air masses, accounting for stronger depletion in  $\delta^{18}\text{O}$  than water vapor evaporated from the relatively warmer ocean.

#### ***Oceanic moisture source*** (inequality (6))

Precipitation transferred from an external moisture source, with the smallest deuterium excess, comprises on average 16% of the annual accumulation with a summer maximum (up to 21%, **Figure 1.21b**) and spring minimum (8%). The lowest mean d-excess (4.6‰) was observed in autumn records; most probably associated with the highest atmospheric moisture saturation over the ocean during water vapor evaporation, further moisture transportation and precipitation over the Altai. The mean and absolute values of  $\delta^{18}\text{O}$ , from accumulation layers (**Figure 1.21b** and **c**) having an “oceanic” moisture origin, are the highest among the three considered clusters, ranging from –9‰ to –15‰. The highest  $\delta^{18}\text{O}$  values may be associated with the highest air temperature during precipitation (*Aizen and others*, 1996; *Lin and others*, 1995; *Yao and Thompson*, 1992; *Yao and others*, 1995) and/or near-by water-vapor formation, e.g. Pacific Ocean.

#### ***Relationship between $\delta^{18}\text{O}$ and air temperature*** (i.e. transfer function)

Multiple picks (**Figure 1.16**) with high  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  isotopic relationships in annual variations of stable-isotope records were the result of: (1) summer maximum air temperature; (2) several sources of moisture including oceanic and inter-land, partially during spring, summer and autumn, and; (3) different trajectories by which air masses were transporting moisture to the Altai mountains.

#### ***Transfer functions and continental/oceanic moisture sources***

The transfer functions (**Figure 1.22a**) were developed for three considered clusters of precipitation origin (section 6.1) using monthly data on air temperatures at the drill site (Equation (1)) and isotopic ratios averaged for seasonal accumulation snow/firn layers (section 5.2). Annual and seasonal air temperatures for each cluster were averaged from the monthly means, if the records related to a considered cluster were observed in these months. Transfer functions (Equations (8–10))


inside each cluster differed insignificantly (range of variation in parentheses) during all seasons (**Figure 1.22a**):

$$\text{“Atlantic”}: \delta^{18}\text{O} = 0.6 T - 13.0 (\pm 0.4) \quad (8)$$

$$\text{“Oceanic”}: \delta^{18}\text{O} = 0.85 (\pm 0.05) T - 10.2 (\pm 0.4) \quad (9)$$

$$\text{“Internal”}: \delta^{18}\text{O} = 1.05 (\pm 0.05) T - 10.0 (\pm 0.6). \quad (10)$$

The most similar relationship, to the co-variance found in precipitation from the North Atlantic (Equation 4), was observed during all seasons in the records clustered under the condition of inequality (5) (Equation 8), pointing to North Atlantic precipitation. Under the same air temperatures, the most negative  $\delta^{18}\text{O}$  values are associated with precipitation from the North Atlantic, while the heaviest values were associated with moisture transferred from an oceanic source (Fig. 11a) that could be the Pacific or Arctic Oceans. Considering that relatively heavy  $\delta^{18}\text{O}$  isotopic ratios are characterized by higher temperatures, we expect this precipitation origin was the Pacific not the Arctic Ocean.



**Figure 1.22** Seasonal relationships between  $\delta^{18}\text{O}$ , content in snow/firn cores and air temperature at 4115 m a.s.l. depending on (a) sources of moisture and (b) prevailing large-scale atmospheric patterns.

The transfer functions developed for precipitation from the inter-land moisture sources lie between the two functions developed for oceanic originated precipitation (**Figure 1.22a**).

The different isotope–temperature slopes (i.e.  $0.6\text{‰}$ ,  $0.85\text{‰}$  and  $1.05\text{‰ }^{\circ}\text{C}^{-1}$ ) in the transfer functions (Equations (8–10)) are the result of a different intensity in the enrichment of snowfall by heavy isotopes through snow evaporation and sublimation. With “Internal” air masses (Equation (10)), low humidity, insignificant amounts of precipitation and snow/wind storms caused intensive evaporation and sublimation of falling snowflakes. “Atlantic” air masses (Equation (8)) are associated with intensive snowfalls, high humidity and relatively windless weather. With these, much of the falling snow was not enriched by evaporation and sublimation.

### ***Transfer functions and large-scale atmospheric patterns***

To verify different sources of moisture nourishing the glacier being studied, we compared mean values of  $\delta^{18}\text{O}$  in snow/firn core seasonal accumulation layers with seasonal mean air temperatures at the drill site (Equation (1)), considering the indices of atmospheric circulation patterns. The sign of the seasonal EA, EA/WR, NAO and PNA values differentiated these relationships (**Figure 1.22b**). The negative spring values of EA and positive summer values of the NAO and EA/WRP indices are associated with moisture brought from the west (i.e. Atlantic Ocean, Mediterranean and Black Seas) to the Siberian Altai. Positive values of PNA indices are associated with moisture brought from the Pacific.

$$\text{'NAO, EA/WRP} > 0; \text{EA} < 0\text{'}: \delta^{18}\text{O} = 0.6 T - 12.1 (\pm 0.1) \quad (11)$$

$$\text{'PNA} > 0\text{'}: \delta^{18}\text{O} = 0.85 (\pm 0.05) T - 10.3 (\pm 1.0) \quad (12)$$

The relationships based on records associated with Atlantic moisture sources (Equation (8), **Figure 1.22a**) are most similar to those that considered the NAO, EA/WRP indices (Equation (11), **Figure 1.22b**). Furthermore, linear transfer functions associated with an oceanic moisture source (Equation (9), **Figure 1.22a**) are comparable with equations relating  $\delta^{18}\text{O}$  and air temperature during the seasons when the PNA indices had positive values (Equation (12), **Figure 1.22b**).

### **Findings**

Analysis of the oxygen–deuterium content in a 21 m snow/firn core allowed us to estimate the annual/seasonal amounts and origins of precipitation nourishing the Belukha plateau region of the Siberian Altai.

Annual accumulation was found to be about 700 mm w.e. at 4115 m a.s.l. for the period from summer 1984 to summer 2001. This is in agreement with the meteorological and snow/firn stratigraphy data.

The mean oxygen isotope ratio of  $-13.6\text{‰}$ , as well as an annual  $\delta^{18}\text{O}$  variation range of  $19.6\text{‰}$ , from the Altai core is in accordance with oxygen-isotope records obtained from central Asia and the northern Tibetan mountains. The slopes in the hydrogen and oxygen relationship (i.e. 8), in the precipitation, snow pits and snow/firn core and d-excess of  $10.63\text{‰}$ , are similar to slopes and d-excess in the GMWL co-variance. However, the clustering analysis of the oxygen–deuterium relationships indicated that both Atlantic and Pacific Oceans and inner-continental water are the sources of moisture to the Siberian Altai. The 17 year snow/firn accumulation, recorded in the 21 m core, comprised 69% of the precipitation transferred from the oceans. In the annual accumulation, more than half was from water vapor evaporated over the northern Atlantic Ocean. The contribution of precipitation transferred from the Pacific Ocean varied from 8% in spring to 21% in summer, when the lowest d-excess ( $4.6\text{‰}$ ) was associated with high atmospheric moisture saturation over the ocean during water-vapor evaporation. Among the three clusters, the highest mean and absolute  $\delta^{18}\text{O}$  values were related to precipitation originating from the Pacific and were associated with relatively higher air temperature during snowfall. The inter-land moisture sources contributed 31% of annual precipitation, with the spring maximum reaching 42%. Low humidity at the center of the continent resulted in the highest d-excess (up to  $25\text{‰}$ ) in the records associated with inter-land moisture sources.

The linear transfer functions, between air temperature and isotopic composition in accumulated precipitation from both external and internal moisture sources, were developed at the seasonal temporal scale. Under the same air temperatures, the lightest  $\delta^{18}\text{O}$  values are associated with precipitation from the Atlantic Ocean, while the heaviest values are associated with that from the Pacific Ocean. Low humidity and snow/wind storms caused significant evaporation and sublimation of falling snowflakes and a bias on exceeding the isotope–temperature slope when air masses transferred moisture from “Internal moisture sources” rather than when air masses originated over the Atlantic. The relationships based on records associated with Atlantic moisture sources are most similar to those which considered the NAO, EA, EA/WRP indices, whereas linear

relationships associated with the Pacific moisture origin are comparable with equations based on records when the PNA indices had positive values.


According to identification of annual layers in the cores, the mean annual snow accumulation for the period from 1997 to 2001 was found to be about 800 mm. Analysis of large-scale atmospheric patterns (NAO and PNA), oxygen and deuterium isotope content in ice core and snow pits assume two sources of moisture on the Altai Mountains.

#### **Further proposed laboratory and analytical research (Altai)**

1. complete deep ice-core (175 m) analysis for major ions, trace elements, stable isotopes ( $^{18}\text{O}$  and  $^2\text{H}$ ), Green House Gases (GHG), black carbon, radionuclide, heavy metals
2. ice-core data calibration and validation
3. data interpretation and simulation
4. technical reports and manuscripts preparation for publication in peer review journals

Necessary time: **2 years (2004-2005)**

## PART II. SOUTH-EASTERN TIBET AND HIMALAYA



**Figure 2.1.** Map Bomi glacierized region and Zuoqiupu Glacier, South-Eastern Tibet-Himalayas.

### Overview

The Bomi mountains are the largest glacier covered area (approximately 28,000 km<sup>2</sup>) in the low latitudes of the Northern Hemisphere (29°30'N; 97°00'E) with the highest peaks over 7,700 m a.s.l. (**Figure 2.1**). The glacial accumulation areas elevated above 5,000 m where glaciers, ice thicknesses are greater than 300 m (Aizen, 1994) are suitable for the recovery of ice core records. The location of this area between Tibetan and Himalayas mountain ranges, on the southeastern margin of the Tibetan Plateau provide an excellent opportunity to

develop climatic records relating to the major circulation systems such as the tropical and subtropical monsoons, Tibetan high pressure system and the westerly jet stream. Furthermore, the

Bomi region, as the montane barriers to southeastern air masses moving across the Tibetan Plateau, plays an important role in determining the climatic processes in the southern periphery of Asia. In addition, the Bomi glaciated area holds the greatest concentration of snow and ice in the low latitudes and constitute a source of fresh water shared by ten very populated countries. Bomi glaciers feed several major Asian rivers: Yangtze, Mekong, Hengduan, Brahmaputra, and play a very important role in the water cycle in this region. For example, glaciers of Gongga Shan (another glaciated area in south-eastern Tibet) tend to trap, accumulate, and transfer about 0.24% atmospheric moisture from the external water cycle returning 0.30% of water per year to the Pacific and Indian oceans. The 0.06% of this external moisture is taken from accumulated glacial resources of this area (Aizen & Aizen, 1997). In contrast to the relatively short-term records of the chemical composition of wet deposition provided by the National Atmospheric Deposition Program and Hydrological BenchMark data bases, glacier ice in low latitudes provides a much longer record of wet and dry deposition (Naftz, et al., 1991). A long-term record of atmospheric deposition in the inhabited Tibetan glacial area is desirable to evaluate the possible relationship between anthropogenic emissions and chemical composition of wet and dry deposition. Furthermore, there exists a network of hydro-meteorological stations that have been in operation for the past 40-50 years in the south-eastern Tibet and Himalaya that provide sufficient data for detailed climatological analyses as well as providing data that can be used to compare and calibrate ice core records.



**Figure 2.2.** Up and down through Tibet to Bomi glaciers. Our expedition truck is going forward. Photo by V. Aizen



### **Field itinerary**

Our first field reconnaissance in south-eastern Tibet-Himalaya started on September 24 from



**Figure 2.3.** Zuoqiupu Glacier, passage to snow-firn plateau, 5800 m a.s.l.  
Photo by V. Aizen

Chengdu the capital of Sichuan Province. Two ‘Toyota’ jeeps and one 6 ton truck (similar to 1944 Studebaker) loaded with expedition gears and food moved to Bomi through Luding, Kagding, Batang and Baxoi. Six days we traveled by unpaved alpine roads up and down from one pass to another ascending over 5,500 m elevation. On September 30 we finally arrived to Bomi village. The next day we established our base camp near Youlong Glacier, the biggest glacier (32 km long) in the Bomi glaciated area. After three days acclimating around base camp (4,500 m

a.s.l.) and working on installing one of the ‘Campbell’ automatic weather stations, we moved to the Zuoqiupu Glacier. The 15 yaks and 20 porters with expedition gears and food went ahead from base camp. The first day we ascended to 5,800 m a.s.l. and the following day installed the second ‘Campbell’ automatic weather station for annual meteorological measurements. After one day, Dr. Stanislav Nikitin, Daniel Joswiak and I took alpine ropes, ice-axes, ice screws, crampons, radio-telephone, and our personal backpacks and went to the glacier to scout best way trough the ice-fall. After several hours we crossed ice-fall, marked the crevasses with red colored wig-wags and screwed down five 80 m alpine ropes for our porters at the most dangerous spots. This day the first portion of our gear and food was transferred to our research site we found at 6,120 m a.s.l. on the upper part of the snow-firn plateau.



**Figure 2.4.** Tibetan porters carrying ice-core in termo-insolated boxes, Zuoqiupu Glacier, 6100 m a.s.l. Photo by V. Aizen

The next day, Tibetan porters moved all our cargo to the advance camp. One week was spent on the plateau accomplishing our research goal, which included ice thickness radio-echo sounding measurements, collecting snow samples, digging snow pits and recovering a 14 m shallow firn core. After we finished our scheduled work we returned to base camp at 4,500 m and continued collecting




stream samples and measuring ice thickness at the Zuoqiupu Glacier terminus area. The 14 m firn core and snow samples were transported down to base camp by our porters in thermo-insolated boxes. At base camp, all boxes were placed on the truck into four chest freezers powered by a mini-electric generator. On October 16 heavy snow forced us to retreat from the alpine area. Four days we then had to drive west to Lhasa along the Brahmaputra River tributaries. The east route was closed, winter had come to Tibet. The jeeps and truck returned to Lanzhou Institute via the Tibetan highway (5 days), we flew from Lhasa to Lanzhou, then to Seattle through Beijing. Firn core and snow samples were transported from Lanzhou to Seattle and the University of Idaho freezer.

## Field results

### Zuoqiupu Glacier radio-echo sounding measurements

Detailed radio-echo sounding measurements were completed at over 100 sites throughout at the accumulation area and glacier terminus in order to determine ice thickness (Figure 2.5). These measurements

were accomplished using a light-weight ice-penetrating radar system (30° direction of aeri-als, 700 MHz frequency, 10 Watt impulse, 50 NSC duration of impulse, -130 Decibel sensitivity of receiving signal, 1-2% error of measurement). Measured ice thickness range from 100 to 150 m, with a depth of 132 m at the shallow firn core site. Several intermediate horizons were apparent in the radio echo sounding reflections that may be associated with the presence of fine-grained sediment of eolian genesis. The radio-echo thickness measurements were surveyed using a GPS tool to coordinate each point of measurement. The recorded GPS measurements were converted from the World Geodetic System 1984 (WGS-84) to the topographic map system (Hvostov, V.V., 1990) using



**Figure 2.5.** Zuoqiupu Glacier, automatic weather stations, snow pits, shallow core sites (a) and the glacier ice thickness measured by ice-penetrated radar (b).

Helmet's parameters calibrated for Zuoqiupu glacier basin. The accuracy of GPS in an independent regime is about 20 m, the accuracy of our converted coordinates is about 3 m, quite high compared to the accuracy of the Chinese topographic map. The Helmet's transformation implements a conversion from one coordinate system (X,Y,Z) to another using seven parameters: - zero coordinates linear displacement (3 parameters), angle of orientation components (3 parameters), and the scaling coefficient (1 parameter). The orthogonal coordinates of the radio-echo sounding points which were not referenced by GPS were calculated by two methods: - linear interpolation, when points are located in longitudinal section between the points referenced by GPS, or through

extrapolation by the angle of cross section profile when each point (control stake) is not more than 50 m from the previous point. The altitudes of radio-echo sounding points on the glacier surface were calculated using Digital Elevation Model (DEM). A Triangulated Irregular Network (TIN) has been developed for the glacier bedrock topography (187 triangles based on 119 echo records).

***Meteorological observations***

Two ‘Campbell Scientific’ automatic weather stations were installed to record hourly measurements in the Zuoqiupu Glacier basin for one year. One station is near the terminus (4,800 m a.s.l.) and the other at the glacier equilibrium level (5,800 m a.s.l.) (**Figure 2.7**). Both stations were assembled to record air temperature, air humidity, wind speed and direction, barometric pressure, solar radiation (short and long wave) and the snow/rain amounts. The stations worked very well through the year 2002/2003. Data was successfully retrieved from the automatic weather stations in October 2003 by Daniel Joswiak, Arjhan Surazakov and Young Han. Using a laptop computer to connect to the CR10X datalogger on the automated weather stations, the hourly records were downloaded and the memory was cleared to allow for another year of recording.

***Shallow firn core and snow sampling***

During three weeks work in the Bomi glacial area, fresh snow and stream water samples were collected in pre-cleaned 200 ml plastic bottles and placed in chest freezers. Snow from six 3-3.5 m snow pits was collected every 5 cm deep, and detailed snow stratigraphy was described and recorded. Density samples were taken every 3-5 cm in snow pits using a 100 cm<sup>3</sup> stainless steel sampler and electronic balance. Temperature was measured by an electronic sensor each 10 cm. A firn core (14 m long and 7.5 cm in diameter) was extracted at an elevation of 6120 m, where the results of radio-echo sounding suggests an ice depth of



**Figure 2.6.** ‘Mini-Felix’ sun powered electro-mechanical drill. Drill barrel cleaning after each run. Zuoqiupu Glacier, 6,120 m a.s.l proposed deep drilling site. Photo by V. Aizen



**Figure 2.7.** ‘Campbell Scientific’ automatic weather station established at 5,800 m, 2 km northeast from proposed drilling site at Zuoqiupu Glacier snow-firn area. Photo by D. Joswiak



**Figure 2.8.** ‘Mini-Felix’ sun powered electro-mechanical drill. Zuoqiupu Glacier, 6,120 m a.s.l proposed deep drilling site. Photo by V. Aizen

about 132 m (**Figure 2.5**). The core was drilled from the bottom of the 3.0 cm snow pit (number 2)

with a 'Mini-Felix' (Swiss made) solar powered electro-mechanical drill (**Figure 2.6 and 2.8**). The diameter, length and weight of each recovered core section were measured for density calculations. Firn-core sections, sealed in pre-cleaned polyethylene bags, were packed in insulated shipping containers and delivered to the University of Idaho freezer for further processing and analysis. Samples from firn core, snow pits, fresh snow, and streams were analyzed for  $^{18}\text{O}$ ,  $^2\text{H}$ , major ions and trace elements. To prevent contamination, the outer portion of each core section was removed in a laminar flow clean bench using sterilized stainless blades.


### Laboratory analyses

Core and snow samples from Zuoqiupu Glacier were melted at room temperature on a laminar flow horizontal clean bench at the U of I. Ultra-clean sample preparation techniques previously described by Kreutz and others (2000) were used to prepare liquid firn samples from the 10 cm sections.

Major ion analyses were performed at the U of I and WSU geochemical laboratories. Anions were determined using a Dionex AI-450 Ion Chromatograph, and cation concentrations were determined by inductively coupled plasma- atomic emission spectroscopy (ICP-AES) with an axial-view torch (Perkin Elmer Optima 3000 XL) in the Department of Geological Sciences at the University of Idaho. Heavy metals and REE concentrations were determined at the Washington State University ICP-MS laboratory using a ThermoFinnigan Element2 Inductively Coupled Plasma-Mass Spectrometer (ICP-MS). Calibration standards for each analysis were made from certified standard acquired from Aldrich Chemical Company and Fisher Scientific. Sets of three to five different concentrations of standards were used to establish calibration curves, with correlation coefficients better than 0.999. Blanks and standards of known concentrations were run periodically throughout the analyses as quality-control checks.

Stable isotope ( $\delta^{18}\text{O}$ ) analysis was performed at the UofI Stable Isotope Laboratory using a Finnigan Delta plus isotope ratio mass spectrometer (IRMS) coupled with Finnigan's GasBench II. The GasBench injects from the headspace five times per vial, and we use the mean of the five  $\text{CO}_2$  peaks. The GasBench flush-fills the headspace of each vial with  $\sim 3000$  ppm  $\text{CO}_2$  in helium for 4 minutes to turn over this gas volume completely. This  $\text{CO}_2$  equilibrates with the oxygen in the water (0.5 mL) over 18 hours at  $26^\circ\text{C}$ . After equilibration, the  $\delta^{18}\text{O}$  of the  $\text{CO}_2$  in this classical "headspace" experiment is almost the same as the water's value. The details of the actual fractionation during equilibration, such as a temperature-based coefficient of fractionation which indicates the amount that the  $\text{CO}_2$  does not match the water's isotopic signature, are not needed since several vials of water of known isotopic composition are run along with each melted firn sample through the identical treatment principle, with the same adjustment applied to each sample within the batch.

### Discussion



**Figure 2. 9.** Anion concentration in snow at different Asian sites.

Small concentrations of  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$  and  $\text{SO}_4^{2-}$  are dominate in the fall and

Located in South-Eastern Tibet, the Zuoqiupu Glacier area has extremely low values of anions (**Figure 2.9**) compared to other Asian sites (Tien Shan and Altai) associated with the greatest amounts of precipitation (**Figure 2.10**), close proximity to moisture sources (Indian Ocean), and location far from terrestrial dust inputs. The low  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  concentrations point to minimal effects from anthropogenic impact in the most recent record recovered from the shallow firn-core. The elevated Cl levels among other anions (**Figure 2.9**) is related to the close proximity to a marine moisture source.

winter. Maximum concentrations of Na and Cl were observed in the spring and summer samples (Fig. 11), during monsoon periods of maximum precipitation.

The firn-core recovered at Zuoqiupu Glacier shows well-preserved annual isotopic variation. The low annual range (8.8‰) (Figure 2.11) results from the warmer monsoon climatic conditions and short transport distance of source moisture. The mean and absolute (maximum and minimum) values are slightly lower than from Altai core, although moisture source transport distance is less.




Figure 2.10. Mean monthly annual precipitation adjusted to the drill sites.




Figure 2.11. South-East Tibet 14 m shallow core (6120 m a.s.l.)

The meteorological records from two automatic weather stations in the Zuoqiupu glacier basin were obtained in November 2003, after one year of recording. This data will be analyzed this year with other long-term meteorological and synoptic data we collected in China during our reconnaissance.

The GPS satellite grounded survey at the terminus of two glaciers in the Bomi area was accomplished in November 2003 and will required additional measurements in the Fall of 2004 in order to understand glacier-ice velocity and link GPS coordinated points with high resolution satellite images. The glaciers in the Bomi area have been retreating at extraordinary rates for the last 20-30 years and may disappear in the near future. The climate and glacier regimes in the Bomi area are absolutely unknown, even though they are one of the most interesting and important phenomena in the northern hemisphere tropics. We are the first geo-scientists visiting the Bomi region to study climate and glaciers, and our goal is to document glacier distribution and couple glacier surface and volume changes with ice-core paleo-climatic reconstructions. We are planning to accomplish our research goals in this region within the next few years.

### PART III. The U of I new ICE-CORE LABORATORY and FREEZER


In order to the process ice-cores and snow/firn samples at the University of Idaho, we have begun to build a dedicated cold ice-core processing laboratory, ice-core storage freezer and ice-core analytical laboratory for ice-stratigraphy and stable isotope analysis. The reason for this effort is to process data quickly and efficiently, to further develop our professional skills and atmosphere at the UofI, and to teach students to work in both field and laboratory environments. Unfortunately, the Moscow-Pullman area with two academic institutions (UofI and WSU) doesn't have the necessary analytical laboratories we require. The geochemical laboratories in both universities are constantly contaminated with huge concentrations chemical elements from geological samples, forcing us to find different way to accomplish our research tasks. Fortunately, last Fall we received a NSF grant to upgrade some of our research facilities. With this grant, we are purchasing a new Delta PLUS XL Mass Spectrometer with continuous flow sample analysis for stable isotopes ( $^2\text{H}$  and  $^{18}\text{O}$ ). This machine in combination with Delta Plus Mass Spectrometer equipped with the gas bench device



will give us a very speedy analysis rate. In the photos below, you may see our new 80 m<sup>3</sup> freezer and ice-core laboratory at the University of Idaho.



Ice-core freezer in the Mines building, UofI



The 100 level clean bench and equipment for core processing in the Ice-Core Laboratory, UofI, Mines building



New Stable Isotope laboratory in the College of Natural Resources, University of Idaho

## **References:**

- Aizen, V. B., E.M. Aizen, J.M. Melack, K.J. Kreutz, L.D. Cecil. Association between atmospheric circulation patterns and firn-ice core records from the Inilchek glacierized area, Central Tien Shan, Asia. *Journal of Geophysical Research*, 2004a (in press).
- Aizen, V.B., E.M. Aizen, K. Fujita, S. A. Nikitin, K.J. Kreutz, L.D. Cecil, N. Takeuchi. Stable-isotope time series and precipitation origin from firn cores and snow samples, Altai glaciers, Siberia. *Journal of Glaciology*, 2004b (in press).
- Aizen, V.B., E.M. Aizen, K. Fujita, K.J. Kreutz, L.DeW. Cecil and S.A. Nikitin. Approaches to ice core climatic reconstruction in central Asia (Tien Shan and Siberian Altai). In Cecil, L.DeW., L.G. Thompson, E.J. Steig and J.R. Green, eds. *Earth paleoenvironments: records preserved in mid and low latitude glaciers*. Dordrecht, *Kluwer Academic Publishers*, p. 38-56, 2003.
- Aizen V.B., Aizen E.M. Hydrological Cycle on North and South Peripheries in Mountain-Glacial Basins of Central Asia. *Hydrological Processes*, Vol. 11, pp. 451-469, 1997.
- Aizen, V.B., E.M. Aizen, J.M. Melack, and T. Martma. Isotopic measurements of precipitation on Central Asian glaciers (southeast Tibet, northern Himalayas, central Tien Shan). *J. Geoph. Res.*, 101, 9185-9196, 1996.
- Aizen, V.B., S.A. Nikitin, Song Guoping. Model of the Dynamics of the Hailuogou Glacier (Southeastern Xizang). *J. Glaciology and Geocryology*, No 5, pp.121-132, 1994.
- Olivier, S. and 10 others. Glaciochemical investigation of an ice core from Belukha glacier, Siberian Altai. *Geophys. Res. Lett.*, 30(19), 2003 (10.1029/2003GL018290.), 2003.

- Barlow, L.K., J.W.C. White, R.G. Barry, J.C. Rogers and P. Grootes.. The North Atlantic oscillation signature in deuterium and deuterium excess signals in the Greenland Ice Sheet Project 2 ice core, 1840–1970. *Geophys. Res. Lett.*, 20(24), 2901–2904, 1993.
- Craig, H. Isotopic variations in meteoric waters. *Science*, 133(3465), 1702–1703, 1961
- Glukh, I.S. and N.K. Kononova. Vzaimosvyaz' izmencivosti mnogoletnih osadkov v Zapadnoi Sibiri I Sredney Asii. [Relationship of long-term precipitation variability in western Siberia and Middle Asia]. *Izv. Akad. Nauk SSSR, Ser. Geogr.*, 3, 23–29, 1982 [In Russian.]
- Dansgaard, W. Stable isotopes in precipitation. *Tellus*, 16, 436-468, 1964.
- Dansgaard, W. and H. Tauber. Glacier oxygen-18 content and Pleistocene ocean temperatures. *Science*, 166, 3904, 499-502, 1969.
- Dansgaard, W., S. Johnsen, H. Clausen, D. Dahl-Jensen, N. Gundestrup, C. Hammer, C. Hvidbeg, J. Steffensen, A. Sveinbjornsdottir, J. Jouzel and G. Bond. Evidence for a General Instability of the Past Climate from a 250 kyr Ice Core Record. *Nature* 364, 218-220, 1993.
- Easterling, D.R., T.C. Peterson, and T.R. Karl,: On the development and use of homogenized climate data sets. *6<sup>th</sup> International Meeting on Statistical Climatology*, June 19-23. Galway, Ireland, 435-438, 1995.
- Friedman, I., A.C. Redfield, B. Shoen and J. Harris. The variations of the deuterium content of natural waters in the hydrological cycle. *Rev. Geophys.*, 2, 177–224, 1964.
- Froehlich, K., J.J. Gibson and P. Aggarwal,: Deuterium excess in precipitation and its climatological significance. *Study of Environmental changes using isotope techniques*, C&S Papers Series 13/P, International Atomic Energy Agency, Vienna, Austria, 54 – 65, 2002.
- Johnsen, S.J.:. Stable isotope homogenization of polar firm and ice. *International Association of Hydrological Sciences Publication*, 118. Symposium in Grenoble 1975 – Isotopes and Impurities in Snow and Ice, 210-219, 1977.
- Jouzel, J., R.B. Alley, K.M. Cuffey, W. Dansgaard, P.M. Grootes, G. Hoffman, S.J. Johnsen, R.D. Koster, D. Peel, C.A. Shuman, M. Stievenard, M. Stuiver and J.W.C., White,: Validity of the temperature reconstruction from water isotopes in ice cores. *J. Geoph. Res.*, 102, 26471-26488, 1997.
- Kendall, C. and J.J. McDonnell, eds.. *Isotope tracers in catchment hydrology*. Amsterdam, etc., Elsevier, 1998.
- Koerner, R.M. and D.A. Fisher. A record of Holocene summer climate from a Canadian high-Arctic ice core. *Nature*, 343, 630-631, 1990.
- Kotlyakov, V.M. (Ed.). *World Atlas of Snow and Ice Resources*, Minsk, 400 p., 1998.
- Kotlyakov, V.M. and F.G. Gordienko.. *Isotopnaya I geochemicheskaya glyaciologiya. [Isotopic and geochemical glaciology]*. Leningrad, Gidrometeoizdat, 1982. [In Russian.]
- Kreutz, K., V. B. Aizen, D. Cecil and C. Wake. Dust deposition and isotopic composition of precipitation recorded in a shallow ice core, Inilchek glacier, central Tien Shan. *J. Glaciology*, 47, 159, 549-554, 2001.
- Lin, P.N., L.G. Thompson, M.E. Davis and E. Mosley-Thompson,: 1000 years of climatic changes in China: ice core  $\delta^{18}\text{O}$  evidence. *Ann. Glaciology*, 21, 189-195, 1995.
- Naftz, D.L., D.D. Susong, P.F. Schuster, L.D. Cecil, M.D. Dettinger, R.L. Michel, and C. Kendall,: Ice core evidence of rapid air temperature increases since 1960 in alpine areas of the Wind River Range, Wyoming, United States. *J. Geoph. Res.*, 107, 2002.
- Naftz, D., J.A.Rice, and J.R.Ranville,. Glacial Ice Composition: A potensial long-term record of the chemistry of atmospheric deposition, Wind River Range, Wyoming. *Water Resources Research*, 27, No.6, p. 1231-1238, 1991.
- Richman, M.B.. Rotation of principal components. *J. Climatol.*, 6(\*), 293–335, 1986.
- Shuman, C.A., R.B. Alley, S. Anandkrishnan, J.W.C. White, P.M. Grootes, and C.R. Stearns. Temperature and accumulation at the Greenland Summit: comparison and high resolution isotope profile and satellite passive microwave brightness temperature trends. *J. Geoph. Res.*, 100 (D5), 9165-9177, 1995.

- Thompson, L.G., E. Mosley-Thompson, M.E. Davis, P.N. Lin, J. Dai and J. Bolzan. A 1000 year ice core climate record from the Guliya Ice Cap, China: its relationship to global climate variability. *Ann. Glaciology*, 21, 175-181, 1995.
- Yao, T., V. Masson, J. Jouzel, M. Stievenard, S. Weizen and J. Keqin,: Relationships between  $\delta^{18}\text{O}$  in precipitation and surface air temperature in the Urumqi River Basin, east Tien Shan Mountains, China. *Geoph. Res. Lett.*, 26 (23), 3473-3476, 1999.
- Yao, T., L.G. Thompson, K. Jiao, E. Mosley-Thompson and Z. Yang,: Recent warming as recorded in the Qinghai-Tibetan Cryosphere. *Ann. Glaciology*, 21, 196-200, 1995.
- Yao, T. and L.G. Thompson. Trends and features of climatic changes in the past 5000 years recorded by the Dunde ice core. *Ann. Glaciology*, 16, 116-119, 1992.

## **Publications:**

- (1) Aizen, V. B., E.M. Aizen, J.M. Melack, K.J. Kreutz, L.D. Cecil. Association between atmospheric circulation patterns and firn-ice core records from the Inilchek glacierized area, Central Tien Shan, Asia. *Journal of Geophysical Research*, 2004 (in press).
- (2) Aizen, V.B., E.M. Aizen, K. Fujita, S. A. Nikitin, K.J. Kreutz, L.D. Cecil, N. Takeuchi. Stable-isotope time series and precipitation origin from firn cores and snow samples, Altai glaciers, Siberia. *Journal of Glaciology*, 2004 (in press).
- (3) Aizen, V.B., E.M. Aizen, K. Fujita, K.J. Kreutz, L.DeW. Cecil and S.A. Nikitin. Approaches to ice core climatic reconstruction in central Asia (Tien Shan and Siberian Altai). In Cecil, L.DeW., L.G. Thompson, E.J. Steig and J.R. Green, eds. *Earth paleoenvironments: records preserved in mid and low latitude glaciers*. Dordrecht, *Kluwer Academic Publishers*, p. 38-56, 2003.
- (4) Kreutz, K. J. C. P. Wake, V. B. Aizen, L. DeWayne Cecil, Hans-Arno Synal. 2003. Seasonal deuterium excess in a Tien Shan ice core: Influence of moisture transport and recycling in Central Asia. *Geophysical Research Letters*, Vol. 30, №. 18, 1922, doi: 10.1029/2003GL017896, 2003.
- (5) Kreutz, K.J., C.P. Wake, V.B. Aizen, L.D. Cecil, J.R. Green, H-A. Synal. Event to Decadal-Scale Glaciochemical Variability on the Inilchek Glacier, Central Tien Shan. ). In Cecil, L.DeW., L.G. Thompson, E.J. Steig and J.R. Green, eds. *Earth paleoenvironments: records preserved in mid and low latitude glaciers*. Dordrecht, Kluwer Academic Publishers, p. 18-24, 2003.
- (6) Aizen, V.B., E.M. Aizen. Spatial and temporal variability of precipitation and snow pack in Tien Shan, Japanese Islands and Sierra Nevada Mountains. In Proceedings: *International Symposium on Disaster Mitigation and Basin-wide Water management*, p. 74-87, 2003.
- (7) Takeuchi, N., A. Takahashi, J. Uetake, T. Yamazaki, V. Aizen, D. Joswiak, A. Surazakov, S. Nikitin. A report on ice core drilling operation at the western plateau of Mt. Belukha in the Altai Mountain Range in 2003. In *Journal of Polar Meteorology and Glaciology*, V. 6, p. 23-47, 2003.
- (8) Fujita, K., N. Takeuchi, V. Aizen, S. Nikitin. Glaciological observations on the plateau of Belukha Glacier in the Altai Mountains, Russia from 2001 to 2003. *Glaciological Bulletin*, V. 4, p. 31-50, 2003.
- (9) Aizen, V.B., E.M. Aizen. Global Climate and Environmental Changes in Alpine Asia (Tibet, Himalaya, Tien Shan, Pamir and Altai). NSF GCOS Workshop, Boulder, Colorado, March 2003 (*presentation*)
- (10) Aizen, V.B., E.M. Aizen. Major peculiarities of climate and ecosystems in the Northern Eurasia. NEESPI NASA. Proceeding: *Conference, Suzdal*, Russia, April 2003.
- (11) Aizen, V.B., V.A. Kuzmichenok, E.M. Aizen, A.B. Surazakov. Tien Shan glacier's recession since the end of 19th century and during last 60 years by land instrumental, aerial photo and satellite remote sensing data. Proceedings: *San Francisco Fall AGU Meeting*, December 2003.



- (12) Aizen, E.M., V.B. Aizen, K.Fujita, S.A.Nikitin, K.J.Kreutz. Snow and ice-core stable isotopes time series and precipitation in Altai Mountains (Siberia). Proceedings: *San Francisco Fall AGU Meeting*, December 2003.
- (13) Aizen, E.M., V. B. Aizen, D. Joswiak. Spatial variation of ionic and isotopic records from snow pits and shallow ice cores in Central and Southeast Asia. Proceedings: *San Francisco Fall AGU Meeting*, December 2003.
- (14) Surazakov, A.B., Yu. Narojniy, S.A. Nikitin, **V. B. Aizen, E. M. Aizen**. Glacier Evolution in the Altai Mountains, South-West Siberia, for the Last Half Century. Proceedings: *San Francisco Fall AGU Meeting*, December 2003.
- (15) Aizen, V.B., E.M. Aizen. Spatial and temporal variability of precipitation and snow pack in Tien Shan, Japanese Islands and Sierra Nevada Mountains. Proceedings: *International Symposium on Disaster Mitigation and Basin-wide Water management*, Niigata, Japan, December 2003.
- (16) Aizen, V. B, E. M.Aizen, J. M. Melack, T. Nakamura, S. Kobayashi. Estimation of the energy used to melt snow in Tien Shan mountains and Japanese Islands. Pergamon Elsevier Science, *Global and Planetary Change*, V. 32, Issue 4, pp. 349-359, 2002.
- (17) Aizen, V. B., E. M. Aizen, V. N. Nikitin. Glacier regime on the northern slope of the Himalaya (Xixibangma glaciers). Pergamon Elsevier Science, *Quaternary International*, V. 97-98, pp. 27-39, 2002.
- (18) Aizen, V.B., E.M. Aizen, K.J. Kreutz, K.Fujita, L.D. Cecil, S.A.Nikitin. Climatic and environmental records from Altai glaciers, Siberia, recovered from ice-cores and snow samples. Proceedings, *San Francisco AGU Fall Meeting*, 2002.
- (19) Kreutz, K. J., C. P. Wake, V. B. Aizen, L. D. Cecil, J. R. Green, H.A. Synal, D. S. Introne. Seasonal deuterium excess in a Tien Shan ice core: influence of moisture transport and recycling in Central Asia. Proceedings: *San Francisco AGU Fall Meeting*, PP42B-0529, 2002.
- (20) Aizen, V.B. Paleoclimatic and Glaciological Reconstructions in Central Asia through collection and analysis of ice-cores and instrumental data in the Tien Shan and Altai Mountains. *Annual Report to DOE/NSF* 2002, 31 p.
- (21) Aizen, V.B. Climatic and Environmental Changes in Tien Shan and Tibet Alpine Areas. Proceedings: *International CliC Conference* in Lanzhou, October 2002.

## APPENDIX

### AGREEMENT ON COLLABORATIVE RESEARCH

University of Idaho (UofI) College of Science P.O. Box 443025 Moscow Idaho 83844-3025, U.S.A. E-mail: aizen@uidaho.edu	Lanzhou, Cold & Arid Regions Environment and Engineering Research Institute (LCAREERI), Chinese Academy of Sciences People Republic of China 260 West Donggang Road, Lanzhou 730000, P.R. of CHINA E-mail: liusy@ns.lzb.ac.cn
--	--

#### LONG-TERM DYNAMIC OF SOUTHERN MONSOONS AND HYDRO-CLIMATOLOGICAL REGIME OF THE SOUTHEASTERN TIBETAN ALPINE BASINS

This document is an advanced Agreement in developing the Collaborative Research Project between the University of Idaho (UofI), and the Lanzhou Cold and Arid Regions Environmental and Engineering Research Institute, Chinese Academy of Sciences, (LCAREERI – the People Republic of China, Principal Institution). The General 3-4 years long-term Project-Agreement on joint investigation of the Climate and Water Resources Changes in South-East Asia between UofI and LCAREERI can be signed after the first research expedition in Southeastern Tibet (Bomi glaciation) in spring or autumn 2002 and upon receiving financial support from the sponsored Agencies of the USA and China for 3-4 years field and analytical research in this area.

#### Objectives of the Agreement

- 1.1 Organization and achievement of three joint glacio-climatic environmental monitoring expeditions to Southeastern Tibet glaciation area (Bomi)
- 1.2 Investigation of the climatic changes and monsoon dynamics in the South and South-East Asia based on hydro-meteorological and synoptic data from the Tibetan Mountains over the last 50 years and three-four year all year around hydro-meteorological and glaciological monitoring in Bomi glaciation area with automatic weather stations
- 1.3 Study the glaciers and snow cover existence climatic regime glaciers mass balance and runoff dynamics during the last 50 to 100 years in tropical monsoon region
- 1.4 Study precipitation, snow, ice and water geochemistry, isotopic and trace metal dynamics in the region during last 50 to 100 years
- 1.5 Development Land-Surface-Atmosphere Model Interaction in South Asia Tibet Monsoon Area emphasizes on recent climatic and environmental changes in the region

## **The UofI 2004 expedition scope**

The purpose of this expedition is to recover several surface to bottom ice-cores in Bomi glaciated area, south-eastern Tibet. Additionally, during one-month expedition we will retrieve meteorological records collected through the year 2003/2004 in two automatic weather stations established in 2002, achieve second year GPS survey on Zuoqiupu and Yalong Glaciers, collect long-term data from nearby hydro-meteorological stations, snow and water samples and tree-cores. The collected data and samples will be processed in field, at the UofI, Nevada Desert Research Institute (DRI) and LCAREERI laboratories through the physical, isotope, geo-chemical and radionuclide analysis. GPS satellite data, tree-core analysis, glacier mass balance calculation, data statistical control and analysis and mathematical simulation will occur in UofI. With growing numbers of precisely collected environmental data in Asia we will begin to examine the climatic appearance of regional to global synoptic-scale atmospheric phenomena like the ENSO and the tropical and sub-tropical monsoon and changes in glaciers mass balance and water resources.

Our particular interest focuses on glaciers of southeastern Tibet (the Bomi region) for several reasons.

- 1) The existence of widespread glaciation in a tropical-monsoon belt.
- 2) The location of the research area on the southeastern margin of the Tibetan Plateau provides an opportunity to collect information relating to the tropical monsoons and Tibetan High.
- 3) To study unique Tibetan glaciers that hold the greatest concentration of snow and ice in the low latitudes and constitute a source of fresh water shared by ten heavily populated neighboring countries, the glaciers that can cause the flood disasters in certain climatic and environmental changes.
- 4) Decadal records of atmospheric deposition in inhabited areas of the Tibetan Plateau are desirable to evaluate the possible relation between anthropogenic emissions (such as Green House Gases) and chemical composition of snow/ice depositions

The proposed one month field research will include:

1. Recover several surface to bottom ice-cores on Zuoqiupu and Yalong Glacier (140-200 m long each) and deliver cores frozen to the University of Idaho freezer;
2. Retrieve data from two automatic weather stations (Campbell Scientific) collected during year 2003/2004 and load stations for one more year observations;
3. Complete second year radio-echo sounding and GPS survey of the glaciers to determine present glaciers ice thickness and the glacier surface parameters and compare data with 2002 and 2003 measurements;
4. Collect new snow, precipitation and stream water samples in the studied area;
5. Collect tree-cores in the Bomi forest zone for cross correlation analysis with ice-cores;
6. Collect long-term meteorological and synoptic data from the region of investigation extending back 40-50 years for further data analysis comparison and calibration.

The results of the preliminary studies will be submitted in special reports and publications in appropriate journals. All results of scientific investigations will be the ownership of UofI, DRI and LCAREERI collaborators.

## Obligations

**A. The UofI and DRI** will provide major scientific instrumentation for field research and the cost of camping equipment and supplies during the expedition according to the Year 2004 budget and scientific schedule. Specifics of the instrumentation and camping equipment are following:

- a) Solar-powered mechanical auger (equipment will be return to U.S.A. every year)
- b) U.S.A. team will be shared shallow ice-core with LCAREERI 50 by 50 after we will recover them
- c) Portable radio echo-sounding instruments for the measurements of ice/firn thickness (return to U.S.A. every year)
- d) Snow and firn sampling and field processing kit and clean plastic bottles to collect and transport samples (return to U.S.A. every year)
- e) GPS equipment to measure glacier ice velocity and the glacier terminus dynamics (return to U.S.A. every year)
- f) Dendro-drills and dendro-containers kit to collect dendro-cores (return to U.S.A. every year)
- g) Insulated firn/ice shipping containers (return to U.S.A. every year).
- h) UofI will invite will invite one post-graduate student for the period of our research in Tibet (4 years) to complete his PhD thesis upon we will get an additional fund and stipend for the student.

**B. The LCAREERI** will provide the following during the agreement and expeditions in year 2002 and through year 2005

- a) Provide and direct qualified, communicative and responsible collaborators that are the specialists in climatology, glaciology, and isotop-geochemistry and will be able to work at altitudes over 4000-6000 m
- b) Assist in collection existing synoptical, meteorological, hydrological and other necessary data and information for the Project fulfilling an initial statistic data control and analysis
- c) Implement and solve all formal acquiescence (at the Government level of the People Republic of China) associated with obtaining permission for:
  - the fulfillment of the scientific investigations at the Tibet territory
  - permission for installation automatic weather stations, GPS and Satellite Phone use in field
  - permission to transport snow, ice and water samples and dendro-cores from Tibet, through China to U.S.A. in 2002-2005.

## The UofI field 2004 budget portion

Budget Category					Total
<b>Field expenditures</b>					
Permission for intrance to Tibet for work	4	persons			\$1 500 /pers \$ 6 000
Rental truck and jeeps	3	vehicles	40	days	\$90 /day \$10 800
Fuel			9000	liters	\$0.9 /liter \$ 8 100
Food in field	12	persons	30	days	\$10 /day \$3 600
Yaks and skimmers	6	yaks	20	days	\$40 /day \$4 800
Cook	1	person	40	days	\$40 /day \$1 600
Porters	6	persons	40	days	\$40 /day \$9 600
Maintainance of the metstations	1	station	12	month	\$300 /month \$3 600
<b>Field expenditures Subtotal:</b>					<b>\$48 100</b>
<b>Supplies</b>					
Base camp equipment					\$4 000
Mountain gear for porters					\$3 500
Mountain gear for the skimmers	6	persons			\$300 /pers \$1 800
<b>Supplies Subtotal:</b>					<b>9 300</b>
<b>Other Direct costs</b>					
Service contract with CAREERI, (China)					
20% from the field cost:			\$57 400	@ 20%	\$11 480
<b>Total Cost:</b>					<b>\$68 880</b>

This budget doesn't include the airfare between USA and Tibet, accommodation and per-diem for the USA team in Chengdu and Bangda, cost of excess luggage in flights between USA and Chengdu and Chengdu-Bangda, cost of the USA team mountaineering and scientific equipment, cost of the laboratory data analysis, and other expenditures.

## The LCAREERI Field 2002 Budget portion

Budget Category					Total
<b>Field expenditures</b>					
Permission for intrance to Tibet for work	9	persons			\$200 /pers \$1,800
Rental truck and jeeps	1	vehicles	40	days	\$90 /day \$3,600
Fuel			3000	liters	\$0.7 /liter \$2,100
Food, and emergency expenditures	12	persons	10	days	\$10 /day \$1,200
Yaks and skimmers	2	yaks	15	days	\$40 /day \$1,200
Porters	2	persons	40	days	\$40 /day \$3,200
Maintainance of the metstations	1	station	12	month	\$300 /month \$3,600
<b>Field expenditures Subtotal:</b>					<b>\$16,700</b>
<b>Supplies</b>					
Base camp equipment					\$1,500
Mountain gear for porters					\$1,400
Mountain gear for the skimmers	2	persons			\$200 /pers \$400
<b>Supplies Subtotal:</b>					<b>3,300</b>
<b>Total Cost:</b>					<b>\$20,000</b>

This preliminary Agreement begins from the time of signing by both collaborators. This agreement is in effect upon new general agreement will be signed but no late than December 31, 2005. The agreement will be extended in details **EACH** year if both parties agree to continue this research.

**The Preliminary-Agreement and field budget has been signed:**

From the University of Idaho  
Leader of the Project  
Prof., Dr. Vladimir B. Aizen

From the Lanzhou Cold and Arid Regions  
Environmental Engineering Research  
Institute, Chinese Academy of Sciences,  
Lanzhou, 730000, P.R. China  
Director, Prof. Dr. Cheng Godung



*Lanzhou (China), Moscow Idaho (U.S.A.)  
October, 2003*