

1 **Remote sensing for prediction of 1-year post-fire ecosystem condition**

2 *Leigh B. Lentile*^{A,D,*}, *Alistair M. S. Smith*^{B,*}, *Andrew T. Hudak*^C, *Penelope Morgan*^B, *Michael J. Bobbitt*^B,
3 *Sarah A. Lewis*^C and *Peter R. Robichaud*^C

4 ^ADepartment of Forestry and Geology, University of the South, Sewanee, TN 37383, USA.

5 ^BDepartment of Forest Resources, University of Idaho, Moscow, ID 83844-1133, USA.

6 ^CRocky Mountain Research Station, US Department of Agriculture Forest Service, Moscow, ID 83843,
7 USA.

8 ^DCorresponding author. Email: lbentil@sewanee.edu

9 **WF07091 TOC** We compare and evaluate the applicability of immediate post-fire estimates of percentage char and
10 vegetation fractions, in addition to NBR and dNBR derived from Landsat ETM+ imagery, to remotely predict 1-year
11 post-fire ecological effects. The char and green fractions are versatile indicators of canopy and subcanopy effects
12 and longer-term effects related to fire behavior.

13 Appropriate use of satellite data in predicting >1 year post-fire effects requires remote measurement of surface
14 properties that can be mechanistically related to ground measures of post-fire condition. The present study of burned
15 ponderosa pine (*Pinus ponderosa*) forests in the Black Hills of South Dakota evaluates whether immediate fractional
16 cover estimates of char, green vegetation, and brown (non-photosynthetic) vegetation within a pixel are improved
17 predictors of 1-year post-fire field measures, when compared with single-date and differenced Normalized Burn
18 Ratio (NBR and dNBR) indices. The modeled estimate of immediate char fraction either equaled or outperformed
19 all other immediate metrics in predicting 1-year post-fire effects. Brown cover fraction was a poor predictor of all
20 effects ($r^2 < 0.30$), and each remote measure produced only poor predictions of crown scorch ($r^2 < 0.20$). Application
21 of dNBR (1 year post) provided a considerable increase in regression performance for predicting tree survival.
22 Immediate post-fire NBR or dNBR produced only marginal differences in predictions of all the 1-year post-fire
23 effects, perhaps limiting the need for prefire imagery. Although further research is clearly warranted to evaluate fire
24 effects data available 2–20 years after fire, char and green vegetation fractions may be viable alternatives to dNBR
25 and similar indices to predict longer-term post-fire ecological effects.

26 WF07091

27 L.B. Lentile *et al.*

28 Fractional cover and fire effects

29 **Additional keywords:** burn severity, char, Landsat ETM+, ponderosa pine, subpixel, unmixing.

30 *Leigh B. Lentile and Alistair M. S. Smith contributed equally to the present paper.

1 **Introduction**

2 The large size, and in many cases, remote nature of many wildfires has made analysis of Earth
3 observation imagery an important and widely applied method for immediate and long-term assessment of
4 fire effects on ecosystems (Morgan *et al.* 2001; Lentile *et al.* 2006). Appropriate use of remote sensing
5 tools and techniques in predicting these fire effects, such as vegetation recovery and successional
6 processes, requires that we investigate the mechanistic, biophysical relationships between remotely
7 sensed metrics of post-fire surface condition, such as changes in reflectance, surface temperature, heights
8 (e.g. using laser altimetry data), or fractional cover; with field measures of ecosystem condition (Lentile
9 *et al.* 2006; Key 2006). Definitions and assessments of post-fire ecosystem condition often use the word
10 ‘severity’, which for the present paper will be described as ‘burn severity’ and includes changes in both
11 soil and vegetation conditions as a result of fire (Lentile *et al.* 2006).

12 Although early remote sensing research to infer the severity of fires focussed on metrics that could be
13 both measured on the ground and inferred by the sensors, such as crown consumption or subsequent tree
14 mortality (Patterson and Yool 1998; Miller and Yool 2002), recent research has predominately focussed
15 on using the differenced Normalized Burn Ratio (dNBR: Key and Benson 2006) spectral index or variants
16 thereof (Holden *et al.* 2005; Miller and Thode 2007). This spectral index effectively measures the relative
17 degree of vegetation and soil/char cover change between pre- and post-fire conditions (Smith *et al.* 2005;
18 Lentile *et al.* 2006). Within North American wildfires, these values have been evaluated predominantly
19 against an ocular field assessment method termed the Composite Burn Index (CBI) (van Wagendonk *et*
20 *al.* 2004; Brewer *et al.* 2005; Cocke *et al.* 2005) with only limited studies modeling or evaluating
21 regressions with specific biological/ecological measures of post-fire effects (De Santis and Chuvieco
22 2007; Hudak *et al.* 2007b; Robichaud *et al.* 2007; Smith *et al.* 2007b).

23 Although fundamentally an ocular measurement, CBI is an integrative measure of post-fire effects
24 across under- and overstorey strata (Key and Benson 2006). Although it is not ideal to compare singular-
25 date field measurements with change-detection indices (Dozier and Strahler 1983), the CBI methodology
26 is often applied because the assessment of fire effects in wildland fires typically occurs in an opportunistic
27 or a rapid response (Lentile *et al.* 2007a) fashion, which limits the likelihood of available prefire data
28 (Lentile *et al.* 2006). Although widely applied by fire management in the production of burned area
29 reflectance classification (BARC) maps and within the national Monitoring Trends in Burn Severity
30 (MTBS) program (e.g. Cocke *et al.* 2005; Epting *et al.* 2005; Miller and Thode 2007), limitations in both
31 the dNBR and CBI methodologies have been highlighted (Roy *et al.* 2006; Smith *et al.* 2007b):

- 1 (i) The CBI measure is calculated in a highly subjective and qualitative manner, with evaluations often
2 conducted without explicit knowledge of prefire ecosystem condition (van Wagtenonk *et al.* 2004)
3 but rather with unburned adjacent areas used as prefire surrogates (Lentile *et al.* 2006);
- 4 (ii) dNBR often exhibits non-linear asymptotic relationships with CBI (van Wagtenonk *et al.* 2004;
5 Cocks *et al.* 2005; Wimberly and Reilly 2007), which leads to scaling challenges. This effect further
6 varies with ecosystem type (Epting *et al.* 2005) and with the spatial resolution of the satellite sensor
7 (van Wagtenonk *et al.* 2004; Holden *et al.* article in review);
- 8 (iii) Contemporary remote sensing studies have shown that the spectral bands used to calculate NBR are
9 not optimal to evaluate the degree of burning (Smith *et al.* 2005; Roy *et al.* 2006);
- 10 (iv) dNBR and CBI have been shown to be suboptimal in woodland, shrub, and grassland environments
11 (Epting *et al.* 2005; Roy *et al.* 2006; De Santis and Chuvieco 2007), resulting in studies considering
12 variants of both CBI (Epting *et al.* 2005; De Santis and Chuvieco 2007) and dNBR, such as the
13 relative dNBR (RdNBR), to assess post-fire effects (Miller and Thode 2007). Although other
14 investigations show no improvement of RdNBR over dNBR when applied in environments beyond
15 those similar to that RdNBR was developed within (Hudak *et al.* 2007b), it has the potential to
16 provide consistently interpretable results across multiple environments (Miller and Thode 2007;
17 Safford *et al.* 2007);
- 18 (v) The remote sensing literature has demonstrated the spectral changes that dNBR highlights are due to
19 differences in the amount of vegetation, soil, and char detected (Smith *et al.* 2005; Roy *et al.* 2006)
20 and although it has been highlighted that dNBR predominately detects changes in vegetation
21 consumed (Hudak *et al.* 2007b) or killed (Miller and Thode 2007), disagreement exists within the fire
22 ecology community as to whether dNBR maps should be used to only infer fire effects on soil and not
23 on vegetation (Odion and Hanson 2006, 2007; Safford *et al.* 2007);
- 24 (vi) Contemporary studies have questioned whether studies should infer post-fire effects from the
25 immediate post-fire NBR, the immediate post-fire dNBR ($NBR_{pre} - NBR_{immediate\ post-fire}$), or the 1-year
26 post-fire dNBR ($NBR_{pre} - NBR_{1-year\ post-fire}$), which can lead to confusion when selecting methods for
27 assessment (Epting *et al.* 2005; Hudak *et al.* 2007b). Specifically, some studies highlight dNBR
28 (Epting *et al.* 2005) or conversely NBR (Bobbe *et al.* 2001; Hudak *et al.* 2007b) to be improved
29 predictors of post-fire effects. Motivation for using dNBR or similar multitemporal imagery includes
30 the potential to minimize classification errors due to sun-sensor geometry, atmospheric effects,
31 phenology, or areas that are spectrally flat such as water or older burns (Bobbe *et al.* 2001; De Santis
32 and Chuvieco 2007).

1 (vii) Roy *et al.* (2006) highlighted that the original application of the dNBR was for burned area mapping
2 (Lopez-Garcia and Caselles 1991), which relies on fundamentally opposite assumptions to methods
3 used to assess a range of biophysical variation within an area (Verstraete and Pinty 1996; Roy *et al.*
4 2006), such as a range of ‘severity’ after a wildfire.

5 Specifically in terms of (vii), numerous authors have remarked that the ultimate goal of any land-cover
6 classification approach is ideally to produce class histograms that are narrowly peaked (low internal
7 variability) but that have well-separated means, such that the different class histograms are less likely to
8 overlap and therefore would exhibit higher spectral separation (Verstraete and Pinty 1996; Pereira 1999;
9 Roy *et al.* 2006). In contrast, when evaluating within-area effects, such as burn severity, a large dynamic
10 range of within-class values is desired to provide detailed characterization of those effects. In essence,
11 although well-separated class means are still desired, the user now needs the individual class histograms
12 to be very wide, or at least exhibit bi- or tri-modal properties, to enable splitting of any particular class
13 into distinct regimes, such as ‘low, moderate and high’. When such bi- or tri-modal properties are not
14 immediately apparent, it is common to use statistical breaks, such as those derived from training data
15 (Pereira 1999; Hudak *et al.* 2007a).

16 As these are mutually exclusive objectives (Verstraete and Pinty 1996; Pereira 1999), dNBR and any
17 other similar spectral index cannot be optimal for characterizing both burned area and post-fire effects
18 related to severity (Roy *et al.* 2006). Owing to mixed results in the application of dNBR to severity
19 assessments in a range of fire types outside the area around Glacier National Park, Montana, for which it
20 was originally developed (Key 2006), it remains apparent that more research is needed to determine
21 whether dNBR is better suited to the assessment of burn area or burn severity.

22 These factors potentially limit the wide-scale applicability of dNBR and similar spectral indices to infer
23 post-fire ecosystem condition and highlight the continued need for further research to evaluate alternative
24 and perhaps more appropriate remote sensing methods (Roy *et al.* 2006). Recent research has highlighted
25 spectral mixture analysis (SMA) as one such alternative approach with potential to meet this need (Lentile
26 *et al.* 2006; Smith *et al.* 2007a, 2007b; Hudak *et al.* 2007b). To date, SMA (common synonyms: linear
27 spectral unmixing, mixture modeling) has predominately been used to map the extent of the burned area
28 (Wessman *et al.* 1997; Cochrane and Souza 1998; Vafeidis and Drake 2005; Smith *et al.* 2007a). It also
29 enables estimation of fractional cover components with each multispectral image pixel, including
30 unburned vegetation (green or senesced), soils, and charred or fully combusted vegetation (Smith and
31 Hudak 2005; Smith *et al.* 2005; Robichaud *et al.* 2007). Although other subpixel methods such as mixture
32 tuned matched filtering (Robichaud *et al.* 2007) and fuzzy classification methods exist (Foody 2000),
33 SMA has widely been applied to the analysis of ecological data (Wessman *et al.* 1997) and its theory and

1 limitations are well documented in the literature (Drake *et al.* 1999; Theseira *et al.* 2003). SMA relies on
2 the assumptions of linear spectral mixing models (Drake *et al.* 1999) and thus the results are inherently
3 scalable across data of different spatial resolutions (Settle and Drake 1993). SMA also can be applied to
4 any type of imagery with multiple reflectance channels in the visible and near-infrared wavelength
5 regions, without reliance on the availability of specific channels (e.g. bands 4 and 7 to calculate dNBR
6 from Landsat TM or ETM+). Furthermore, it allows production of measures that are directly analogous to
7 traditional 'field severity' assessments of % green, % brown, and % black (Lentile *et al.* 2006).

8 Smith *et al.* (2007b) observed in a recent preliminary study that in comparison with the immediate
9 post-fire metrics of dNBR and fractional green cover, the estimate of fractional char cover applied to a
10 mixture of aspen and ponderosa pine plots produced marginally improved predictions of two 1-year post-
11 fire effect measures (% live trees and organic litter weight). However, in contrast, Hudak *et al.* (2007b)
12 observed that green fractional cover was an equal or improved correlate to multiple post-fire effects when
13 compared with immediate post-fire NBR and dNBR. This conflicting result of Hudak *et al.* (2007) could
14 in part be due to the ponderosa pine and aspen stands exhibiting different spectral and fire effect
15 characteristics, highlighting that these data be reevaluated for single-species stands. Moreover, the
16 apparent ability of a remote sensing method to reasonably predict a subcanopy post-fire effect offers
17 considerable promise to remotely assess ecological indicators of the fire intensity and severity (Smith *et*
18 *al.* 2007b). Therefore, further research is warranted to assess whether similar fractional cover estimates of
19 the char, green vegetation, and also senesced vegetation (or a mixture thereof), can also predict a wider
20 range of both canopy and subcanopy post-fire effects, beyond the two 1-year post-fire measures
21 previously evaluated. Therefore, following on from Smith *et al.* (2007b) and Hudak *et al.* (2007b), the
22 objectives of the present study are:

- 23 (1) Evaluate whether SMA-derived estimates of fractional char, green, and brown vegetation covers
24 immediately post-fire are improved predictors, over immediate NBR and dNBR, for a wide variety of
25 both canopy (four) and subcanopy (nine) ecological indicators measured in 66 ponderosa pine plots 1
26 year after fire;
- 27 (2) Evaluate whether these immediate fractional measures are improved correlates of 1-year post-fire
28 conditions when compared with dNBR calculated from pre- and 1-year post-fire imagery;
- 29 (3) Evaluate the degree of redundancy (if any) due to the different fractional cover measures when
30 predicting the 1-year post-fire effects. This will enable us to evaluate whether a combination approach
31 based on several different cover metrics has potential; and
- 32 (4) Evaluate whether any improvement in prediction performance is achieved by using the immediate
33 post-fire dNBR over NBR. Although applications of dNBR are widely presented to managers via

1 BARC maps, this analysis will enable us to determine whether the prefire imagery provides additional
2 information in predicting longer-term post-fire effects.

3 **Methods**

4 *Study area*

5 The current study focussed on the ~33 800-ha Jasper Fire, which occurred in 2000 in the Black Hills of
6 western South Dakota, USA. Within the fire, latitudes range from 43°41'35" to 43°55'48"N and
7 longitudes range from 103°46'1" to 104°0'47"W. Elevations range from ~1500 to 2100 m. The Black
8 Hills are an isolated mountain range on the Northern Great Plains physiographic province in western
9 South Dakota and north-eastern Wyoming (Fig. 1). As the easternmost extension of the Rocky
10 Mountains, the Black Hills were formed by regional uplift between ~35 and 65 million years ago. This
11 uplift produced an elliptical dome with an older crystalline core surrounded by younger, steeply dipping
12 sedimentary deposits (Shepperd and Battaglia 2002). The Limestone Plateau surrounds the core and the
13 area burned by the Jasper Fire is located on the south-western extent of this fertile plateau. The area
14 within the Jasper Fire perimeter is characterized by relatively continuous ponderosa pine (*Pinus*
15 *ponderosa*) stands, although occasional quaking aspen (*Populus tremuloides* Michx.) clones and
16 grasslands also exist. A complete description of the study area and fire regime is provided in Lentile
17 (2004) and Lentile *et al.* (2005).

18 *Remote sensing data and methods*

19 Three Landsat ETM+ images of the study area were acquired (18 August 1999; 14 September 2000; 24
20 September 2001). Each image was corrected to top-of-atmosphere reflectance using the standard
21 calibration equations. The Normalized Burn Ratio (NBR), defined as the normalized difference of
22 Landsat bands 4 and 7 (band4 – band7)/(band4 + band7), was then determined for each image. Two forms
23 of dNBR were applied, namely the ‘immediate dNBR’, which used the prefire and immediate post-fire
24 image, and the ‘1-year post-fire dNBR’, which, as the name suggests, used the prefire and 1-year post-fire
25 image in the dNBR calculation. Both the immediate dNBR and 1-year post-fire dNBR were calculated by
26 subtracting the post-fire image from the prefire image. Rather than classifying dNBR values using
27 arbitrary thresholds, we used the continuous dNBR values in subsequent regression analyses. Following
28 Cocke *et al.* (2005), each dNBR image was then scaled by multiplying each value by 1000. For this
29 analysis, the immediate post-fire Landsat ETM+ imagery was additionally converted into ground-
30 reflectance using the standard method of ‘dark-body subtraction’ using the minimum-band pixel values as
31 selected by the ITT Visual Information Systems for *ENVI* software package (Boulder, CO, USA).

1 The estimation of the fractional cover of char (Fig. 1), brown vegetation, and green vegetation within
2 each Landsat pixel was determined using spectral mixture analysis (Settle and Drake 1993). Although
3 other linear and non-linear spectral unmixing methods have been applied to the assessment of post-fire
4 affected surfaces (Smith *et al.* 2005; Robichaud *et al.* 2007), the accuracy and ease of implementation of
5 linear compared with the complexity of non-linear models has led to the widespread adoption of linear
6 SMA in Earth observation studies (Drake *et al.* 1999; Chen *et al.* 2004). Importantly, the principal
7 assumption of linear mixture models, namely that a mixture of 50% of A + 50% of B will have a spectral
8 reflectance of $((A + B)/2)$ over all analyzed wavelengths (e.g. 0.3–2.5 μm), has been shown to be broadly
9 valid when considering mixtures of unburned and burned surface components (Cochrane and Souza 1998;
10 Smith *et al.* 2005; Vafeidis and Drake 2005). The classical linear spectral unmixing model is defined by
11 Theseira *et al.* (2002) as:

$$R_n = \sum_{c=1}^n (a_{nc} x_c) + e_n \quad (1)$$

13 where, R_n is the reflectance of the pixel in the n th spectral band, x_c is the proportion of the c th component
14 in the pixel, a_{nc} is the spectral endmember of the c th component for the n th spectral band, and e_n denotes
15 the pixel noise term.

16 Generic spectra of senesced vegetation, green vegetation, and char (Fig. 2) were used as the authors of
17 several past studies have remarked that these spectral reflectance curves are broadly similar across a wide
18 range of environments (Elvidge 1990; Landmann 2003; Smith *et al.* 2005, 2007a, 2007b; Hudak *et al.*
19 2007b). As detailed by Smith *et al.* (2005), the spectral reflectance functions for representative green
20 vegetation, senesced grasses, and char (among other surfaces) were collected with a GER-3700
21 spectroradiometer (Spectravista Corporation, Poughkeepsie, NY). The GER-3700 was set up at a height
22 of 0.75 m directly above (i.e. at nadir) the vegetation and char samples. Measurements were acquired in
23 full sunlight within a 3° field of view; for each spectrum, eight spectral measurements were acquired over
24 a 5-s interval and the mean calculated. Furthermore, for each sample of vegetation and char (five of each),
25 three such sets of measurements were acquired from different vantage points to capture variability due to
26 shading and increase the signal-to-noise ratio of the resultant spectral reflectance curves. The spectral
27 radiance measurements were converted into reflectance by normalizing the response against a Spectralon
28 reflectance panel, which provides near 100% reflectance over all the 0.3–2.5- μm wavelengths.

29 As outlined in Smith *et al.* (2007b), the spectral components of green vegetation, senesced or non-
30 photosynthetic vegetation, and charcoal were selected as they are ubiquitous surface characteristics that
31 are present in most global savanna-type fire-prone environments. We acknowledge that application of
32 non-site- and species-specific spectral reflectance curves may not provide an optimal unmixing result.

1 However, these generic spectra were applied as we sought to explore surface components that could
2 potentially be quickly applied to global savanna-type fires to assess post-fire effects. A common spectral
3 component used in spectral mixture analysis is that of the dominant soil. However, we deliberately sought
4 non-soil components, as (1) soils types are highly variable within individual fires, let alone across a series
5 of fires, and (2) soil spectra vary considerably between different soil types and in soils with varying
6 organic or moisture characteristics (Huete and Escadafal 1991; Nagler *et al.* 2000). In the present study,
7 we did not consider shade as an endmember as it does not have a unique spectral reflectance curve, but
8 rather exhibits a range of curves associated with ever-darkened versions of the surface component being
9 shaded. We acknowledge that in future regional fires, using locally collected spectral data would be most
10 appropriate. For example, for a detailed comparison of spectral endmembers from other wildfires, please
11 refer to the FRAMES online resource. FRAMES can be accessed at <http://frames.nbii.gov/> (accessed 26
12 January 2008). The endmembers can be accessed using keywords 'Spectral Library'.

13 Linear spectral unmixing was applied using the *IDL/ENVI* ver 4.2 module with the 'sum to 1'
14 constraint applied (Drake *et al.* 1999), which ensures that all component fractions within a pixel sum to
15 unity, although individual class fractions may be negative or exceed 1. To input the spectral reflectance
16 curves for use in the ITT Visual Information Systems for the *ENVI* software package, it was first
17 necessary to interpolate the spectral data to 1-m steps and then convolve the data with the band spectral
18 response functions of the Landsat 7 sensor (Smith *et al.* 2005). This provides *ENVI* with six values, where
19 each value corresponds to the associated reflectance bands, rather than 2300 continuous values. The
20 wavelength ranges associated with bands 4 and 7 (i.e. the NBR bands) of the Landsat sensors are shown
21 in Fig. 2.

22 Each individual band reflectance, dNBR, and associated fractional cover estimate was then extracted at
23 each plot location using the *ARC* software package (ESRI, Redlands, CA, USA).

24 *Field measurements*

25 Lentile (2004) and Keyser (2007) sampled three ~800-ha study areas that contained a mosaic of fire
26 effects in the north, central, and southern portions of the Jasper Fire perimeter. In June 2001, before the
27 fall of fire-scorched needles, 66 0.28-ha (30-m radius) permanent study sites were established in
28 ponderosa pine stands within the study areas. Within these burned stands, nine sites were located in areas
29 exhibiting evidence of surface fire behavior with low initial post-fire tree mortality; 24 sites were located
30 in ponderosa pine stands exhibiting moderate fire behavior, consisting of surface fire with individual tree
31 torching resulting in moderate initial post-fire tree mortality; and 33 were located in severely burned
32 ponderosa pine stands where all trees were killed. Each site consisted of three 0.031-ha (10-m radius)
33 plots. Plots were located at bearings 0°, 135°, and 225° azimuth 20 m from the site center. Study sites

1 were similar in respect to preburn species composition, aspect, slope (5–13%), elevation, and soil type
2 (Lentile 2004; Keyser *et al.* 2006; Lentile *et al.* 2006; Keyser 2007). These data were recently presented
3 in combination with 14 additional aspen study sites within a preliminary assessment of char fraction
4 measures (Smith *et al.* 2007b). Prior studies have observed that pine and aspen plots differ in their
5 spectral properties and response to fire (Brown and Smith 2000; Keyser *et al.* 2005), so the present study
6 excluded the aspen plots from the analysis.

7 On these plots, data on the fire effects on the canopy, boles, and around the bases of individual trees >5
8 cm diameter at breast height (DBH) (1.4 m above soil surface) were collected. Tree survival was
9 calculated based on the proportion of trees surviving the fire compared with trees alive before the fire on
10 each plot (% live tree). Trees with no green foliage were considered dead. Bark thickness was sampled at
11 breast height at two different locations on the bole to compute an average bark thickness per tree. In
12 addition, we measured total tree height and prefire crown base height. Crown base height was measured at
13 the point of branch-bole attachment of the lowest prefire live whorl. We identified prefire crown base
14 height from the position of scorched needles in the case where no foliage consumption occurred and fine
15 branch structure in the case where consumption of needles occurred. Scorched needles were easily
16 distinguishable from non-scorched needles as they were brown or orange in color. Crown injury was
17 measured on individual trees and included the proportion of the prefire live crown that was affected by
18 crown scorch (% crown scorch) or crown consumption (% crown consumption). The cumulative effects
19 of crown scorch and consumption are represented by the total crown fire effects. We measured the
20 percentage of the bole circumference scorched below 30 cm above the soil to the nearest 5% as an
21 indicator of stem damage (basal scorch %). We measured the percentage of the bole circumference
22 charred below 30 cm to the nearest 5% as an indicator of stem and cambial damage (basal char %).
23 Charred bark was distinguished from scorched bark as it was metallic black in color (similar to the color
24 and texture of charcoal) and was eroded to the point that the bark no longer contained grooves or furrows,
25 whereas scorched bark was completely intact and black or gray in color. We measured the percentage of
26 the bole circumference scorched at 100 cm to the nearest 5% as an indicator of stem damage (bole scorch
27 at 1 m %). For further details, see Lentile (2004), Keyser *et al.* (2006), and Keyser (2007). A typical
28 ponderosa pine stand 1 year following the Jasper Fire is shown in Fig. 3.

29 Following Ryan and Noste (1985), the percentage low, moderate, and high ground char in a 1-m radius
30 area around the base of each tree was measured. Line transects (30 m) were laid at 90° and 270° bearings
31 with the site center as the midpoint. Depths of forest floor litter/duff depth and the percentage low,
32 moderate, and high ground char (Ryan and Noste 1985) for a 0.025-m² surface area were measured at 30
33 points at 2-m intervals along these transects. An index of burn severity (BI) was defined as a weighted

1 sum of the product of the proportion of the ground area charred, with the degree of char scaled from low
2 (1) to high (3). Within each of the individual tree plots, we characterized the forest floor and soil effects.
3 As these measurements were not originally intended for the purposes of a char or remote sensing analysis,
4 we assigned a proportion low, moderate, and high burn severity based on widely applied descriptions of
5 field severity (Ryan and Noste 1985). To calculate the Burn Index, we multiplied the % low times 100;
6 the % moderate times 200; and the % high times 300; and then summed these scores. BI was calculated
7 within a 1-m radius area around the base of each tree within plots (Total BI 1 m tree), and for each of the
8 30 forest-floor points located at 2-m intervals along the transect (Floor BI). At six additional points offset
9 from the transect, samples were collected, and later oven-dried and weighed to estimate forest floor
10 biomass (litter organic weight). This same suite of measurements was conducted at nine adjacent,
11 unburned sites in order to provide an estimate of fire-induced changes on the forest floor.

12 *Statistical analysis*

13 Simple linear regressions were applied to assess the predictive ability of the fractions and NBR indices to
14 predict the 1-year post-fire effects. The regressions were tested for significance at the 95% level and the
15 standard error calculated (Tables 2–4). Linear regressions were also determined to assess the ability of 1-
16 year post-fire dNBR to indicate 1-year post-fire effects. Furthermore, both single and multiple linear
17 regressions were applied within the *SPSS* software package (SPSS Inc., Chicago, Illinois) to assess the
18 degree of redundancy in the information provided by each of the fractional cover measures in predicting
19 the 1-year post-fire effect measures (Table 5).

20 *General limitations of pixel-based remote sensing methods and SMA*

21 Two main limitations exist when relating pixel-based remotely sensed data to ecological effects, namely
22 assumed independence of neighboring pixels and that the observed signal only represents the spectral
23 contributions of the components that occupy that specific pixel (Townshend 1981; Cracknell 1998). These
24 problems arise from the arbitrary definition of a pixel: a typically square unit somewhat related to the
25 circular field of view of the optical sensor that will likely not have edges matching up with actual
26 ecological or physical boundaries (Foody *et al.* 1997; Cracknell 1998). As a result, unless the relative
27 contributions of surface components are inferred, assumption of a homogeneous pixel can lead to
28 classification and subsequent model propagation errors (Foody *et al.* 1997).

29 The principal assumption specific to spectral mixture analysis is that the combined reflectance of the
30 pixel is a ‘linear combination’ of the reflectances of the individual pixel components, weighted by the
31 relative proportion they occupy in the pixel (Drake and White 1991; Settle and Drake 1993; Foody *et al.*
32 1997). In general, the linear mixing assumption is valid when the surface components exist as sufficiently

1 large discrete areas that are optically thick (i.e. no light is transmitted through to a lower land-cover type),
2 such that the photons only interact with a single land-cover type (Drake *et al.* 1999; Qin and Gerstl 2000).
3 However, non-linear mixing does occur in environmental applicants (Borel and Gerstl 1994), but can be
4 minimized by using visible and short-wave infrared wavelengths (0.3–2.5 μm) associated with low
5 canopy transmissions (Drake *et al.* 1999).

6 Clark and Lucey (1983) observed that mixtures containing dark and light components mix non-linearly
7 owing to the multiple scatters being preferentially reflected by lighter surfaces and absorbed by darker
8 surfaces. Such examples lead to the appearance of greater proportions of darker components within a
9 pixel than actually exist (Foody *et al.* 1997). However, as noted earlier in the text, post-fire mixtures
10 including the extreme ‘real’ ecological examples of white mineral ash (>70% reflectance) and black char
11 (<20% reflectance) have been shown to be approximated to a generally linear mixture model (Smith *et al.*
12 2005). A further limitation of spectral mixture analysis is that unique spectral endmembers are required
13 for each land cover class and the results are highly sensitive to how those endmembers are selected
14 (Atkinson *et al.* 1997; Theseira *et al.* 2003). Errors can also arise when endmember spectra are missing or
15 incorrectly defined.

16 **Results**

17 *General description of post-fire effects*

18 The direct and cumulative effects of fire on ponderosa pine trees were much greater on high-severity than
19 on low- or moderate-severity sites (Table 1). Approximately 1, 22, and 100% of trees that were alive
20 before the fire were killed in pine stands burned by low-, moderate-, and high-severity fire. The entire
21 bole was scorched, and canopy foliage and small branches were completely consumed in areas of high-
22 severity fire. Bole and crown scorch was more extensive on moderate- than on low-severity sites.
23 Approximately 75% of the crown was scorched or consumed on moderate-severity compared with ~20%
24 on low-severity sites. On average, 80% of the base of each tree bole was scorched on low- and moderate-
25 severity sites, and 2.2 times more char was found on the base of each tree on moderate-severity sites
26 relative to low-severity sites. Post-fire bark thickness (s.e.) was 1.5 (0.1), 1.2 (0.1), and 0.7 (0.1) cm in
27 low-, moderate-, and high-severity sites.

28 Fire effects on the forest floor were most substantial in areas of high burn severity where litter and duff
29 were almost completely consumed. Total ‘BI 1 m tree’ was 141 on low-, 223 on moderate-, and 290 on
30 high-severity sites on a BI scale of 100 to 300. Floor BI was 119 on low-, 186 on moderate-, and 246 on
31 high-severity sites on the same BI scale. Average litter depths (s.e.) were 1.2 (0.3), 0.5 (0.2), and 0.2 (0.1)
32 cm on low-, moderate-, and high-severity compared with 4.8 (0.5) cm on unburned sites. Fire reduced
33 litter depths by ~76, 91, and 97% on low-, moderate-, and high-severity sites 1 year after fire. On average,

1 there were 2.3 and 6.6 times more duff on unburned sites than on sites burned with low and moderate
2 severity. No duff remained 1 year after fire on high-severity sites. Litter organic weights (s.e.) were 1266
3 (264), 684 (173), 459 (93), and 82 (45) g m⁻² in unburned, low-, moderate-, and high-severity sites.

4 *Prediction of 1-year post-fire effects*

5 Fractional char cover either equaled or outperformed all other remote metrics as a predictor of 1-year
6 post-fire effects, except for the relation between 1-year post-fire dNBR and percentage live tree ($r^2 =$
7 0.74) (Tables 2, 3, 4). Each remote metric poorly characterized crown scorch, with the char fraction and
8 dNBR methods producing statistically significant but poor relationships ($r^2 < 0.17$, $P < 0.031$). The 1-year
9 post-fire crown scorch on trees will likely be similar to scorch measured immediately post-fire. The
10 results illustrate that fractional char cover is a reasonable predictor of several canopy and subcanopy
11 measures (but not all: Tables 2, 3). In terms of canopy measures, fractional char cover produced
12 reasonable predictions of % live trees ($r^2 = 0.69$) and % crown consumption ($r^2 = 0.65$), and was
13 comparable with the results obtained using the 1-year post-fire dNBR measure. However, the improved
14 performance of the 1-year post-fire dNBR measure might be expected because both the imagery and field
15 measures are effectively coincident measures of the same condition. In terms of subcanopy measures,
16 fractional char cover strongly predicted % bole scorch ($r^2 = 0.72$) and weight of organic litter ($r^2 = 0.71$),
17 while fractional green cover produced weaker but reasonable predictions ($r^2 = 0.60$ and $r^2 = 0.64$,
18 respectively). Both the char and green cover fraction predictions surpassed the immediate post-fire dNBR
19 predictions of these 1-year post-fire effects.

20 When we compared the immediate post-fire NBR with immediate post-fire dNBR, inclusion of the
21 prefire data did not improve the prediction of the 1-year post-fire measures, except for marginal
22 improvements in predicting % bole scorch ($r^2 = 0.53$). As it is not possible for any spectral ratio like
23 dNBR to outperform a regression containing both of the two component bands (Lawrence and Ripple
24 1998), which in this case are NBR_{pre} and $NBR_{immediate\ post-fire}$, we think that the majority of the useful
25 predictive information is contained within the $NBR_{immediate\ post-fire}$ data, thus potentially limiting the need
26 for using NBR_{pre} to predict post-fire effects. These results concur with prior studies that relate both NBR
27 and dNBR with CBI in forest and woodland environments (Epting *et al.* 2005) and are further supported
28 by the correlation between the $dNBR_{immediate}$ data and each of NBR_{pre} ($r = 0.93$) and $NBR_{immediate}$ ($r = 0.34$)
29 data.

30 Immediate dNBR was a reasonable predictor of % live trees ($r^2 = 0.53$), % bole scorch ($r^2 = 0.50$), and
31 weight of organic litter ($r^2 = 0.59$). Although 1-year post-fire dNBR outperformed immediate post-fire
32 dNBR for most of the post-fire effects, the immediate measure did produce a marginally improved
33 prediction, in terms of the coefficient of determination, of the depth of the 1-year post-fire litter. Again,

1 these general results of higher coefficient of determination in using the 1-year post-fire dNBR are as we
2 expected, as this index incorporates data that are effectively coincident with the 1-year post-fire field
3 measures. Furthermore, in a comparison with the preliminary data presented in [Smith et al. \(2007b\)](#), we
4 did observe that removal of the 14 aspen plots from the regressions considerably improved the r^2 values
5 and reduced the variability in the results. These results confirm our premise that coniferous cover types
6 should be separately evaluated from deciduous cover types. For instance, the example of char fraction v .
7 litter organic weight improved from $r^2 = 0.55$ (s.e. = 4.78) ([Smith et al. 2007b](#)) to $r^2 = 0.71$ (s.e. = 3.99)
8 (present study). Similar improvement was observed using immediate dNBR data ([Smith et al. 2007b](#)).

9 **Discussion**

10 *Remote prediction of post-fire effects*

11 To predict implies to 'forecast a situation that is yet to occur'. Therefore, it is not appropriate to predict
12 field measures of post-fire effects with 1-year post-fire dNBR, as this is effectively measured
13 concurrently with the 1-year post-fire field measures. Thus, the regressions herein were presented solely
14 for the purpose of determining the 'potential inference ability' of the 1-year post-fire dNBR, not to
15 forecast conditions 2, 5, or 10 years after fire ([Table 4](#)). Timely prediction of field-based ecological
16 indicators of 1-year post-fire effects must instead be achieved through the use of methods measured either
17 during or immediately following the fire event, as it is not practical to wait a year before making a 1-year
18 post-fire prediction.

19 These results demonstrate that immediate dNBR was a poorer indicator of 1-year post-fire ecological
20 effects than char cover fraction. Furthermore, immediate dNBR was in many cases a poorer indicator of
21 1-year post-fire ecological effects. The ability of immediate dNBR to reasonably predict 1-year post-fire
22 % live crown is because the index is sensitive to the quantity of green and senesced vegetation
23 (highlighted by Landsat band 4 values), and to a lesser extent, the quantity (and moisture content) of
24 exposed soil or char cover (highlighted by Landsat band 7 values) present within the immediate post-fire
25 pixel ([Eva and Lambin 1998a, 1998b](#); [Stroppiana et al. 2002](#); [Smith et al. 2005](#); [Lentile et al. 2006](#); [Key
26 2006](#)). In instances where either the canopy component is relatively untouched or completely consumed
27 (e.g. in a stand-replacing fire), the 1-year post-fire canopy conditions may still represent the same relative
28 amount of green vegetation. In contrast, the understorey immediately following the fire will be dominated
29 by char and mineral ash, which 1 year later will have been removed by wind and water or occluded by
30 vegetation regrowth or scorched needlecast ([Smith and Hudak 2005](#)). As such, the contribution of band 7
31 to the 1-year post-fire dNBR might simply be adding noise to the predictions of the subcanopy fire
32 effects.

1 The majority of spectral indices are designed to highlight complementary changes in two or more
2 bands. For NBR-based indices, we expect a lowering of reflectance in band 4 between the pre- and post-
3 fire dates to correspond to a complementary increase in the value of band 7. Therefore, we would expect a
4 significant correlation between these band differences: i.e. $(b_{4_{\text{post}}} - b_{4_{\text{pre}}})$ and $(b_{7_{\text{post}}} - b_{7_{\text{pre}}})$. To test this
5 assumption, correlations between the differences in band value pairs were calculated and are presented in
6 **Table 6**. Although the correlations of both band 4 and band 7 differences were both significant at the 95%
7 confidence interval, the correlation between the band 4 difference and the band 7 difference was
8 noticeably lower when using the 1-year post-fire image value.

9 We further calculated a measure of the signal to noise ratio (SNR) for each of the band 4 and band 7
10 difference pairs (**Table 6**). This measure was determined by calculating both the mean and standard
11 deviations of the band 4 and band 7 values of all the 66 plots for each image. In a similar manner to the
12 SNR-based M-statistic (**Pereira 1999**), SNR was calculated by:

$$13 \quad SNR_{band(int)} = \frac{|\mu_f - \mu_i|}{\sigma_f + \sigma_i} \quad (2)$$

14 where $band(int)$ denotes the temporal interval between the prefire condition and either the immediate or
15 1-year post values for the band of interest (in this case bands 4 or 7); μ and σ denote the mean and
16 standard deviations of all the 66 fire-affected plots for the band of interest; and i denotes the prefire data,
17 and f denotes the post-fire image values.

18 The $SNR_{band\ 4(immediate)} = 3.2$, whereas the $SNR_{band\ 4(1-year)} = 1.4$. The SNR for band 7 changed in a similar
19 manner, with $SNR_{band\ 7(immediate)} = 0.75$ and $SNR_{band\ 7(1-year)} = 0.47$. It is clear that both the band 4 and band
20 7 values have notable decreases in their SNR when using the 1-year post-fire image values compared with
21 the usage of the immediate post-fire imagery. These results lend support to the proposition that the
22 application of 1-year post-fire band values may not be optimal for assessing post-fire effects.

23 These effects would be less pronounced where canopy closure remains high (unburned or low degree
24 of fire effects) or in stand-replacing fires where the understory vegetation could be replaced by bare soil.
25 The unexpected ability of the immediate NBR and dNBR to predict the 1-year post-fire measure of
26 organic litter weight could be an indirect effect of the combined impact of scorched canopies with
27 extensive surface fires. In such fires, we would expect the surface material to be consumed and scorched
28 needles to fall as new litter for the 1-year post-fire measurement, as highlighted in several studies
29 (**Robichaud and Brown 2000**; **Pannkuk and Robichaud 2003**; **Robichaud 2004**). In contrast, low-severity
30 fires and stand-replacing fires would result in high and low organic litter weights, respectively.

1 Measures of both the immediate post-fire char and green vegetation fractions are good predictors of
2 several 1-year post-fire canopy and several subcanopy measures. Most notable, several of these 1-year
3 post-fire measures appear to be potential surrogates of fire intensity. Specifically, % bole scorch can be
4 considered a proxy to flame length, while scorch to 1 m and organic litter weight might each relate to rate
5 of spread, and average bark thickness might similarly relate to fire duration. Therefore, these fractional
6 measures have the potential to inform managers regarding tree mortality (via canopy condition and
7 average bark thickness) and may provide viable proxies of fire intensity to Burned Area Emergency
8 Response (BAER) teams tasked with deciding where post-fire mitigation efforts are needed.

9 Of the fractions considered, char fraction was marginally better over green fraction for predicting
10 several metrics of fire severity 1 year after fire. This leads to the question, ‘Are these two fractions
11 providing redundant or complimentary information?’ This is an important distinction, as the degree to
12 which the information is unique to each fraction would indicate whether or not a composite metric of the
13 two fractions could be used to produce an improved burn severity remote sensing method. To answer this
14 question, we first assessed the correlation between the component fractions (Fig. 4), and second assessed
15 the variability that each fraction term accounted for in the predictions of each of the 1-year post-fire field
16 measures (Table 5).

17 The correlations of the green fraction with both the char and the senesced vegetation fractions were
18 significant and high ($r = \sim 0.85$); however, the char and brown fractions were only poorly correlated ($r =$
19 0.46) (Fig. 4). Although the fractions are relative, the brown (senesced) component produced ‘negative’
20 fractions, suggesting that this term is not optimal and perhaps is accounting for the lack of a specific soil
21 endmember. However, without detailed soil maps and spectral reflectance curves for each soil type, it
22 would be difficult to replace the brown endmember with representative soil endmembers.

23 In multiple linear regressions of the component fractions against the organic litter weight, only the char
24 fraction and the combination of the char with the green fraction produced significant results at the 95%
25 confidence level. In the example of the organic litter weight measure, the char fraction alone accounted
26 for 71% of the variance, with the addition of the green fraction only accounting for an additional 2% of
27 the variability. In each of the two cases of bole scorch and percentage live tree, addition of the green
28 fraction within the regression was not significant and only accounted for an additional 2% variance
29 explained over the char fraction. These results limit the likelihood that a combination index using both the
30 char and green vegetation fractions would provide significantly improved predictions over just the char
31 fraction alone.

1 *Management and science implications*

2 Immediate post-fire assessments, particularly those that utilize only immediate post-fire dNBR
3 techniques, can be misleading. The post-fire environment will change greatly within 1 year, some aspects
4 of which may be predictable whereas others may be related to local and regional climate. Char fractional
5 cover may be a viable alternative to dNBR to predict longer-term post-fire ecological effects, especially
6 when the prediction is needed in a timely manner. For instance, BAER teams must make post-fire
7 rehabilitation treatment recommendations within 7 days following fire containment. Second, post-fire
8 recovery generally is more rapid in less severely burned areas. However, commonly applied dNBR
9 techniques provide very little information about the effects of fire on the forest floor and soil. As such,
10 char fraction is particularly useful in fire regimes where some, but not all, of the overstorey tree and shrub
11 canopy is consumed. The mosaic of relatively small patches of severely burned forests interspersed within
12 less severely burned forests, a common signature of surface fires and mixed-severity fire regimes, exerts a
13 strong influence on post-fire landscape heterogeneity and rates of recovery. In some extensive areas of
14 high-severity fire, post-fire vegetation dynamics may not follow the same trajectory as less severely
15 burned areas, and a cover type conversion from forests to shrubs or meadows may occur.

16 From a management perspective, streamlined assessment of fire effects on overstorey, understorey, and
17 forest floor environments can be used to predict areas likely to develop vegetation structure different from
18 prefire conditions, and will facilitate post-fire monitoring and mitigation ([Lentile et al. 2007b](#)).
19 Identification of desirable attributes of fire behavior and positive post-fire effects may improve restoration
20 strategies. For example, recognition of initial fire effects likely to result in tree death may facilitate
21 selection of which trees to salvage-harvest or leave as potential seed sources. In some burned areas,
22 reforestation or seeding are probably unnecessary and could interfere with natural successional dynamics.
23 Furthermore, severely burned areas with low rates of recovery may indicate areas that require immediate
24 attention or are highly vulnerable to displacement of native flora by invasive species. If a cover type
25 conversion from ponderosa pine to shrub-dominated communities is desirable for wildlife habitat
26 diversity, then large patches of high severity may lend themselves to this objective. Longer-interval, large
27 fire events, such as the Jasper Fire, may be critical in maintaining landscape heterogeneity and diversity.
28 Openings in a previously dense, closed-canopy forest may represent a desirable departure from prefire
29 conditions and a return of some attributes of historical landscape function. Rapid landscape
30 characterization that can be mechanistically related to ground measures of post-fire ecosystem condition
31 may provide much needed management guidance and decision support following large fire events.

32 The post-fire effects measured in the field typically reflect fine-scale processes, but also impact coarse
33 spatial (watershed to regional) and temporal (decadal) scales. For such measures to be applicable in

1 describing ecosystem recovery and condition across a range of scales and ecosystems, they should
2 physically relate to pools and fluxes of biophysical variables (e.g. the carbon and water cycles). Although
3 the mechanistic relations between fire effects and the carbon and water cycles are not currently well
4 defined, the results of the present study support the argument that cover fractions are potentially versatile
5 measures of post-fire ecological impact that also influence the terrestrial carbon and water cycles (Table
6 7).

7 **Conclusions**

8 The previous study of Smith *et al.* (2007b) reported for the Jasper Fire that the modeled estimate of the %
9 char was a slightly improved predictor over immediate dNBR of two 1-year post-fire field measures,
10 namely the % live tree and the organic litter weight. However, given the conflicting findings of Hudak *et al.*
11 (2007) with respect to fractional green cover estimates and the recognition that the initial study
12 incorrectly analyzed both ponderosa pine and aspen stands together, further analysis of this dataset was
13 warranted. Specifically, assessment to investigate whether the char and other modeled estimates of the
14 immediate post-fire fractional covers (green and senesced vegetation) could also predict an expanded
15 variety of both canopy (four) and subcanopy (nine) post-fire effects. These field measures were selected
16 based on whether they could provide a reasonable bridge between the fire intensity and the fractional
17 cover estimates. An investigation was also included of whether a combination of different fractional cover
18 estimates could act as improved predictor of the post-fire effects, or whether the different fractional cover
19 estimates account for the same variability. For the sake of completeness with contemporary remote
20 sensing post-fire effects research, the present study further investigated whether the dNBR indices, both
21 immediate and 1-year post-fire, were improved predictors when compared with the fractional cover
22 estimates.

23 The results demonstrated that although the char cover fraction either equaled or outperformed all other
24 immediate measures in predicting 1-year post-fire effects, the green fractional cover was a reasonable
25 predictor for several of the post-fire measures. Although the char and green cover fractions provided
26 improved predictions of the 1-year post-fire effects over the immediate post-fire NBR and dNBR
27 measures, predictions incorporating both the char and green fractional covers only accounted for ~2%
28 more variability than that achieved using the char fraction cover alone. This result combined with the
29 ineffectiveness of the brown fractional cover highlight the limited utility for a combination approach
30 based on several different cover metrics. The comparison of immediate post-fire NBR with dNBR
31 showed that the inclusion of the prefire NBR data did not provide any notable improvement in the
32 predictions of the 1-year post-fire measures. Therefore, perhaps future studies may not need to consider
33 prefire imagery in order to predict several 1-year post-fire canopy and subcanopy effects.

1 Although in the present study, application of the char and green fractional covers to predict 1-year
2 post-fire effects were an improvement over NBR and dNBR, we do not suggest that this approach or that
3 of any other spectral index currently existing will be a panacea for evaluating burn severity in all fire-
4 affected environments (e.g. savannah grasslands, temperate forests, boreal forests, woodlands, chaparral,
5 scrublands). In contrast, it is more likely that a suite of methods will need to be identified, where each
6 separate method in this suite will be optimal for predicting >1 year post-fire effects in a single
7 environment. To enable robust national and global burn severity products, further research is warranted to
8 identify and evaluate these methods.

9 The principal limitation of the current study is that it only represents information from a single wildfire
10 at one point in time. Research is clearly warranted to repeat this analysis on data collected from fires 5,
11 10, or even 20 years post-fire, to evaluate the potential for inferences from immediate post-fire remote
12 sensing data to predict long-term ecological responses to fire, such as succession processes and carbon
13 accumulation. Future research should also evaluate how changes in surface cover fractions relate to both
14 these differenced indices and to changes in remotely sensed fractional cover. This could be achieved via
15 the analysis of prescribed fires, where it is possible to collect information on the prefire fractional cover
16 of flammable and non-flammable materials.

17 **Acknowledgements**

18 The present fieldwork component of the current study was funded by the Black Hills National Forest through In-
19 Service Agreement No. 0203-01-007, Monitoring Fire Effects and Vegetation Recovery on the Jasper Fire, Black
20 Hills National Forest, South Dakota to Rocky Mountain Research Station and Colorado State University. The
21 subsequent research was supported in part by funds provided by the Rocky Mountain Research Station, Forest
22 Service, US Department of Agriculture (03-JV-11222065-279) and the USDA/USDI Joint Fire Science Program
23 (Projects 03-2-1-02 and 05-4-1-07). We thank the Associate Editor, Mark Cochrane and the anonymous reviewers
24 whose comments greatly improved the present manuscript.

25 **References**

- 26 <jrn>Atkinson PM, Cutler MEJ, Lewis H (1997) Mapping sub-pixel proportional land cover with AVHRR imagery.
27 *International Journal of Remote Sensing* 18, 917–935. doi:10.1080/014311697218836</jrn>
- 28 <other>Bobbe T, Finco MV, Quayle B, Lannon K (2001) Field measurements for the training and validation of
29 burn severity maps from spaceborne, remotely sensed imagery. Final Project Report, Joint Fire Sciences Program
30 2001-2</other>
- 31 <jrn>Borel CC, Gerstl SAW (1994) Non-linear spectral mixing models for vegetative and soil surfaces. *Remote*
32 *Sensing of Environment* 47, 403–416. doi:10.1016/0034-4257(94)90107-4</jrn>

- 1 <jrn>Brewer CK, Winne JC, Redmond RL, Opitz DW, Magrich MV (2005) Classifying and mapping wildfire
2 severity: a comparison of methods. *Photogrammetric Engineering and Remote Sensing* **71**(11), 1311–1320.</jrn>
- 3 <other>Brown JK, Smith JK (Eds) (2000) Wildland fire in ecosystems: effects of fire on flora. USDA Forest
4 Service, Rocky Mountain Research Station, General Technical Report RMRS-GTR-42. (Ogden, UT)</other>
- 5 <jrn>Chen X, Vierling L, Rowell E, DeFelice T (2004) Using Lidar and effective LAI to evaluate IKONOS and
6 Landsat 7 ETM+ vegetation estimates in a ponderosa pine forest. *Remote Sensing of Environment* **91**, 14–26.
7 [doi:10.1016/j.rse.2003.11.003](https://doi.org/10.1016/j.rse.2003.11.003)</jrn>
- 8 <jrn>Clark RN, Lucey PG (1983) Spectral properties of ice–particulate mixtures and implications for remote
9 sensing 1. Intimate mixtures. *Journal of Geophysical Research* **89**(B7), 6341–6348.
10 [doi:10.1029/JB089iB07p06341](https://doi.org/10.1029/JB089iB07p06341)</jrn>
- 11 <jrn>Cochrane MA, Souza CM (1998) Linear mixture model classification of burned forests in the Eastern Amazon.
12 *International Journal of Remote Sensing* **19**(17), 3433–3440. [doi:10.1080/014311698214109](https://doi.org/10.1080/014311698214109)</jrn>
- 13 <jrn>Cocke AE, Fulé PZ, Crouse JE (2005) Comparison of burn severity assessments using differenced Normalized
14 Burn Ratio and ground data. *International Journal of Wildland Fire* **14**(2), 189–198.
15 [doi:10.1071/WF04010](https://doi.org/10.1071/WF04010)</jrn>
- 16 <jrn>Cracknell AP (1998) Synergy in remote sensing – what’s in a pixel? *International Journal of Remote Sensing*
17 **19**(11), 2025–2047. [doi:10.1080/014311698214843](https://doi.org/10.1080/014311698214843)</jrn>
- 18 <jrn>Crockford RH, Richardson DP (2000) Partitioning of rainfall into throughfall, stemflow and interception:
19 effect of forest type, ground cover and climate. *Hydrological Processes* **14**, 2903–2920. [doi:10.1002/1099-1085\(200011/12\)14:16/17<2903::AID-HYPI26>3.0.CO;2-6](https://doi.org/10.1002/1099-1085(200011/12)14:16/17<2903::AID-HYPI26>3.0.CO;2-6)</jrn>
- 20
- 21 <jrn>De Santis A, Chuvieco E (2007) Burn severity estimation from remotely sensed data: performance of
22 simulation versus empirical models. *Remote Sensing of Environment* **108**, 422–435.
23 [doi:10.1016/j.rse.2006.11.022](https://doi.org/10.1016/j.rse.2006.11.022)</jrn>
- 24 <jrn>Doerr SH, Cerda A (2005) Fire effects on soil system functioning: new insights and future challenges.
25 *International Journal of Wildland Fire* **14**, 339–342. [doi:10.1071/WF05094](https://doi.org/10.1071/WF05094)</jrn>
- 26 <edb>Dozier J, Strahler AH (1983) Ground investigations in support of remote sensing. In ‘Manual of Remote
27 Sensing’. (Eds RN Colwell, DS Simonett, GA Thorley) pp. 959–986. (American Society of Photogrammetry:
28 Falls Church, VA)</edb>
- 29 <jrn>Drake NA, Mackin S, Settle JJ (1999) Mapping vegetation, soils, and geology in semiarid shrublands using
30 spectral matching and mixture modeling of SWIR AVIRIS imagery. *Remote Sensing of Environment* **68**, 12–25.
31 [doi:10.1016/S0034-4257\(98\)00097-2](https://doi.org/10.1016/S0034-4257(98)00097-2)</jrn>
- 32 <conf>Drake NA, White K (1991) Linear mixture modelling of Landsat Thematic Mapper data for mapping the
33 distribution and abundance of gypsum in the Tunisian Southern Atlas. In ‘Spatial Data 2000: Proceedings of a

1 Joint Conference of the Photogrammetric Society, the Remote Sensing Society, the American Society for
2 Photogrammetry and Remote Sensing, Christ Church, Oxford'. (Ed. I Dowman) pp. 168–177.</conf>

3 <jrn>Elvidge CD (1990) Visible and near-infrared reflectance characteristics of dry plant materials. *International*
4 *Journal of Remote Sensing* **11**, 1775–1795. [doi:10.1080/01431169008955129](https://doi.org/10.1080/01431169008955129)</jrn>

5 <jrn>Epting J, Verbyla D, Sorbel B (2005) Evaluation of remotely sensed indices for assessing burn severity in
6 interior Alaska using Landsat TM and ETM+. *Remote Sensing of Environment* **96**, (3–4), 328–339. </jrn>

7 <jrn>Eva H, Lambin EF (1998a) Burnt area mapping in Central Africa using ATSR data. *International Journal of*
8 *Remote Sensing* **19**(18), 3473–3497. [doi:10.1080/014311698213768](https://doi.org/10.1080/014311698213768)</jrn>

9 <jrn>Eva H, Lambin EF (1998b) Remote sensing of biomass burning in tropical regions: sampling issues and
10 multisensor approach. *Remote Sensing of Environment* **64**, 292–315. [doi:10.1016/S0034-4257\(98\)00006-6](https://doi.org/10.1016/S0034-4257(98)00006-6)</jrn>

11 <jrn>Foody GM, Lucas RM, Curran PJ, Honzak M (1997) Non-linear mixture modelling without end-members
12 using an artificial neural network. *International Journal of Remote Sensing* **18**(4), 937–953.
13 [doi:10.1080/014311697218845](https://doi.org/10.1080/014311697218845)</jrn>

14 <jrn>Foody GM (2000) Estimation of sub-pixel land cover composition in the presence of untrained classes.
15 *Computers & Geosciences* **26**(4), 469–478. [doi:10.1016/S0098-3004\(99\)00125-9](https://doi.org/10.1016/S0098-3004(99)00125-9)</jrn>

16 <jrn>Goforth BR, Graham RC, Hubbert KR, Zanner CW, Minnich RA (2005) Spatial distribution and properties of
17 ash and thermally altered soils after high-severity forest fire, southern California. *International Journal of*
18 *Wildland Fire* **14**, 343–354. [doi:10.1071/WF05038](https://doi.org/10.1071/WF05038)</jrn>

19 <jrn>Holden Z, Smith AMS, Morgan P, Rollins MG, Gessler PE (2005) Evaluation of novel thermally enhanced
20 spectral indices for mapping fire perimeters and comparisons with fire atlas data. *International Journal of Remote*
21 *Sensing* **26**(21), 4801–4808. [doi:10.1080/01431160500239008](https://doi.org/10.1080/01431160500239008)</jrn>

22 <edb>Hudak AT, Morgan P, Bobbitt M, Lentile L (2007a) Characterizing stand-replacing harvest and fire
23 disturbance patches in a forested landscape: a case study from Cooney Ridge, Montana. In ‘Understanding Forest
24 Disturbance and Spatial Patterns: Remote Sensing and GIS Approaches’. (Eds MA Wulder and SE Franklin), pp.
25 209–231 (Taylor & Francis: London)</edb>

26 <jrn>Hudak AT, Morgan P, Bobbitt MJ, Smith AMS, Lewis SA, Lentile LB, Robichaud PR, Clark JT, McKinley
27 RA (2007b) The relationship of multispectral satellite imagery to immediate fire effects. *Fire Ecology* **3**, 64–
28 90.</jrn>

29 <jrn>Huete AR, Escadafal R (1991) Assessment of biophysical soil properties through spectral decomposition
30 techniques. *Remote Sensing of Environment* **35**, 149–159. [doi:10.1016/0034-4257\(91\)90008-T](https://doi.org/10.1016/0034-4257(91)90008-T)</jrn>

31 <jrn>Key CH (2006) Ecological and sampling constraints on defining landscape fire severity. *Fire Ecology* **2**(2),
32 178–203.</jrn>

- 1 <edb>Key CH, Benson NC (2006) Landscape assessment: ground measure of severity, the Composite Burn Index;
2 and remote sensing of severity, the Normalized Burn Ratio. In 'FIREMON: Fire Effects Monitoring and
3 Inventory System'. (Eds DC Lutes, RE Keane, JF Caratti, CH Key, NC Benson, S Sutherland, LJ Gangi) USDA
4 Forest Service, Rocky Mountain Research Station General Technical Report, RMRS-GTR-164-CD, pp. LA1–51.
5 (Ogden, UT).</edb>
- 6 <ths>Keyser TL (2007) Changes in forest structure, community composition, and development in ponderosa pine
7 forests following a mixed-severity wildfire in the Black Hills, SD, USA. PhD thesis, Colorado State
8 University.</ths>
- 9 <jrn>Keyser TL, Smith FW, Lentile LB, Shepperd WD (2006) Modeling post-fire mortality of ponderosa pine
10 following a mixed-severity wildfire in the Black Hills: the role of tree morphology and direct fire effects. *Forest
11 Science* **52**, 530–539.</jrn>
- 12 <jrn>Keyser TL, Smith FW, Shepperd WD (2005) Trembling aspen response to a mixed-severity wildfire in the
13 Black Hills, South Dakota, USA. *Canadian Journal of Forest Research* **35**, 2679–2684. doi:10.1139/x05-
14 180</jrn>
- 15 <jrn>Landmann T (2003) Characterizing sub-pixel Landsat ETM+ fire severity on experimental fires in the Kruger
16 National Park, South Africa. *South African Journal of Science* **99**, 357–360.</jrn>
- 17 <jrn>Lawrence RL, Ripple WJ (1998) Comparisons among vegetation indices and bandwise regression in a highly
18 disturbed, heterogeneous landscape: Mount St Helens, Washington. *Remote Sensing of Environment* **64**, 91–102.
19 doi:10.1016/S0034-4257(97)00171-5</jrn>
- 20 <ths>Lentile LB (2004) Causal factors and consequences of mixed-severity fire in Black Hills ponderosa pine
21 forests. PhD Thesis, Colorado State University.</ths>
- 22 <jrn>Lentile LB, Smith FW, Shepperd WD (2005) Patch structure, fire-scar formation and tree regeneration in a
23 large mixed-severity fire in the South Dakota Black Hills, USA. *Canadian Journal of Forest Research* **35**, 2875–
24 2885. doi:10.1139/x05-205</jrn>
- 25 <jrn>Lentile LB, Smith FW, Shepperd WD (2006a). The influence of topography and forest structure on patterns of
26 mixed-severity fire in the South Dakota Black Hills. *International Journal of Wildland Fire* **15**(4), 557–566.
27 doi:10.1071/WF05096</jrn>
- 28 <jrn>Lentile LB, Holden ZA, Smith AMS, Falkowski MJ, Hudak AT, Morgan P, Lewis SA, Gessler PE, Benson
29 NC (2006b) Remote sensing techniques to assess active fire and post-fire effects. *International Journal of
30 Wildland Fire* **15**, 319–345. doi:10.1071/WF05097</jrn>
- 31 <other>Lentile LB, Morgan P, Hardy C, Hudak A, Means R *et al.* (2007a) Value and challenges of conducting rapid
32 response on wildland fires. USDA Forest Service, Rocky Mountain Research Station, General Technical Report,
33 RMRS-GTR-193. (Fort Collins, CO)</other>

- 1 <jrn>Lentile LB, Morgan P, Hudak AT, Bobbitt MJ, Lewis SA, Smith AMS, Robichaud PR (2007b). Post-fire burn
2 severity and vegetation response following eight large wildfires across the western US. *Fire Ecology* **3**, 91–
3 108.</jrn>
- 4 <jrn>Lewis SA, Lentile LB, Hudak AT, Robichaud PR, Morgan P, Bobbitt MJ (2007) Mapping ground cover using
5 hyperspectral remote sensing after the 2003 Simi and Old wildfires in Southern California. *Fire Ecology* **3**, 109–
6 128.</jrn>
- 7 <jrn>Lewis SA, Wu JQ, Robichaud PR (2006) Assessing burn severity and comparing soil water repellency,
8 Hayman Fire, Colorado. *Hydrological Processes* **20**, 1–16. [doi:10.1002/hyp.5880](https://doi.org/10.1002/hyp.5880)</jrn>
- 9 <jrn>Litton CM, Ryan MG, Knight DH, Stahl PD (2003) Soil–surface carbon dioxide efflux and microbial biomass
10 in relation to tree density 12 years after a stand-replacing fire in a lodgepole pine ecosystem. *Global Change*
11 *Biology* **9**, 680–696. [doi:10.1046/j.1365-2486.2003.00626.x](https://doi.org/10.1046/j.1365-2486.2003.00626.x)</jrn>
- 12 <jrn>Lopez-Garcia MJ, Caselles V (1991) Mapping burns and natural reforestation using Thematic Mapper data.
13 *Geocarto International* **6**, 31–37. [doi:10.1080/10106049109354290](https://doi.org/10.1080/10106049109354290)</jrn>
- 14 <jrn>Miller JD, Thode AE (2007) Quantifying burn severity in a heterogeneous landscape with a relative version of
15 the delta Normalized Burn Ratio (dNBR). *Remote Sensing of Environment* **109**, 66–80.
16 [doi:10.1016/j.rse.2006.12.006](https://doi.org/10.1016/j.rse.2006.12.006)</jrn>
- 17 <jrn>Miller JD, Yool SR (2002) Mapping forest post-fire canopy consumption in several overstory types using
18 multi-temporal Landsat TM and ETM data. *Remote Sensing of Environment* **82**, 481–496. [doi:10.1016/S0034-
19 4257\(02\)00071-8](https://doi.org/10.1016/S0034-4257(02)00071-8)</jrn>
- 20 <jrn>Morgan P, Hardy CC, Swetnam T, Rollins MG, Long LG (2001) Mapping fire regimes across time and space:
21 understanding coarse and fine-scale fire patterns. *International Journal of Wildland Fire* **10**, 329–342.
22 [doi:10.1071/WF01032](https://doi.org/10.1071/WF01032)</jrn>
- 23 <jrn>Nagler PL, Daughtry CST, Goward SN (2000) Plant litter and soil reflectance. *Remote Sensing of Environment*
24 **71**, 207–215. [doi:10.1016/S0034-4257\(99\)00082-6](https://doi.org/10.1016/S0034-4257(99)00082-6)</jrn>
- 25 <jrn>Odion DC, Hanson CT (2006) Fire severity in conifer forests of the Sierra Nevada, California. *Ecosystems* **9**,
26 1177–1189. [doi:10.1007/s10021-003-0134-7](https://doi.org/10.1007/s10021-003-0134-7)</jrn>
- 27 <jrn>Odion DC, Hanson CT (2007) Fire severity in the Sierra Nevada revisited: conclusions robust to further
28 analysis. *Ecosystems* **11**, 12–15. [doi:10.1007/s10021-007-9113-0](https://doi.org/10.1007/s10021-007-9113-0)</jrn>
- 29 Pannkuk CK, Robichaud PR (2003). Effectiveness of needle cast at reducing erosion after forest fires. *Water*
30 *Resource Research* **39**, 1333–1343. [doi:10.1029/2003WRR002318](https://doi.org/10.1029/2003WRR002318)
- 31 <jrn>Patterson MW, Yool SR (1998) Mapping fire-induced vegetation mortality using Landsat Thematic Mapper
32 data: a comparison of linear transformation techniques. *Remote Sensing of Environment* **65**, 132–142.
33 [doi:10.1016/S0034-4257\(98\)00018-2](https://doi.org/10.1016/S0034-4257(98)00018-2)</jrn>

- 1 <jrn>Pereira JMC (1999) A comparative evaluation of NOAA/AVHRR vegetation indexes for burned surface
2 detection and mapping. *IEEE Transactions on Geoscience and Remote Sensing* **37**(1), 217–226.
3 [doi:10.1109/36.739156](https://doi.org/10.1109/36.739156)</jrn>
- 4 <jrn>Qin W, Gerstl SAW (2000) 3-D scene modelling of semi-arid vegetation cover and its radiation regime.
5 *Remote Sensing of Environment* **74**, 145–162. [doi:10.1016/S0034-4257\(00\)00129-2](https://doi.org/10.1016/S0034-4257(00)00129-2)</jrn>
- 6 <jrn>Robichaud PR (2004) Post-fire rehabilitation: are we learning what works? *Southwest Hydrology* **3**, 20–
7 21.</jrn>
- 8 <conf>Robichaud PR, Brown RE (2000) What happened after the smoke cleared: onsite erosion rates after a
9 wildfire in Eastern Oregon. In ‘Proceedings of the Annual Summer Specialty Conference (Track 2: Wildland
10 Hydrology)’, 30 June–2 July 1999, Bozeman, MT. pp. 419–426. (American Water Resources Association:
11 Herndon, VA)</conf>
- 12 <jrn>Robichaud PR, Lewis SA, Laes DYM, Hudak AT, Kokaly RF, Zamudio JA (2007) Post-fire soil burn severity
13 mapping with hyperspectral image unmixing. *Remote Sensing of Environment* **108**, 467–480.
14 [doi:10.1016/j.rse.2006.11.027](https://doi.org/10.1016/j.rse.2006.11.027)</jrn>
- 15 <jrn>Roy DP, Boschetti L, Trigg SN (2006) Remote sensing of fire severity: assessing the performance of the
16 normalized burn ratio. *IEEE Geoscience and Remote Sensing Letters* **3**(1), 112–116.
17 [doi:10.1109/LGRS.2005.858485](https://doi.org/10.1109/LGRS.2005.858485)</jrn>
- 18 <conf>Ryan KC, Noste NV (1985) Evaluating prescribed fires. In ‘Proceedings of the Symposium and Workshop
19 on Wilderness Fire’, 15–18 November 1983, Missoula MT. (Eds JE Lotan, BM Kilgore, WC Fischer, RW
20 Mutch) USDA Forest Service, Intermountain Forest and Range Experiment Station, General Technical Report
21 INT-GTR-182, pp. 230–238. (Ogden, UT)</conf>
- 22 <jrn>Safford HD, Miller JD, Schmidt D, Roath B, Parsons A (2007) BAER soil burn severity maps do not measure
23 fire effects to vegetation: a comment on Odion and Hanson (2006). *Ecosystems* **11**, 1–11. [doi:10.1007/s10021-](https://doi.org/10.1007/s10021-007-9094-z)
24 [007-9094-z](https://doi.org/10.1007/s10021-007-9094-z)</jrn>
- 25 <jrn>Settle JJ, Drake NA (1993) Linear mixing and the estimation of ground cover proportions. *International*
26 *Journal of Remote Sensing* **14**(6), 1159–1177. [doi:10.1080/01431169308904402](https://doi.org/10.1080/01431169308904402)</jrn>
- 27 <other>Shepperd WD, Battaglia MA (2002) Ecology, silviculture, and management of Black Hills ponderosa pine.
28 USDA Forest Service, Rocky Mountain Research Station, General Technical Report RMRS-97. (Fort Collins,
29 CO)</other>
- 30 <jrn>Smith AMS, Hudak AT (2005) Estimating combustion of large downed woody debris from residual white ash.
31 *International Journal of Wildland Fire* **14**, 245–248. [doi:10.1071/WF05011](https://doi.org/10.1071/WF05011)</jrn>
- 32 <jrn>Smith AMS, Wooster MJ, Drake NA, Dipotso FM, Falkowski MJ, Hudak AT (2005) Testing the potential of
33 multi-spectral remote sensing for retrospectively estimating fire severity in African savannahs environments.
34 *Remote Sensing of Environment* **97**(1), 92–115. [doi:10.1016/j.rse.2005.04.014](https://doi.org/10.1016/j.rse.2005.04.014)</jrn>

- 1 <jrn>Smith AMS, Drake NA, Wooster MJ, Hudak AT, Holden ZA, Gibbons CJ (2007a) Production of Landsat
2 ETM+ reference imagery of burned areas within Southern African savannahs: comparison of methods and
3 application to MODIS. *International Journal of Remote Sensing* **28**, 2753–2775.
4 [doi:10.1080/01431606009547044](https://doi.org/10.1080/01431606009547044)</jrn>
- 5 <jrn>Smith AMS, Lentile LB, Hudak AT, Morgan P (2007b) Evaluation of linear spectral unmixing and Δ NBR for
6 predicting post-fire recovery in a N. American ponderosa pine forest. *International Journal of Remote Sensing*
7 **28**, 5159–5166. [doi:10.1080/0143160701395161](https://doi.org/10.1080/0143160701395161)</jrn>
- 8 <jrn>Stroppiana D, Pinnock S, Pereira JMC, Gregoire JM (2002) Radiometric analysis of SPOT-VEGETATION
9 images for burnt area detection in Northern Australia. *Remote Sensing of Environment* **82**, 21–37.
10 [doi:10.1016/S0034-4257\(02\)00021-4](https://doi.org/10.1016/S0034-4257(02)00021-4)</jrn>
- 11 <jrn>Theseira MA, Thomas G, Sannier CAD (2002) An evaluation of spectral mixture modeling applied to a semi-
12 arid environment. *International Journal of Remote Sensing* **23**, 687–700. [doi:10.1080/0143160010019652](https://doi.org/10.1080/0143160010019652)</jrn>
- 13 <jrn>Theseira MA, Thomas G, Taylor JC, Gemmill F, Varjo J (2003) Sensitivity of mixture modelling to end-
14 member selection. *International Journal of Remote Sensing* **24**(7), 1559–1575.
15 [doi:10.1080/0143160210146631](https://doi.org/10.1080/0143160210146631)</jrn>
- 16 <jrn>Townshend JGR, Huang C, Kalluri SNV, Defries RS, Liang D (2000) Beware of per-pixel characterization of
17 land cover. *International Journal of Remote Sensing* **21**(4), 839–843. [doi:10.1080/01431600210641](https://doi.org/10.1080/01431600210641)</jrn>
- 18 <jrn>Trumbore S (2006) Carbon respired by terrestrial ecosystems – recent progress and challenges. *Global Change*
19 *Biology* **12**(2), 141–153. [doi:10.1111/j.1365-2486.2006.01067.x](https://doi.org/10.1111/j.1365-2486.2006.01067.x)</jrn>
- 20 <jrn>Vafeidis AT, Drake NA (2005) A two-step method for estimating the extent of burnt areas with the use of
21 coarse-resolution data. *International Journal of Remote Sensing* **26**(11), 2441–2459.
22 [doi:10.1080/0143160500034102](https://doi.org/10.1080/0143160500034102)</jrn>
- 23 <jrn>Verstraete MM, Pinty B (1996) Designing optimal spectral indices for remote sensing applications. *IEEE*
24 *Transactions on Geoscience and Remote Sensing* **34**(5), 1254–1265. [doi:10.1109/36.536541](https://doi.org/10.1109/36.536541)</jrn>
- 25 <jrn>van Wageningen JW, Root RR, Key CH (2004) Comparison of AVIRIS and Landsat ETM+ detection
26 capabilities for burn severity. *Remote Sensing of Environment* **92**, 397–408. [doi:10.1016/j.rse.2003.12.015](https://doi.org/10.1016/j.rse.2003.12.015)</jrn>
- 27 <jrn>Wessman CA, Bateson CA, Benning TL (1997) Detecting fire and grazing patterns in tallgrass prairie using
28 spectral mixture analysis. *Ecological Applications* **7**(2), 493–511. [doi:10.1890/1051-0761\(1997\)007\[0493:DEAGPI\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1997)007[0493:DEAGPI]2.0.CO;2)</jrn>
- 29
30 <jrn>Wimberly MC, Reilly MJ (2007) Assessment of fire severity and species diversity in the southern
31 Appalachians using Landsat TM and ETM+ imagery. *Remote Sensing of Environment*.
32 [doi:10.1016/j.rse.2006.03.019](https://doi.org/10.1016/j.rse.2006.03.019)</jrn>

1 Manuscript received 12 July 2007, accepted 21 August 2008

2 **Table 1. Direct fire effects measured on the boles and in crowns of trees after the Jasper Fire**

3 All values are mean \pm standard error ($n = 66$). All were significant at the 99% confidence interval

Burn severity (%)	Crown scorch (%)	Crown consumption (%)	Bole scorch at 1 m (%)	Bole scorch (%)	Basal scorch (%)	Basal char (%)
Low	19.5 (3.3)	0.1 (0.1)	15.2 (2.1)	35.3 (7.6)	80.2 (4.1)	8.7 (3.6)
Moderate	69.7 (4.3)	4.9 (2.2)	41.9 (3.2)	87.1 (2.6)	79.3 (6.2)	19.2 (6.2)
High	8.8 (7.3)	90.6 (7.3)	99.7 (0.2)	99.9 (0.1)	48.2 (9.3)	51.8 (9.3)

4 **Table 2. Prediction results from immediate remote sensing and fractional cover measures**

5 All regressions, except those denoted by **, were significant at the 95% level. s.e. denotes the standard
 6 error of the estimate. Prediction statistics ($n = 66$) between immediate post-fire remote fractional
 7 measures (char fraction, green vegetation fraction) with 1-year post-fire field measures

Ground predictor (y)	Remote measures (x)					
	Fraction char cover			Fraction green cover		
	r^2	s.e.	Equation	r^2	s.e.	Equation
<i>Canopy variables</i>						
% Live tree	0.69	23.17	$-483 \times x + 487$	0.59	25.65	$254 \times x - 15$
Crown scorch	0.17	31.73	$-201 \times x + 224$	**		
Crown consumption	0.65	26.25	$499 \times x - 422$	0.42	33.83	$-227 \times x + 89$
Total crown fire effects	0.57	18.88	$298 \times x - 197$	0.55	19.16	$-168 \times x + 114$
<i>Subcanopy variables</i>						
Bole scorch	0.72	18.27	$411 \times x - 318$	0.60	22.12	$-212 \times x + 108$
Basal char	0.33	28.77	$277 \times x - 206$	0.32	28.82	$-157 \times x + 84$
Basal scorch	0.21	4.81	$35 \times x + 65$	0.16	4.97	$-17 \times x + 101$
Average bark thickness	0.48	0.28	$-3.6 \times x + 4.3$	0.35	0.32	$1.8 \times x + 0.51$
Bole scorch at 1 m	0.43	20.29	$243 \times x - 144$	0.44	20.02	$-141 \times x + 112$
Total BI 1 m tree	0.64	36.96	$289 \times x - 396$	0.56	41.12	$-365 \times x + 320$
Floor BI	0.44	49.37	$607 \times x - 339$	0.31	55.04	$-277 \times x + 284$
Litter depth	0.49	0.24	$-3.4 \times x + 3.5$	0.39	0.27	$1.7 \times x + 0.04$
Litter organic weight	0.71	3.99	$-80 \times x + 82$	0.64	4.40	$44 \times x - 1.70$

8 **Table 3. Prediction statistics ($n = 66$) between immediate post-fire brown (senesced vegetation)
 9 fraction and immediate post-fire NBR with 1-year post-fire field measures**

10 All regressions, except those denoted by ** were significant at the 95% level. s.e. denotes the standard
 11 error of the estimate

Ground predictor (y)	Remote measures (x)					
	Fraction brown cover			Immediate post-fire NBR		
	r^2	s.e.	Equation	r^2	s.e.	Equation
<i>Canopy variables</i>						
% Live tree	0.24	36.37	$-270 \times x + 0.6$	0.54	28.44	$0.11 \times x + 52.69$
Crown scorch	**			0.07	33.71	$0.03 \times x + 41.95$
Crown consumption	0.1	42.22	$13 \times x + 39$	0.43	33.67	$-0.10 \times x + 27.79$
Total crown fire effects	0.28	24.36	$196 \times x + 107$	0.50	20.25	$-0.71 \times x + 69.74$
<i>Subcanopy variables</i>						
Bole scorch	0.23	30.56	$218 \times x + 94$	0.53	23.81	$-0.09 \times x + 52.05$
Basal char	0.16	32.12	$186 \times x + 77$	0.31	29.11	$-0.07 \times x + 42.30$
Basal scorch	**			0.17	4.952	$-0.01 \times x + 96.82$

Average bark thickness	0.11	0.37	$-1.6 \times x + 0.7$	0.40	0.31	$0.001 \times x + 0.99$
Bole scorch at 1 m	0.24	23.43	$171 \times x + 106$	0.31	22.27	$-0.53 \times x + 75.53$
Total BI 1 m tree	0.23	54.16	$392 \times x + 299$	0.54	41.89	$-0.16 \times x + 223.21$
Floor BI	0.09	63.25	$250 \times x + 259$	0.34	53.72	$-0.14 \times x + 206.05$
Litter depth	0.14	0.32	$-1.7 \times x + 0.2$	0.39	0.27	$0.001 \times x + 0.50$
Litter organic weight	0.28	6.23	$-48 \times x + 1$	0.57	4.83	$0.02 \times x + 10.06$

Table 4. Prediction statistics ($n = 66$) between immediate and 1-year post-fire dNBR with 1-year post-fire field measures

All models were significant at the 95% confidence level. s.e. denotes the standard error of the estimate

Ground predictor (y)	Remote measures (x)					
	Immediate post-fire dNBR			1-year post-fire dNBR		
	r^2	s.e.	Equation	r^2	s.e.	Equation
<i>Canopy variables</i>						
% Live tree	0.53	28.65	$-0.10 \times x + 105$	0.74	21.24	$-0.13 \times x + 97$
Crown scorch	0.07	33.58	$-0.03 \times x + 58$	0.16	31.98	$-0.05 \times x + 61$
Crown consumption	0.44	33.46	$0.10 \times x - 23$	0.62	27.42	$0.12 \times x - 16$
Total crown fire effects	0.49	20.51	$-0.07 \times x + 35$	0.55	19.27	$0.07 \times x + 44$
<i>Subcanopy variables</i>						
Bole scorch	0.50	24.50	$0.08 \times x + 10$	0.68	19.58	$0.10 \times x + 17$
Basal char	0.28	29.77	$0.06 \times x + 11$	0.34	28.36	$0.07 \times x + 18$
Basal scorch	0.19	4.88	$0.01 \times x + 93$	0.22	4.78	$0.01 \times x + 94$
Average bark thickness	0.43	0.30	$-0.001 \times x + 1$	0.48	0.28	$-0.001 \times x + 1.3$
Bole scorch at 1 m	0.31	22.37	$0.05 \times x + 50$	0.43	20.17	$0.06 - x + 56$
Total BI 1 m tree	0.53	42.46	$0.15 \times x + 146$	0.63	37.48	$0.17 \times x + 164$
Floor BI	0.40	51.29	$0.14 \times x + 132$	0.50	46.73	$0.16 \times x + 148$
Litter depth	0.42	0.26	$-0.001 \times x + 1$	0.36	0.28	$-0.001 \times x + 0.74$
Litter organic weight	0.59	4.71	$-0.02 \times x + 19$	0.63	4.47	$-0.02 \times x + 16$

Table 5. Prediction statistics ($n = 66$) between immediate post-fire remote fractional measures (char fraction, green vegetation fraction) with 1-year post-fire field measures

All regressions, except those denoted by **, were significant at the 95% level. s.e. denotes the standard error of the estimate

Ground predictor (y)	Fraction char cover and fraction green cover			
	r^2	s.e.	Char fraction Significance	Green fraction Significance
<i>Canopy variables</i>				
% Live tree	0.71	22.75	0.00	**
Crown scorch	0.23	30.82	0.00	0.03
Crown consumption	0.66	26.32	0.00	**
Total crown fire effects	0.61	18.13	0.01	0.02
<i>Subcanopy variables</i>				
Bole scorch	0.74	18.09	0.00	**
Basal charring	0.35	28.44	**	**
Basal scorch	0.22	4.85	0.049	**
Average bark thickness	0.48	0.29	0.001	**
Bole scorch at 1 m	0.47	19.65	**	0.03
Total BI 1 m tree	0.66	36.36	0.00	**
Floor BI	0.44	49.78	0.00	**
Litter depth	0.50	0.25	0.001	**
Litter organic weight	0.73	3.83	0.00	0.03

Table 6. Pearson correlation coefficient (r) and Signal to Noise Ratio (SNR) analysis for bands 4 and 7

	Variable 1	Variable 2	r	Significance
Prefire and immediate post-fire	Band 4 difference	Band 7 difference	-0.52	0.00
Prefire and 1 year post-fire	Band 4 difference	Band 7 difference	-0.33	0.01
SNR calculations using pre- and immediate post-fire imagery		SNR _{band 4(Immediate)} = 3.2 SNR _{band 7(Immediate)} = 0.75		
SNR calculations using pre and 1-year post-fire imagery		SNR _{band 4(1-year)} = 1.4 SNR _{band 7(1-year)} = 0.47		

Table 7. Relation between % cover measures of burn severity and carbon (C) and water (H₂O) cycles

ET denotes evapotranspiration

Ecological metrics	Fire-effects reference(s)	Linkages to C and H ₂ O cycles
Tree survival/mortality	Miller and Yool (2002) Litton <i>et al.</i> (2003); Trumbore (2006)	C accumulation/ET rates
Bare soil	Goforth <i>et al.</i> (2005)	Plant establishment/soil respiration rates
Reddened soil	Doerr and Cerda (2005)	Infiltration, water repellency, and erosion
Exposed litter	Lewis <i>et al.</i> (2006) Crockford and Richardson (2000)	Plant establishment/water repellency Surface evaporation
White ash	Smith <i>et al.</i> (2005)	C volatilization/water repellency
Coarse woody debris	Smith and Hudak (2005)	C volatilization/erosion

Fig. 1. The location of the Jasper Fire, South Dakota (USA). The image insert is the fractional char cover image produced using the immediate post-fire Landsat image.

Fig. 2. Generic spectral reflectance curves of green vegetation, senesced vegetation, and char (black ash). Spectra were acquired by Smith *et al.* (2005). The data gap ~1.8 μm represents the dominant water absorption feature where data quality is insufficient for analysis. The two shaded columns highlight the general wavelength ranges of Landsat band 4 (0.76–0.90 μm) and band 7 (2.08–2.35 μm) for both TM and ETM.

Fig. 3. Left image shows a 1-year post-fire view of a ponderosa pine forest burned by non-stand replacing fire. Right image highlights a typical scorched bole, as measured in the field. A color version of this figure is available from the journal website.

Fig. 4. Correlation of fractional cover measures where dark lines depict the main linear trend. Scatter plot of fractional green and senesced vegetation cover ($r = -0.86$) and scatter plot of fractional char cover with green vegetation ($r = -0.85$).