

PETERSEN PLANE ARRANGEMENTS AND A SURFACE WITH MULTIPLE 7-SECANTS

HIROTACHI ABO, HOLGER P. KLEY, AND CHRIS PETERSON

ABSTRACT. We study configurations of 2-planes in \mathbb{P}^4 that are combinatorially described by the Petersen graph. We discuss conditions for configurations to be locally Cohen-Macaulay and describe the Hilbert scheme of such arrangements. An analysis of the homogeneous ideals of these configurations leads, via linkage, to a class of smooth, general type surfaces in \mathbb{P}^4 . We compute their numerical invariants and show that they have the unusual property that they admit (multiple) 7-secants. Finally, we demonstrate that the construction applied to Petersen arrangements with additional symmetry leads to surfaces with exceptional automorphism groups.

1. INTRODUCTION

Linkage (or liaison) theory can provide a bridge between combinatorially interesting varieties such as ℓ -plane arrangements (equidimensional unions of projective subspaces of \mathbb{P}^n) and geometrically interesting ones: smooth projective varieties. This paper studies one such bridge, connecting the class of 2-plane arrangements in \mathbb{P}^4 whose incidence structure may be described by the Petersen graph with a previously unpublished class of smooth surfaces.

We begin, in Section 2, with a review of the relevant combinatorics of the Petersen graph. Next, in Section 3, we associate a graph to an arrangement: the vertices are the linear spaces in the arrangement and two vertices are joined by an edge if the spaces meet in greater than the expected dimension. Our study of the geometry of 2-plane arrangements begins with a necessary combinatorial condition (Corollary 3.10) on the associated graph to insure that the underlying arrangement is locally Cohen-Macaulay (ICM). Since linkage preserves the ICM property, such a result is essential if we are interested in linking to smooth surfaces.

The discussion leads naturally to the consideration of 2-plane arrangements whose associated graph is the Petersen graph, which begs the following questions: do such arrangements exist, and are they (generally) locally Cohen-Macaulay? In Section 4, we answer both questions in the affirmative (Propositions 4.9) via an explicit and combinatorially interesting construction and show (Proposition 4.10) that in fact, all ICM Petersen arrangements arise in this manner. The modular nature of the construction has immediate benefits. For example, we are able to give (Theorem 4.11) detailed information about the Hilbert scheme of such arrangements.

2000 *Mathematics Subject Classification.* 14N20, 14J50, 14J25, 14M06, 05E20, 14J10.

Key words and phrases. Plane Arrangements, Petersen Graph, Automorphisms, 7-Secants, General Type Surface, Liaison, Group Actions, Linkage.

The second and third authors were partially supported by NSF grant MSPA-MCS-0434351.

Furthermore, the explicit construction allows a detailed description of the homogeneous ideal of an ICM Petersen arrangement; this is the subject of Section 5. In particular, we find (Theorem 5.7) that the ideal always contains six quintic forms. Thus, it is not surprising that Petersen arrangements may be linked, via a general pair of quintics, to degree fifteen surfaces in \mathbb{P}^4 . It should be noted that Rane-stad and Aure (independently and unpublished) explored such a construction while considering the classification of low-degree surfaces in \mathbb{P}^4 .

These linked, degree fifteen surfaces are the subject of Section 6. In particular, we show that they are smooth and of general type (Theorem 6.2). By exploiting the link between combinatorics and geometry once more, we show (Theorem 6.6) that the surfaces so obtained have the surprising property that they admit a one-dimensional family of 6-secant lines and fifteen 7-secant lines. Finally, in Theorem 7.1, we show how to generate examples of these surfaces with non-trivial automorphism groups.

Please note that all computer algebra scripts used in examples and computations throughout the present article may be found at [1].

Notation. For simplicity, fix an algebraically closed field K of characteristic zero. (In fact, the reader will note that all results will hold over sufficiently large algebraic extensions of the prime field in all but finitely many positive characteristics as well.)

For n a non-negative integer, let $\mathbb{P}^n := \mathbb{P}(K^{n+1})$ denote the projective space of dimension n over K . By a *linear (sub)space* (of \mathbb{P}^n) we mean $\mathbb{P}(V)$ for some vector subspace $V \subset K^{n+1}$. If $\dim V = \ell + 1$, we also refer to $\mathbb{P}(V)$ as an ℓ -*plane* in \mathbb{P}^n . Finally, for linear subspaces $\Lambda_1 = \mathbb{P}(V)$ and $\Lambda_2 = \mathbb{P}(W)$ of \mathbb{P}^n , let $\overline{\Lambda_1, \Lambda_2} = \mathbb{P}(V + W)$ denote the linear space they span.

Acknowledgments. The authors would like to thank Rick Miranda, Kristian Rane-stad, Bernd Sturmfels and Ravi Vakil for stimulating conversations and questions.

2. THE PETERSEN GRAPH

In this section, we recall a few definitions from graph theory. We then define the Petersen graph and outline some of its standard properties.

Definition 2.1. Let Γ be a graph with vertex set V and edge set E .

- A *matching* in Γ is a subset $E' \subset E$ such that no two edges in E' share a vertex.
- A *perfect matching* in Γ is a matching that covers every vertex in Γ .
- An *independent set of vertices* of Γ is a subset $S \subset V$ such that no pair of vertices in S form an edge of Γ .
- A *maximal independent set of vertices* is an independent set of vertices which is not properly contained in any other independent set of vertices.
- The *girth* of Γ is the length of the shortest, non-trivial cycle in Γ . If Γ contains no non-trivial cycles, it is said to have *infinite girth*.

By a slight abuse of notation, we write $\{0, 1, 2, 3, 4\} = \mathbb{Z}/(5)$ for (the elements of) the integers modulo 5.

The *Petersen graph* is the graph Γ_{Pete} with vertex set

$$V(\Gamma_{\text{Pete}}) := \{0, 1, 2, 3, 4, 0', 1', 2', 3', 4'\} = \{i, i' : i \in \mathbb{Z}/(5)\}$$

and edge set

$$\begin{aligned} E(\Gamma_{\text{Pete}}) &:= \{01, 12, 23, 34, 40, 00', 11', 22', 33', 44', 0'2', 1'3', 2'4', 3'0', 4'1'\} \\ &= \{i(i+1), ii', i'(i+2)' : i \in \mathbb{Z}/(5)\}. \end{aligned}$$

One standard way to realize Γ_{Pete} is shown in Figure 1. Notice that Γ_{Pete} has diameter 2 and girth 5.




FIGURE 1. The Petersen graph

Henceforth, let $\rho \in \text{Aut}(\Gamma_{\text{Pete}})$ be the “rotation by $2\pi/5$,” i.e., the automorphism whose action on $V(\Gamma_{\text{Pete}})$ is given by

$$\rho(i) = i + 1 \text{ and } \rho(i') = (i + 1)' \text{ for } i \in \mathbb{Z}/(5),$$

let $\tau \in \text{Aut}(\Gamma_{\text{Pete}})$ be the automorphism whose action on $V(\Gamma_{\text{Pete}})$ is given (in cycle notation) by

$$\tau := (00')(13'42')(21'34')$$

and let $\omega \in \text{Aut}(\Gamma_{\text{Pete}})$ be the automorphism whose action on $V(\Gamma_{\text{Pete}})$ is given (in cycle notation) by

$$\omega := (21')(34')(2'3').$$

Referring to Figure 1, τ more or less “turns Γ_{Pete} inside-out.” Also, observe that

$$\sigma := \tau^2 = (14)(23)(1'4')(2'3') \in \text{Aut}(\Gamma_{\text{Pete}})$$

is the “reflection across the vertical.”

We identify the maximal independent vertex set

$$V_0 := \{1, 4, 2', 3'\} \subset V(\Gamma_{\text{Pete}})$$

of Γ_{Pete} and observe that any 4-element maximal independent vertex set of Γ_{Pete} is one of

$$V_j := \rho^j V_0$$

for $j = 0, \dots, 4$. Notice that each vertex of Γ_{Pete} lies in precisely two of the V_j and that any two of the V_j intersect in exactly one vertex of Γ_{Pete} . Consequently, $\text{Aut}(\Gamma_{\text{Pete}})$ acts faithfully on $\{V_0, \dots, V_4\}$. Furthermore, ρ acts as a 5-cycle on $\{V_0, \dots, V_4\}$ while ω acts as the transposition $(V_1 V_4)$ on $\{V_0, \dots, V_4\}$, so we recover the well-known

Proposition 2.2. *The automorphism group of the Petersen graph is isomorphic to the symmetric group on 5 letters.*

Similarly, we identify the perfect matching

$$(2.1) \quad E_5 := \{00', 11', 22', 33', 44'\} \subset E(\Gamma_{\text{Pete}}),$$

and observe that every perfect matching of Γ_{Pete} is either E_5 or one of

$$(2.2) \quad E_j := \rho^j \omega E_5 \quad \text{for } j = 0, \dots, 4.$$

Clearly, $\text{Aut}(\Gamma_{\text{Pete}})$ acts transitively on $\{E_0, \dots, E_5\}$, so the group

$$G := \text{Stab}(E_5) \leq \text{Aut}(\Gamma_{\text{Pete}})$$

has order 20. One easily checks that $\rho, \tau \in G$, and concludes that

$$(2.3) \quad G = \langle \rho, \tau \rangle \leq \text{Aut}(\Gamma_{\text{Pete}}).$$

Finally, we identify

$$(2.4) \quad D := \langle \rho, \sigma \rangle = \langle \rho, \tau^2 \rangle \leq G,$$

which is isomorphic to the dihedral group of order 10.

3. ARRANGEMENTS AND THEIR INCIDENCE GRAPHS

In this section we define our basic objects of study and prove a simple but fundamental proposition about their algebro-geometric structure.

We begin with a trivial but useful observation about linear spaces:

Remark 3.1. Let P, Q and $R \subset \mathbb{P}^n$ be linear spaces. If

$$\dim(P \cap R) + \dim(Q \cap R) - \dim R \geq \dim(P \cap Q),$$

then $(P \cap Q) \subset R$. In particular, if R is a 2-plane meeting P and Q in lines and if $P \cap Q = \{p\}$, then $p \in R$.

Definition 3.2. For positive integers $\ell < n$, an ℓ -plane arrangement in \mathbb{P}^n is a finite collection of ℓ -dimensional linear subspaces of \mathbb{P}^n .

Notation 3.3. Let \mathcal{A} be an ℓ -plane arrangement in \mathbb{P}^n , let $S = K[x_0, \dots, x_n]$ be the homogeneous coordinate ring of \mathbb{P}^n , let $H \subset \mathbb{P}^n$ be a hyperplane, and let $p \in \mathbb{P}^n$. Then we write

- $V_{\mathcal{A}} := \bigcup_{\Lambda \in \mathcal{A}} \Lambda$ for the corresponding projective variety in \mathbb{P}^n ,
- $I_{\mathcal{A}} \subset S$ for the homogeneous ideal of $V_{\mathcal{A}}$ and $\mathcal{J}_{\mathcal{A}}$ for its ideal sheaf in $\mathcal{O}_{\mathbb{P}^n}$,
- $\mathcal{A}_p := \{\Lambda \in \mathcal{A} : p \in \Lambda\}$ for the subarrangement of ℓ -planes in \mathcal{A} which pass through p ,
- $H \cap \mathcal{A} := \{H \cap \Lambda : \Lambda \in \mathcal{A}\}$ for the *hyperplane section* of \mathcal{A} by H , and
- $\overline{p, \mathcal{A}} := \{\overline{p, \Lambda} : \Lambda \in \mathcal{A}\}$ for the *cone over \mathcal{A} with vertex p* .

Remark 3.4. We freely use three simple facts about cones and hyperplane sections of ℓ -plane arrangements:

- If \mathcal{A} is an ℓ -plane arrangement in \mathbb{P}^n and if $H \subset \mathbb{P}^n$ is a hyperplane not containing any of the members of \mathcal{A} , then $H \cap \mathcal{A}$ is an $(\ell - 1)$ -plane arrangement in H .
- If \mathcal{A} is an $(\ell - 1)$ -plane arrangement in a hyperplane $H \subset \mathbb{P}^n$ and if $p \notin H$, then $\overline{p, \mathcal{A}}$ is an ℓ -plane arrangement in \mathbb{P}^n .
- For $p \in \mathbb{P}^n$ and $H \subset \mathbb{P}^n$ a general hyperplane, $\mathcal{A}_p = \overline{p, H \cap \mathcal{A}}$.

In general, we will say that an arrangement has a certain geometric property if its corresponding variety does. In particular, we have:

Definition 3.5. Let \mathcal{A} be an ℓ -plane arrangement in \mathbb{P}^n and let $p \in V_{\mathcal{A}}$ be a point corresponding to the prime ideal $\mathfrak{p} \subset S$. We say that \mathcal{A} is *arithmetically Cohen-Macaulay (aCM)* if the homogeneous coordinate ring $S/I_{\mathcal{A}}$ is a Cohen-Macaulay ring. We say that \mathcal{A} is *locally Cohen-Macaulay (lCM) at p* if the local ring $(S/I_{\mathcal{A}})_{\mathfrak{p}}$ is Cohen-Macaulay. Finally, we say that \mathcal{A} is *locally Cohen-Macaulay* if it is lCM at all closed points $p \in V_{\mathcal{A}}$.

The following proposition is useful in understanding when an arrangement is lCM at a point.

Proposition 3.6. *Let \mathcal{A} be an ℓ -plane arrangement in \mathbb{P}^n with $\ell \geq 2$. Let $p \in V_{\mathcal{A}}$. Let H be a hyperplane not containing any plane of \mathcal{A} . The following are equivalent:*

- (i) \mathcal{A} is lCM at p .
- (ii) \mathcal{A}_p is lCM at p .
- (iii) \mathcal{A}_p is aCM.
- (iv) $H \cap \mathcal{A}_p$ is aCM.

Proof. The equivalence of (i) and (ii) is clear. The equivalence of (iii) and (iv) is an immediate corollary of a result of Huneke and Ulrich [6]. Finally, the equivalence of (ii) and (iv) follows because \mathcal{A}_p is a cone over $H \cap \mathcal{A}_p$, and a variety is aCM if and only if the cone over the variety is lCM at the vertex of the cone. \square

Example 3.7. Consider distinct 2-planes $P_1, P_2, P_3, P_4 \subset \mathbb{P}^4$ not all contained in a hyperplane. Suppose that $P_1 \cap P_3 = P_2 \cap P_4 = P_1 \cap P_4 = \{p\}$ consists of a single point while $P_1 \cap P_2, P_2 \cap P_3$, and $P_3 \cap P_4$ are lines (necessarily passing through p by Remark 3.1). Then the arrangement $\{P_1, P_2, P_3, P_4\}$ is a cone over a projection into \mathbb{P}^3 of a degenerate rational normal curve of degree 4. Such curves are not aCM and hence the arrangement $\{P_1, P_2, P_3, P_4\}$ is not lCM.

It is useful to keep track of those pairs of planes in an ℓ -plane arrangement that are in special position with respect to each other. We represent this data via a graph.

Definition 3.8. If \mathcal{A} is an ℓ -plane arrangement in \mathbb{P}^n , the *incidence graph* of \mathcal{A} is the graph $\Gamma(\mathcal{A})$, with vertices the planes of \mathcal{A} and edges the pairs of planes with special intersection. In other words, $\Gamma(\mathcal{A})$ has vertex set \mathcal{A} , and edge set $\{\Lambda\Lambda' : \dim \Lambda \cap \Lambda' > n - 2\ell\}$.

Remark 3.9. Because it captures non-generic behavior only between *pairs* of ℓ -planes, the graph $\Gamma(\mathcal{A})$ may not completely characterize the geometry of the arrangement \mathcal{A} . For example, in \mathbb{P}^4 , both the arrangement of three 2-planes all containing a common line and the arrangement of three 2-planes meeting pairwise in three distinct lines have the complete graph on three vertices as their incidence

graphs. (These arrangements could, however, be distinguished by an *incidence complex*.)

Corollary 3.10. *Let \mathcal{A} be a 2-plane arrangement in \mathbb{P}^4 and let $p \in V_{\mathcal{A}}$. If $\Gamma(\mathcal{A}_p)$ is disconnected, then \mathcal{A} is not ICM at p . As a partial converse, if $\Gamma(\mathcal{A}_p)$ is connected and of order at most three, then \mathcal{A} is ICM at p .*

Proof. Let H be a hyperplane not passing through p . Then \mathcal{A}_p is a cone over the arrangement $\mathcal{L} = \mathcal{A}_p \cap H$ of lines in $H = \mathbb{P}^3$, and by Proposition 3.6, \mathcal{A} is ICM at p if and only if \mathcal{L} is aCM. Now $\Gamma(\mathcal{A}_p)$ is connected if and only if $V_{\mathcal{L}}$ is connected, and disconnected curves are not aCM. Conversely, suppose $V_{\mathcal{L}}$ is connected and has at most three components. Then $V_{\mathcal{L}}$ is either a complete intersection, a cone over three non-collinear points in a plane, or a degenerate twisted cubic, all of which are aCM. \square

Remark 3.11. Let \mathcal{A} be a 2-plane arrangement in \mathbb{P}^4 . Corollary 3.10 suggests that we should *expect* \mathcal{A} to be ICM if $\Gamma(\mathcal{A})$ is connected and of diameter at most 2. On the other hand, Example 3.7 suggests that if the diameter of $\Gamma(\mathcal{A})$ exceeds 2, then \mathcal{A} may well fail to be ICM. If $\Gamma(\mathcal{A})$ is disconnected, \mathcal{A} fails to be ICM in all cases.

4. PETERSEN ARRANGEMENTS

For the remainder of this paper, a *plane* means a 2-plane. In Remark 3.11, we observed that a plane arrangement \mathcal{A} in \mathbb{P}^4 , constructed in such a way that $\Gamma(\mathcal{A})$ has diameter 2, is expected to be ICM. Among the graphs of finite girth, only those of girth ≤ 5 may have diameter ≤ 2 . Thus, graphs of girth 5 and diameter 2 are a natural place to look for interesting examples of arrangements.

The simplest such graph is the pentagon, and one can readily construct a plane arrangement \mathcal{A} in \mathbb{P}^4 such that $\Gamma(\mathcal{A})$ is isomorphic to the pentagon. This plane arrangement appears as a degenerate case in the family containing the elliptic quintic scrolls. Here elliptic quintic scrolls are ruled surfaces over elliptic curves embedded in \mathbb{P}^4 in such a way that all the fibers have degree 1. These surfaces determine one of the few known families of smooth irregular surfaces in \mathbb{P}^4 . The (degenerate) elliptic quintic scroll $V_{\mathcal{A}}$ is cut out by five cubic hypersurfaces and can be linked in the complete intersection of two such cubic hypersurfaces to the Veronese surface (which is known as the only smooth non-degenerate surface in \mathbb{P}^4 that is not linearly normal).

As we observed in Section 2, the Petersen graph Γ_{Pete} likewise has diameter 2 and girth 5.

Definition 4.1. A (two-)plane arrangement \mathcal{A} in \mathbb{P}^4 is a *Petersen arrangement* if $\Gamma(\mathcal{A}) \simeq \Gamma_{\text{Pete}}$. A *labeling* of a Petersen arrangement \mathcal{A} is an isomorphism $\psi: \Gamma_{\text{Pete}} \xrightarrow{\sim} \Gamma(\mathcal{A})$.

Several natural questions are: Do Petersen arrangements exist? If so, how big is the family of such arrangements? Do ICM Petersen arrangements exist? If so, can liaison theory be used to produce (smooth) surfaces from Petersen arrangements? Do Petersen arrangements with extra structure (such as a prescribed automorphism group) exist? The remainder of this article addresses these questions.

4.1. Construction of Petersen Arrangements. Let $\mathcal{L} = \{L_i\}_{i \in \mathbb{Z}/(5)}$ be an indexed collection of lines in \mathbb{P}^4 . Suppose that \mathcal{L} satisfies

Condition 1. L_0, \dots, L_4 are pairwise skew.

Then for each pair $\{i, j\}$ of distinct elements of $\mathbb{Z}/(5)$, L_i and L_j span a hyperplane

$$H_{i,j} := \overline{L_i, L_j}.$$

Clearly, $H_{i,j} = H_{j,i}$.

Suppose that \mathcal{L} also satisfies

Condition 2. No three lines of \mathcal{L} are contained in a hyperplane.

Lemma 4.2. *The following are equivalent conditions on a collection \mathcal{L} satisfying Condition 1:*

- (i) *Condition 2.*
- (ii) *For all choices of distinct $i, j, k \in \mathbb{Z}/(5)$, the hyperplanes $H_{i,j}$ and $H_{i,k}$ are distinct.*
- (iii) *For all choices of distinct $i, j, k \in \mathbb{Z}/(5)$, there is a unique “trisecant” line $M_{i,j,k}$ meeting L_i, L_j and L_k .*
- (iv) *For all choices of three distinct $i, j, k \in \mathbb{Z}/(5)$, there is a unique plane containing L_i which meets L_j and L_k .*

Proof. This is immediate. The key observations are that

$$(4.1) \quad M_{i,j,k} = H_{i,j} \cap H_{j,k} \cap H_{k,i},$$

and that

$$(4.2) \quad H_{i,j} \cap H_{i,k} = \overline{L_i, M_{i,j,k}}$$

is the unique plane containing L_i and meeting L_j and L_k . □

For each $i \in \mathbb{Z}/(5)$, we define two planes

$$(4.3) \quad P_i := H_{i,i+1} \cap H_{i,i-1} \quad \text{and} \quad P'_i := H_{i,i+2} \cap H_{i,i-2}$$

and set

$$(4.4) \quad \mathcal{A} = \mathcal{A}(\mathcal{L}) := \{P_i, P'_i : i \in \mathbb{Z}/(5)\}.$$

We would like to show that \mathcal{A} is a Petersen arrangement. There are, however, two potential problems: $\Gamma(\mathcal{A})$ could contain Γ_{Pete} as a proper subgraph, and some of the planes in \mathcal{A} could coincide. To eliminate these possibilities, we require that \mathcal{L} additionally satisfy

Condition 3. No plane containing a line of \mathcal{L} meets three other lines of \mathcal{L} .

Lemma 4.3. *The following are equivalent conditions on a collection \mathcal{L} satisfying Conditions 1 and 2:*

- (i) *Condition 3.*
- (ii) $L_\ell \cap \overline{L_i, M_{i,j,k}} = \emptyset$ for all choices of distinct $i, j, k, \ell \in \mathbb{Z}/(5)$.
- (iii) $H_{i,j} \cap H_{i,k} \cap H_{i,\ell} = L_i$ for all choices of distinct $i, j, k, \ell \in \mathbb{Z}/(5)$.
- (iv) $M_{i,j,k} \cap M_{i,j,\ell} = \emptyset$ for all choices of distinct $i, j, k, \ell \in \mathbb{Z}/(5)$.

Proof. This is immediate in light of Lemma 4.2. □

We can now establish

Proposition 4.4. *Let $\mathcal{L} = \{L_i\}_{i \in \mathbb{Z}/(5)}$ be an indexed collection of lines in \mathbb{P}^4 satisfying Conditions 1–3. Then $\mathcal{A}(\mathcal{L})$ is a Petersen arrangement with labeling $\psi_{\mathcal{L}}: \Gamma_{\text{Pete}} \rightarrow \Gamma(\mathcal{A})$ given by $\psi_{\mathcal{L}}(i) = P_i$ and $\psi_{\mathcal{L}}(i') = P'_i$. Moreover, $P_i \cap P'_i = L_i$.*

Proof. Since $\psi_{\mathcal{L}}$ is a bijection on vertices, it suffices to show that it induces a bijections of edges, which we establish as a series of claims. Let $i \in \mathbb{Z}/(5)$.

Claim. $\dim(P_i \cap P_{i+1}) = 1$.

Proof. By construction,

$$P_i \cap P_{i+1} = H_{i-1,i} \cap H_{i,i+1} \cap H_{i+1,i+2}$$

which is at least one-dimensional. On the other hand, P_i contains L_i and P_{i+1} contains L_{i+1} and since these lines are skew by Condition 1, $P_i \neq P_{i+1}$.

Claim. $\dim(P'_i \cap P'_{i+2}) = 1$.

Proof. This is essentially identical to the previous case.

Claim. $\dim(P_i \cap P_{i+2}) = 0$.

Proof. Combining (4.2) with (ii) of Lemma 4.3 shows that $P_i = \overline{L_i, M_{i-1,i,i+1}}$ does not meet $L_{i+2} \subset P_{i+2}$. It follows that P_i and P_{i+2} cannot meet in a line.

Claim. $\dim(P'_i \cap P'_{i+1}) = 0$.

Proof. This is essentially identical to the previous case..

Claim. $P_i \cap P'_i = L_i$.

Proof. Using (iii) of Lemma 4.3,

$$\begin{aligned} P_i \cap P'_i &= H_{i-1,i} \cap H_{i,i+1} \cap H_{i-2,i} \cap H_{i,i+2} \\ &= L_i \cap H_{i,i+2} = L_i. \end{aligned}$$

Claim. $\dim(P_i \cap P'_j) = 0$ for all $j \neq i$.

Proof. We consider two cases: $P_i \cap P'_{i+1}$ and $P_i \cap P'_{i+2}$. The other two cases are essentially identical. Keeping in mind that indices are calculated modulo five, we have

$$P_i \cap P'_{i+1} = H_{i-1,i} \cap H_{i,i+1} \cap H_{i-1,i+1} \cap H_{i+1,i-2}.$$

Notice that three terms involve the index $i+1$. Applying (iii) of Lemma 4.3, we reduce to

$$P_i \cap P'_{i+1} = H_{i-1,i} \cap L_{i+1}$$

which must be a point by Condition 2.

For the second case, an analogous argument shows that

$$\begin{aligned} P_i \cap P'_{i+2} &= H_{i-1,i} \cap H_{i,i+1} \cap H_{i,i+2} \cap H_{i+2,i-1} \\ &= H_{i+2,i} \cap L_i \end{aligned}$$

is also a point. □

Remark 4.5. The symmetric group $\mathfrak{S} = \mathfrak{S}_{\mathbb{Z}/(5)}$ acts naturally on the indexing of a collection $\mathcal{L} = \{L_i\}_{i \in \mathbb{Z}/(5)}$ of lines in \mathbb{P}^4 satisfying Conditions 1–3. Let $\mathfrak{F} \leq \mathfrak{S}$ be the subgroup of order 20 generated by the 5-cycle (01234) and the 4-cycle (1342). Recall the group $G \leq \text{Aut}(\Gamma_{\text{Pete}})$ of (2.3) and observe that the mapping (01234) $\mapsto \rho$ and (1342) $\mapsto \tau$ extends to a group isomorphism $f: \mathfrak{F} \xrightarrow{\sim} G$. Now for $s \in \mathfrak{S}$, we have $\mathcal{A}(s\mathcal{L}) = \mathcal{A}(\mathcal{L})$ if and only if $s \in \mathfrak{F}$, in which case the corresponding labelings are related by $\psi_{s\mathcal{L}} = \psi_{\mathcal{L}}(f(s))$.

Of course, labeled Petersen arrangements lead to line arrangements via a reverse construction. Specifically, let \mathcal{A} be a Petersen arrangement and $\psi: \Gamma_{\text{Pete}} \xrightarrow{\sim} \Gamma(\mathcal{A})$ a labeling. Then we have the indexed collection of lines

$$\mathcal{L}(\mathcal{A}, \psi) = \{\psi(i) \cap \psi(i')\}_{i \in \mathbb{Z}/(5)}.$$

Remark 4.6. The lines of $\mathcal{L}(\mathcal{A}, \psi)$ correspond to the edges in the perfect matching $\psi(E_5) = \{\psi(ii')\}_{i \in \mathbb{Z}/(5)}$ of $\Gamma(\mathcal{A})$. Likewise, for $\alpha \in \text{Aut}(\Gamma_{\text{Pete}})$, the lines of $\mathcal{L}(\mathcal{A}, \psi\alpha)$ correspond to the edges in the perfect matching $\psi\alpha(E_5)$ of $\Gamma(\mathcal{A})$. Keeping in mind the notation of Remark 4.5, $\mathcal{L}(\mathcal{A}, \psi\alpha) = \mathcal{L}(\mathcal{A}, \psi)$ as sets if and only if $\alpha \in G = \text{Stab}(E_5) \leq \text{Aut}(\Gamma_{\text{Pete}})$, in which case $\mathcal{L}(\mathcal{A}, \psi\alpha) = f^{-1}(\alpha)\mathcal{L}(\mathcal{A}, \psi)$ as indexed sets.

Unfortunately, not every Petersen arrangement can be obtained as in (4.4):

Example 4.7. Let \mathcal{A} be any Petersen arrangement and let $H \subset \mathbb{P}^4$ be a general hyperplane. Choose a point $p \notin H$. Then

$$\mathcal{B} := \overline{p, (\mathcal{A} \cap H)}$$

is a Petersen arrangement. Choose a labeling $\psi: \Gamma_{\text{Pete}} \xrightarrow{\sim} \Gamma(\mathcal{B})$. Then the lines of $\mathcal{L}(\mathcal{B}, \psi)$ all pass through the vertex p and thus, don't satisfy Condition 1. Since ψ was arbitrary, this shows that it is impossible to realize \mathcal{B} as $\mathcal{A}(\mathcal{L})$ for any indexed collection \mathcal{L} of lines.

4.2. LCM Petersen Arrangements. Let $\mathcal{L} = \{L_i\}_{i \in \mathbb{Z}/(5)}$ be an indexed collection of lines in \mathbb{P}^4 satisfying Conditions 1–3, and let $\mathcal{A} = \mathcal{A}(\mathcal{L})$ as in (4.4). While Proposition 4.4 shows that \mathcal{A} is a Petersen arrangement, \mathcal{A} needn't be locally Cohen-Macaulay. In light of Proposition 3.6, the study of LCM plane arrangements in \mathbb{P}^4 can be reduced to the study of aCM line arrangements in \mathbb{P}^3 . For our purposes, the following will suffice:

Lemma 4.8. *Let \mathcal{L} be an arrangement of lines in \mathbb{P}^3 . If $\Gamma(\mathcal{L})$ is isomorphic to either*

- (i) *the Petersen graph Γ_{Pete} ,*
- (ii) *the pentagon C_5 , or*
- (iii) *the complete bipartite graph $K_{1,3}$,*

then $V_{\mathcal{L}}$ is not arithmetically Cohen-Macaulay.

Proof. In case (i), $V_{\mathcal{L}}$ is a curve of degree 10 and arithmetic genus 6. In case (ii), $V_{\mathcal{L}}$ is a curve of degree 5 and arithmetic genus 1. Finally, in case (iii), $V_{\mathcal{L}}$ is a curve of degree 4 and arithmetic genus 0. In each case, Riemann-Roch now implies that $H^1(\mathcal{J}_{V_{\mathcal{L}}}(1)) \neq 0$. Consequently, $V_{\mathcal{L}}$ is not aCM. \square

For instance, the arrangement \mathcal{B} of Example 4.7, being a cone over a non-aCM line arrangement, fails to be LCM by Proposition 3.6.

As in the previous section, let $\mathcal{L} = \{L_i\}_{i \in \mathbb{Z}/(5)}$ be an indexed collection of lines in \mathbb{P}^4 satisfying Conditions 1–3, and let $\mathcal{A} = \mathcal{A}(\mathcal{L})$ be the resulting Petersen arrangement.

Let P be a plane in \mathcal{A} . Then there exist three planes in \mathcal{A} that are adjacent to P in $\Gamma(\mathcal{A})$. Conditions 1–3 do not imply that the lines along which these planes meet P form a triangle in P . Indeed, if these three lines all pass through a single $p \in P$, then \mathcal{A}_p fails to be LCM at p by case (iii) of Lemma 4.8. Keeping in mind the definition of the “trisecant” lines $M_{i,j,k}$ in Lemma 4.2, we therefore consider

Condition 4. For all $i \in \mathbb{Z}/(5)$, both $M_{i-2,i-1,i} \cap M_{i,i+1,i+2} = \emptyset$ and $M_{i-2,i,i+1} \cap M_{i-1,i,i+2} = \emptyset$.

Notice the similarity of this Condition 4 to Condition 3 via (iv) of Lemma 4.3.

Finally, in order to rule out non-ICM points arising as vertices of cones over the line arrangement of case (ii) of Lemma 4.8, we consider:

Condition 5. $\bigcap_{i \in \mathbb{Z}/(5)} H_{i,i+1} = \bigcap_{i \in \mathbb{Z}/(5)} H_{i,i+2} = \emptyset$.

Proposition 4.9. *Let \mathcal{L} be an indexed collection of lines in \mathbb{P}^4 satisfying Conditions 1–5. Then $\mathcal{A} = \mathcal{A}(\mathcal{L})$ is a locally Cohen-Macaulay Petersen arrangement.*

Proof. By Corollary 3.10, it suffices to show that $\Gamma(\mathcal{A}_p)$ is connected of order at most 3 for all $p \in V_{\mathcal{A}}$. All cases of this assertion are either obvious (i.e., when p lies in exactly one plane of \mathcal{A} or in exactly two planes of \mathcal{A} that meet in a line) or can be deduced from the

Claim. If $P, Q \in \mathcal{A}$ with $P \cap Q = \{p\}$, then there exists a unique $R \in \mathcal{A}$ meeting both P and Q in lines, in which case $\mathcal{A}_p = \{P, Q, R\}$.

Note that p cannot lie in three or more planes meeting pairwise in lines as the Petersen graph contains no 3-cycles. Thus, if \mathcal{A}_p consists of three or more planes, there must be at least two of them which meet in a point, in which case the claim applies.

Proof of claim. $P \cap Q = \{p\}$ means that P and Q are not adjacent in $\Gamma(\mathcal{A})$. Since the Petersen graph has no 4-cycles and any two of its vertices lie on a 5-cycle, there is a unique $R \in \mathcal{A}$ meeting both P and Q in lines. These two lines meet at a point of R which, by Remark 3.1 is necessarily p , so we are reduced to showing that no other $T \in \mathcal{A}$ contains p . Suppose to the contrary, that such a T exists. Then either

- (i) T is adjacent to either P or Q ,
- (ii) T is adjacent to none of P, Q and R , or
- (iii) T and R are adjacent.

In case (i), assume without loss of generality that P and T are adjacent. Then there exists a unique plane $U \in \mathcal{A}$ such that $C := \Gamma(\{T, P, R, Q, U\})$ is a 5-cycle in $\Gamma(\mathcal{A})$. By Remark 3.1, $p \in U$ as well, so

$$(4.5) \quad P \cap R \cap Q \cap U \cap T = \{p\}.$$

Now either $C = \Gamma(\{P_i : i \in \mathbb{Z}/(5)\})$ or $C = \Gamma(\{P'_i : i \in \mathbb{Z}/(5)\})$ or C contains two of the edges of E_5 . In the first two cases, (4.5) contradicts Condition 5, while in the third, it contradicts Condition 1.

In case (ii), P and Q are distance 2 from T , and there exist unique planes U and $U' \in \mathcal{A}$ adjacent to T such that P and Q are adjacent to U and U' respectively in $\Gamma(\mathcal{A})$. In other words, $C := \Gamma(\{T, U, P, R, Q, U'\})$ is a 6-cycle in $\Gamma(\mathcal{A})$. By repeated applications of Remark 3.1, all planes of C must pass through p . Since any 6-cycle in Γ_{Pete} must contain two of the edges of E_5 , this contradicts Condition 1 as in the previous case.

Finally, in case (iii), either $R = P_i$ or $R = P'_i$ for some $i \in \mathbb{Z}/(5)$. If $R = P_i$, then without loss of generality, $P = P_{i-1}$, $Q = P_{i+1}$ and $T = P'_i$. Then, using (4.3) and (4.1),

$$\{p\} = P \cap Q \cap R \cap T = M_{i-2,i-1,i} \cap M_{i,i+1,i+2},$$

contradicting Condition 4. If $R = P'_i$, an analogous argument again leads to a contradiction of Condition 4. \square

As it turns out, every locally Cohen-Macaulay Petersen arrangement arises as a result of our construction:

Proposition 4.10. *Let \mathcal{A} be a locally Cohen-Macaulay Petersen arrangement and $\psi: \Gamma_{\text{Pete}} \xrightarrow{\sim} \Gamma(\mathcal{A})$ a labeling. Then $\mathcal{L}(\mathcal{A}, \psi)$ satisfies Conditions 1–5, $\mathcal{A} = \mathcal{A}(\mathcal{L}(\mathcal{A}, \psi))$ and $\psi = \psi_{\mathcal{L}(\mathcal{A}, \psi)}$.*

Proof. Recall that we set $P_i = \psi(i)$, $P'_i = \psi(i')$ and $L_i = P_i \cap P'_i$ for all $i \in \mathbb{Z}/(5)$ and that $\mathcal{L} = \{L_i\}_{i \in \mathbb{Z}/(5)}$.

For any distinct $i, j \in \mathbb{Z}/(5)$, there is a unique $Q \in \mathcal{A}$ such that $\{P_i, P'_i, P_j, P'_j, Q\}$ are the vertices of a 5-cycle in $\Gamma(\mathcal{A})$. If $L_i \cap L_j \neq \emptyset$, then using Remark 3.1, $\{P_i, P'_i, P_j, P'_j, Q\}$ is a cone over a line arrangement with incidence graph isomorphic to the pentagon C_5 . If any other plane of \mathcal{A} also passes through the vertex p of this cone, then, by repeated application of Remark 3.1, all planes of \mathcal{A} have p in common. Thus, either \mathcal{A}_p is a cone over a line arrangement with pentagon incidence graph, or it is a cone over a line arrangement with Petersen incidence graph. In either case, \mathcal{A} is not ICM at p by Lemma 4.8. Thus, $L_i \cap L_j = \emptyset$, Condition 1 is satisfied and the hyperplanes $H_{i,j} = \overline{L_i, L_j}$ are defined.

Since P_i and P_{i+1} are adjacent in $\Gamma(\mathcal{A})$ for all i , they span a hyperplane. This hyperplane must contain L_i and L_{i+1} so that

$$(4.6) \quad H_{i,i+1} = \overline{P_i, P_{i+1}} \quad \text{for all } i \in \mathbb{Z}/(5).$$

Similarly,

$$(4.7) \quad H_{i,i+2} = \overline{P'_i, P'_{i+2}} \quad \text{for all } i \in \mathbb{Z}/(5).$$

Now suppose \mathcal{L} fails Condition 2. By Lemma 4.2, there exist distinct $i, j, k \in \mathbb{Z}/(5)$ such that L_i, L_j , and L_k lie in some hyperplane H . Without loss of generality, we may assume that either $i = j - 1$ and $k = j + 1$ or that $i = j - 2$ and $k = j + 2$. Using (4.6), the first case yields

$$\overline{P_{j-1}, P_j} = H_{j-1,j} = H = H_{j,j+1} = \overline{P_j, P_{j+1}}.$$

Thus, P_{j-1} and P_{j+1} must meet in a line, contradicting their non-adjacency in $\Gamma(\mathcal{A})$. In the second case, a parallel argument using (4.7) shows that P'_{j-2} meets P'_{j+2} in a line, contradicting their non-adjacency in $\Gamma(\mathcal{A})$. Thus, Condition 2 is satisfied and $P_i = H_{i-1,i} \cap H_{i,i+1}$ while $P'_i = H_{i-2,i} \cap H_{i,i+2}$ for all i .

Suppose Condition 3 fails. Then by Lemma 4.3, there exist distinct indices $i, j, k, \ell \in \mathbb{Z}/(5)$ such that $Q = H_{i,j} \cap H_{i,k} \cap H_{i,\ell}$ has dimension 2. Up to obvious reindexing, there are just two cases to consider: either $j = i - 1$, $k = i + 1$ and $\ell = i + 2$, or $j = i + 1$, $k = i + 2$ and $\ell = i + 3$. In the first of these, $P_i = Q \subset H_{i,i+2} \supset P'_{i+2}$, contradicting the non-adjacency of P_i and P'_{i+2} . In the second case, $Q = P'_i \subset H_{i,i+1} \supset P_{i+1}$, contradicting the non-adjacency of P_{i+1} and P'_i . Thus, Condition 3 is satisfied. Note that we have also established that $\mathcal{A} = \mathcal{A}(\mathcal{L}(\mathcal{A}, \psi))$ and $\psi = \psi_{\mathcal{L}(\mathcal{A}, \psi)}$.

Finally, if $\mathcal{L}(\mathcal{A}, \psi)$ fails Conditions 4 or 5, then $V_{\mathcal{A}}$ contains a point p such that \mathcal{A}_p is a cone over a line arrangement with incidence graph isomorphic to Γ_{Pete} , C_5 , or $K_{1,3}$, contradicting, via Lemma 4.8, that \mathcal{A} is ICM. \square

4.3. Examples. Conditions 1–5 are non-empty open conditions on the set of indexed collections \mathcal{L} of five lines in \mathbb{P}^4 , so ICM Petersen arrangements exist. To construct a specific example, choose random 2×5 matrices A_i of rank 2 to represent the lines $L_i \in \mathcal{L}$ and check by matrix manipulation and rank computations of matrices whether \mathcal{L} satisfies Conditions 1–5.

Alternatively, use a computer algebra system such as MACAULAY2. Represent the lines L_i via their ideals, each of which is generated by picking three random linear forms in five variables. In sufficiently large characteristic, Conditions 1–5 can be verified ideal-theoretically. Also, one can check that \mathcal{A} is locally Cohen-Macaulay directly by looking at the second syzygy module of $I_{\mathcal{A}}$; see [1].

Once such a sufficiently generic example has been constructed, one can check computationally that at the corresponding point, the Zariski tangent space to the Hilbert scheme of \mathbb{P}^4 has dimension 30. Again, see [1]. Now combining Remarks 4.5 and 4.6 and Propositions 4.9 and 4.10, we see that a parameter space for locally Cohen-Macaulay Petersen arrangements is $U/\text{Aut}(\Gamma_{\text{Pete}})$, where $U \subset \prod_{\mathbb{Z}/(5)} \mathbb{G}(1, 4)$ is the open set of indexed collections of lines satisfying Conditions 1–5 and $\text{Aut}(\Gamma_{\text{Pete}})$ acts freely, *though only the subgroup G acts by permutation of indices*.

Given that $\dim U = 30$, we have established:

Theorem 4.11. *Let \mathcal{A} be a general locally Cohen Macaulay Petersen arrangement. Then $V_{\mathcal{A}}$ lies on a unique component of the Hilbert scheme of \mathbb{P}^4 which is birational to $U/\text{Aut}(\Gamma_{\text{Pete}})$. Furthermore, the Hilbert scheme is smooth at $V_{\mathcal{A}}$ of dimension 30 and $V_{\mathcal{A}}$ cannot be smoothed in \mathbb{P}^4 .*

Example 4.12. Recall the dihedral subgroup $D = \langle \rho, \sigma \rangle$ of $\text{Aut}(\Gamma_{\text{Pete}})$ defined in (2.4). Over K , construct a 5-dimensional representation V of D with basis $\{e_i\}_{i \in \mathbb{Z}/(5)}$, by letting ρ act as the 5-cycle $(e_0 e_1 e_2 e_3 e_4)$ and σ as the product of transpositions $(e_1 e_4)(e_2 e_3)$. Then σ has eigenvalues 1 and -1 with corresponding eigenspaces $\langle e_0, e_1 + e_4, e_2 + e_3 \rangle$, and $\langle -e_1 + e_4, -e_2 + e_3 \rangle$, respectively. Consider the σ -invariant two-dimensional subspace W of V spanned by $v_1 = e_0 + e_2 + e_3$ and $v_2 = -e_1 - e_2 + e_3 + e_4$.

Now in the induced projective representation $\mathbb{P}(V)$, the line $L_0 := \mathbb{P}(W)$ is also σ -invariant. For $i \in \mathbb{Z}/(5)$, set $L_i := \rho^i(L_0)$. Then as a set, $\mathcal{L} = \{L_i\}_{i \in \mathbb{Z}/(5)}$ is invariant under the action of $D \hookrightarrow \text{PGl}(V)$, and it is straightforward to show that \mathcal{L} satisfies Conditions 1–5. Thus $\mathcal{A}(\mathcal{L})$ is a locally Cohen-Macaulay Petersen arrangement (see [1] for a direct verification), invariant under the action of D . Note, however, that the corresponding labelings change in the obvious way: $\psi_{\delta\mathcal{L}} = \psi_{\mathcal{L}} \circ \delta$ for $\delta \in D$.

5. GENERATORS OF $I_{\mathcal{A}}$

Let \mathcal{A} be a locally Cohen-Macaulay Petersen arrangement and fix a labeling $\psi: \Gamma_{\text{Pete}} \rightarrow \Gamma(A)$. As before, let $P_i = \psi(i)$ and $P'_i = \psi(i')$ for $i \in \mathbb{Z}/(5)$. The ideal $I_{\mathcal{A}}$ is obtained as the intersection of the ideals of the planes in \mathcal{A} and, by means of an example, one can determine that a minimal set of generators for $I_{\mathcal{A}}$ consists of six quintics and five sextics. The main goal of this section is to provide a combinatorial description of these generators. Among other things, this description will allow us to show that there are exactly five 6-secant lines to $V_{\mathcal{A}}$.

Let u and v be adjacent vertices of Γ_{Pete} and $e = uv$ the corresponding edge. Then the planes $\psi(u), \psi(v) \in \mathcal{A}$ span a hyperplane which we denote

$$H^{u,v} = H^e := \overline{\psi(u), \psi(v)}.$$

For each of the six perfect matchings E_i of Γ_{Pete} (see Section 2), we define a quintic hypersurface in \mathbb{P}^4 by

$$Q_i := \bigcup_{e \in E_i} H^e.$$

Let $q_i \in S$ be a quintic polynomial that defines Q_i . We will prove that $q_i \in (I_{\mathcal{A}})_5$. For each $i \in \mathbb{Z}/(5)$, define

$$(5.1) \quad l_i := H^{i,i'} \cap H^{(i+1)',(i-1)'} \cap H^{(i+2),(i-2)}.$$

Lemma 5.1. *The l_i satisfy:*

- (i) l_i is a line.
- (ii) $l_i \not\subset V_{\mathcal{A}}$.
- (iii) The l_i are pairwise disjoint.

Proof. The techniques are similar and rely on knowing that \mathcal{A} is ICM and Petersen. For (i), without loss of generality consider the case $i = 0$. The three hyperplanes $H^{0,0'}$, $H^{1,4'}$ and $H^{2,3}$ are distinct. (If, e.g., $H^{0,0'} = H^{2,3}$, then P_0 and P_2 meet in a line, contradicting that \mathcal{A} is Petersen.) So if l_0 is not a line, it is a plane. Suppose that were the case. Consider the three points $P_0 \cap P_1'$, $P_1' \cap P_2$ and $P_2 \cap P_0$; by Remark 3.1, all three must lie in P_1 . Since \mathcal{A} is ICM, they cannot coincide. (See case (iii) of Lemma 4.8.) They cannot be collinear since no two of P_1' , P_2 , and P_0 meet in a line. Thus, they span P_1 . Since each of them lies in two of the hyperplanes defining l_0 , they also span l_0 and we conclude $P_1 = l_0$. But this would give, e.g., $\dim P_1 \cap P_0' = 1$, again contradicting that \mathcal{A} is Petersen.

To prove (ii), suppose to the contrary, for example, that $l_0 \subset P_0$. (The remaining cases are similar.) Then the four points $P_0 \cap P_1'$, $P_0 \cap P_2$, $P_0 \cap P_4'$ and $P_0 \cap P_3$ lie on l_0 , and they must be distinct since \mathcal{A} is ICM. Now by Remark 3.1, the first two of these lie on P_1 and we deduce $l_0 = P_0 \cap P_1$. Working with the third and fourth points, we see that $l_0 = P_0 \cap P_4$ as well, contradicting that $\dim P_1 \cap P_4$ consists of a point.

Finally, to prove (iii), suppose, for example, that l_i and l_{i+1} share a point. Then P_i' , P_{i+1}' and P_{i-2} must intersect, which, as we saw in Section 4.2, contradicts the assumption that \mathcal{A} is an ICM Petersen arrangement. The remaining cases follow in a similar manner. \square

Finally, suppose that the arrangement \mathcal{A} satisfies:

Condition 6. For each perfect matching E_i of Γ_{Pete} , $\bigcap_{e \in E_i} H^e = \emptyset$.

Then for each $i \in \mathbb{Z}/(5)$,

$$p_i := \bigcap_{e \in E_i \setminus E_5} H^e,$$

is a point and we have the following lemma:

Lemma 5.2. *The union of $V_{\mathcal{A}}$, the five lines $\{l_i\}$ and the five points $\{p_i\}$ cut out by the five quintic hypersurfaces Q_0, \dots, Q_4 .*

Proof. Let $Q = Q_0 \cap \dots \cap Q_4$ denote the intersection of the five quintic hypersurfaces and let $I_Q = (q_0, \dots, q_4)$ denote the ideal generated by the five quintics. Then a point p of \mathbb{P}^4 lies in Q if and only if for each $i \in \mathbb{Z}/(5)$, there exists at least one $e \in E_i$ such that H^e contains p . A point p of Q lies in $V_{\mathcal{A}}$ if and only if for two distinct elements i and j of $\mathbb{Z}/(5)$, there exist $e \in E_i$ and $f \in E_j$ such that H^e and H^f contain p and intersect in one of the planes in \mathcal{A} . (The latter condition is equivalent to the condition that the edges e and f share a common vertex.)

Thus, a point p lies in the closure of the complement of $V_{\mathcal{A}}$ in Q (whose ideal is the ideal quotient $I_Q : I_{\mathcal{A}}$) if $p \in \bigcap_{e \in E} H^e$ for some matching E (not necessarily perfect) of Γ_{Pete} which is minimal among matchings E such that $E \cap E_i \neq \emptyset$ for all $i \in \mathbb{Z}/(5)$. Any such E is either $E_i \setminus E_5$, $\{ii', (i+1)'(i-1)', (i+2)'(i-2)'\}$, or E_5 . However, by Condition 6, the last subset cannot occur. Thus, the residual variety consists precisely of the union of the p_i and the l_i , which completes the proof. \square

Theorem 5.3. *The union of $V_{\mathcal{A}}$ and the five lines $\{l_i\}$ is scheme-theoretically cut out by the six quintic hypersurfaces Q_0, \dots, Q_5 .*

Proof. For the set-theoretic statement, it is enough, by Lemma 5.2, to show that if $i \in \mathbb{Z}/(5)$ and $p_i \notin V_{\mathcal{A}}$, then $p_i \notin H^e$ for all $e \in E_5$. Condition 6 implies that $H^{i,i'}$ does not contain p_i . On the other hand, if $j \neq i$, then either $(j)(j+1) \in E_i$ or $(j-1)(j) \in E_i$, so that $p_i \in H^{j,j'}$ would imply that $p_i \in P_j = H^{j,j'} \cap H^{j-1,j} = H^{j,j'} \cap H^{j,j+1}$, contradicting that $p_i \notin V_{\mathcal{A}}$. The scheme theoretic statement now follows because each Q_i is a product of linear forms. \square

Corollary 5.4. *There are exactly five 6-secant lines to $V_{\mathcal{A}}$.*

Proof. Let l be a 6-secant line to $V_{\mathcal{A}}$. Then any quintic hypersurface containing $V_{\mathcal{A}}$ contains l . It follows from Theorem 5.3 that $l = l_i$ for some $i \in \mathbb{Z}/(5)$. But for each $i \in \mathbb{Z}/(5)$, the line l_i meets the six planes $P_i, P_{i'}, P_{(i+1)'}, P_{(i-1)'}, P_{i+2}$ and P_{i-2} , but does not meet the lines $P_i \cap P_{i'}, P_{(i+1)'}, P_{(i-1)'}$ and $P_{i+2} \cap P_{i-2}$ and hence does not lie in the planes it meets. So l_i is a 6-secant to $V_{\mathcal{A}}$. \square

Remark 5.5. The general ICM Petersen arrangement satisfies Condition 6. One can verify, however, that the arrangement of Example 4.12 does not. In that case, the five quintics q_0, \dots, q_4 are linearly independent, but q_5 can be written as a linear combination of q_0, \dots, q_4 . We will discuss this example further in the last section.

We recall a well-known fact:

Lemma 5.6. *If four 2-planes in \mathbb{P}^4 meet pairwise in distinct points, then there exists a unique cubic hypersurface containing them.*

Now for each $i \in \mathbb{Z}/(5)$, let C_i be the unique cubic hypersurface in \mathbb{P}^4 containing the four planes $P_{i+1}, P_{(i+2)'}, P_{(i-2)'}, P_{i-1}$ (which correspond to the maximal independent vertex set V_i defined in Section 2) and set

$$S_i := H^{i,i'} \cup H^{(i+1)',(i-1)'} \cup H^{(i+2),(i-2)} \cup C_i.$$

Observe that $V_{\mathcal{A}} \cup l_i \subset S_i$.

Theorem 5.7. *$V_{\mathcal{A}}$ is scheme-theoretically cut out by the six quintic hypersurfaces Q_0, \dots, Q_5 and any two different sextic hypersurfaces S_j and S_k .*

Proof. Since the l_j are pairwise skew, it suffices to show that $V_{\mathcal{A}} \cup l_j$ is scheme-theoretically cut out by Q_0, \dots, Q_5 and S_j for every $j \in \mathbb{Z}/(5)$. Without loss of generality, we may assume that $j = 0$. Let $i \neq 0$. If l_i does not lie in S_0 , then, since l_i is a 6-secant to $V_{\mathcal{A}}$,

$$(l_i \cap S_0) \subset V_{\mathcal{A}}$$

and we are done by Theorem 5.3. Thus, our task is to prove that l_i does not lie in S_0 .

Here we will only show that l_1 does not lie in S_0 ; essentially the same proof applies to the remaining cases. Suppose, to the contrary, that $l_1 \subset S_0$. Then l_1 lies in C_0 or in one of the hyperplanes $H^{0,0'}$, $H^{1,4'}$ or $H^{2,3}$. Given that $l_1 = H^{1,1'} \cap H^{2',0'} \cap H^{3,4}$, we deduce that l_1 lies in either $P'_0 = H^{0,0'} \cap H^{0',2'}$, $P_{1'} = H^{1,1'} \cap H^{1',4'}$ or $P_3 = H^{2,3} \cap H^{3,4}$, contradicting Lemma 5.1.

Finally, suppose that $l_1 \subset C_0$. Recall that l_1 meets (but is not contained in) P_1 , P_4 and P'_2 . Since C_0 contains these three planes, C_0 must be the union of the three distinct hyperplanes spanned by l_1 and these planes. Thus the fourth plane, P'_3 that C_0 contains must lie in one of these hyperplanes and therefore meets either P_1 , P_4 , or P'_2 in a line, contradicting that \mathcal{A} is Petersen. \square

Remark 5.8. Let \mathcal{A} be a Petersen arrangement and let $I_{\mathcal{A}}$ be its vanishing ideal. One can show that $I_{\mathcal{A}}$ is *pl-generated*, i.e., that it is generated by products of linear forms. Recall that every quintic polynomial q_i we defined is a product of linear forms. Furthermore, five partitions of $E(\Gamma_{\text{Pete}})$ arise naturally from the 6-cycles of Γ_{Pete} . To each of these set partitions, we can attach a sextic polynomial which is a product of linear forms. One then shows that these quintics and sextics together generate the same ideal as the q_i and the forms vanishing on the S_j . Arrangements defined by products of linear forms have been studied by Björner, Peeva and Sidman [2].

6. SMOOTH GENERAL-TYPE SURFACES OF DEGREE 15

Once again, let $\mathcal{A} = \{P_i, P'_i\}_{i \in \mathbb{Z}_5}$ be a (labeled) ICM Petersen arrangement. We have shown that the vector space formed by quintic hypersurfaces containing $V_{\mathcal{A}}$ has dimension 6, so $V_{\mathcal{A}}$ can be linked in the complete intersection of two quintic hypersurfaces to a surface of degree 15. In this section, we will study the general surface arising in this way. First, however, we recall (from [8] or [7], for example) the main definition and some basic results of liaison theory.

Definition 6.1. Let X and X' be surfaces in \mathbb{P}^4 with no irreducible component in common. Then X and X' are said to be *linked* by a complete intersection of type (m, n) if there exist hypersurfaces Y and Y' of degrees m and n respectively such that $Y \cap Y' = X \cup X'$ scheme-theoretically.

In this situation, X is ICM if and only if X' is ICM, $X \cap X'$ is a curve, and there are two standard *exact sequences of linkage*,

$$0 \longrightarrow \omega_X \longrightarrow \mathcal{O}_{X \cup X'}(m+n-5) \longrightarrow \mathcal{O}_{X'}(m+n-5) \longrightarrow 0$$

and

$$0 \longrightarrow \omega_X \longrightarrow \mathcal{O}_X(m+n-5) \longrightarrow \mathcal{O}_{X \cap X'}(m+n-5) \longrightarrow 0.$$

The first sequence yields the relation between the Euler-Poincaré characteristics:

$$(6.1) \quad \chi(\mathcal{O}_{X'}) = \chi(\mathcal{O}_{X \cup X'}) - \chi(\omega_X(5-m-n)).$$

Letting d and d' denote the degrees of X and X' and π and π' their sectional genera, the corresponding sequence for linkage of curves in \mathbb{P}^3 yields the relation

$$(6.2) \quad \pi - \pi' = \frac{1}{2}(m + n - 4)(d - d').$$

Theorem 6.2. *Let $V_{\mathcal{A}}$ be a general ICM Petersen arrangement. Then the surface X linked to $V_{\mathcal{A}}$ by two general quintic hypersurfaces is smooth, non-minimal and of general type, with degree $d = 15$, sectional genus $\pi = 21$ and Euler-Poincaré characteristic $\chi = \chi(\mathcal{O}_X) = 5$. Moreover, X is embedded in \mathbb{P}^4 by the linear system*

$$\left| H_{\min} - \sum_{i=0}^4 l_i \right|,$$

where H_{\min} is the hyperplane class of the minimal model of X and the l_i are the lines of (5.1).

Proof. As we have shown in Section 5, $(I_{\mathcal{A}})_5$ is generated by six quintics. Thus, $V_{\mathcal{A}}$ is linked—via a complete intersection of two general elements of $(I_{\mathcal{A}})_5$ —to a surface X of degree 15. Furthermore, a computation with MACAULAY2 (see [1]) shows that X is smooth for a particular choice of \mathcal{A} and quintics in $I_{\mathcal{A}}$. Hence X is smooth in general.

By (6.2), X has sectional genus $\pi = 21$ and by (6.1), X has Euler-Poincaré characteristic $\chi = 5$.

Let H and K be a hyperplane section and a canonical divisor of X respectively. Recall the double point formula for surfaces in \mathbb{P}^4 [5]:

$$d^2 - 10d - 5H.K - 2K^2 + 12\chi = 0.$$

Using this formula, it follows that $K^2 = 5$, showing that X is of general type. Recall that the lines l_0, \dots, l_4 of (5.1) are 6-secants to $V_{\mathcal{A}}$. From the first standard linkage sequence, we obtain:

$$0 \longrightarrow \mathcal{J}_{X \cup V_{\mathcal{A}}}(5) \longrightarrow \mathcal{J}_{\mathcal{A}}(5) \longrightarrow \mathcal{O}_X(K) \longrightarrow 0.$$

Since $X \cup V_{\mathcal{A}}$ is a complete intersection of two quintic hypersurfaces and since $\mathcal{J}_{\mathcal{A}}(5)$ is globally generated by its sections outside $\bigcup l_i$, the canonical bundle ω_X is also globally generated outside $\bigcup l_i$. As a consequence, $\bigcup l_i$ is the fixed part of the canonical linear series, and l_0, \dots, l_4 are the exceptional lines of X . In particular, the embedding linear system of X has the desired form. \square

Remark 6.3. Let $\phi_{|H+K|}: X \rightarrow \mathbb{P}^N$ be the adjunction map, where $N = \pi + p_a - 1 = 21 + 4 - 1 = 24$. Then the minimal model X_{\min} of X is obtained as the image of X under $\phi_{|H+K|}$. Let H_{\min} be its hyperplane class and let K_{\min} be its canonical divisor. Then X_{\min} has degree $H_{\min}^2 = (H + K)^2 = 70$. Since $H_{\min}.K_{\min} = (H + K).K = 30$, the sectional genus of X_{\min} is 51. We note that the canonical model of X_{\min} is a singular surface of degree 10 in \mathbb{P}^3 .

Remark 6.4. For a general surface X obtained in this way, $H^0(\mathcal{J}_X(5)) = 2$. Therefore, since the family of ICM Petersen arrangements has dimension 30 and since the space of quintics containing each Petersen arrangement has dimension 6, the family of smooth general type surfaces of degree 15 obtained from the Petersen arrangements via linkage has dimension $38 = 30 + 2(6 - 2)$. On the other hand, a standard computation shows that $\chi(N_{X/\mathbb{P}^4}) = 35$. We conclude that $\dim H^1(N_{X/\mathbb{P}^4}) \geq 3$, but at present, we are unable to show that equality holds in general. Consequently,

we cannot rule out the possibility that the Hilbert scheme of \mathbb{P}^4 may be obstructed at the general surface X obtained via our construction.

Now let $P \in \mathcal{A}$. Then exactly three distinct planes of \mathcal{A} meet P in lines, and since \mathcal{A} is lCM, these three lines meet pairwise in three distinct points r_1 , r_2 , and r_3 of P .

Lemma 6.5. *There exists a quintic plane curve $Q \subset P$, not passing through r_1 , r_2 or r_3 , such that $X \cap P = Q \cup \{r_1, r_2, r_3\}$ scheme-theoretically.*

Proof. Set $\mathcal{B} = \mathcal{A} \setminus \{P\}$ and let $U = X \cup V_{\mathcal{B}}$. Then P and U are linked (by the same complete intersection of degree $(5, 5)$ that links $V_{\mathcal{A}}$ and X), so $C = P \cap U$ is a curve and the second standard exact sequence of linkage becomes

$$(6.3) \quad 0 \longrightarrow \omega_P \longrightarrow \mathcal{O}_P(5) \longrightarrow \mathcal{O}_C(5) \longrightarrow 0.$$

Consider the divisorial exact sequence:

$$(6.4) \quad 0 \longrightarrow \mathcal{O}_P(-C) \longrightarrow \mathcal{O}_P \longrightarrow \mathcal{O}_C \longrightarrow 0.$$

Let $L \subset P$ be a line so that $\omega_P = \mathcal{O}_P(-3L)$. By comparing (6.3) with (6.4) twisted by $\mathcal{O}_P(5)$, we conclude that $C \equiv 8L$ in P . Now there are exactly three planes of \mathcal{B} that intersect P in lines, so the intersection $X \cap P$ contains a curve $Q \equiv 8L - 3L = 5L$, i.e., a plane curve of degree 5.

Now let $K = K_X$ be a canonical divisor for X . Recall that the equivalence of Q for the intersection product $X \cdot P$ in \mathbb{P}^4 is given by a formula of Todd:

$$(X \cdot P)^Q = (K \cdot Q)_X + (K_P \cdot Q)_P - (K_{\mathbb{P}^4} \cdot Q)_{\mathbb{P}^4} - K_Q;$$

see [3, Example 9.1.7]. Since Q is a quintic plane curve, $K_P \cdot Q = -15$, $K_{\mathbb{P}^4} \cdot Q = -25$, and $\deg K_Q = 2 \cdot 6 - 2 = 10$. So the length of the residual scheme is

$$(6.5) \quad \begin{aligned} \deg(X \cdot P) - \deg(X \cdot P)^Q &= \deg X - (K \cdot Q) + 15 - 25 + 10 \\ &= 15 - K \cdot Q. \end{aligned}$$

(In the preceding discussion, $V \cdot W$ denotes intersection product while $V.W$ denotes intersection number.)

Consider the exact sequence of linkage:

$$0 \longrightarrow \omega_X \longrightarrow \mathcal{O}_X(5) \longrightarrow \mathcal{O}_D(5) \longrightarrow 0,$$

where $D = X \cap V_{\mathcal{A}}$. By comparing this sequence with

$$0 \longrightarrow \mathcal{O}_X(-D) \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_D \longrightarrow 0,$$

we deduce that $D \equiv 5H - K$ in X and therefore

$$K \cdot D = K \cdot (5H - K) = 120.$$

On the other hand, C is the sum of 10 quintics, one in each plane of \mathcal{A} , and we deduce that $K \cdot Q = 120/10 = 12$. So by (6.5), the residual scheme to Q in $P \cap X$ has length $15 - 12 = 3$.

Now $V_{\mathcal{B}}$ fails to be lCM precisely at r_1 , r_2 and r_3 and since $V_{\mathcal{B}}$ and $X \cup P$ are linked (by the same pair of quintics that link X to $V_{\mathcal{A}}$), $X \cup P$ fails to be lCM precisely at r_1 , r_2 , r_3 . It follows that the residual scheme to Q in $P \cap X$ is supported at the r_i , completing the proof. \square

Theorem 6.6. *The surface X admits a two-dimensional family of 5-secant lines, a one-dimensional family of 6-secant lines, and fifteen 7-secant lines.*

Proof. Keeping the notation, we see from Lemma 6.5 that any line in $P \in \mathcal{A}$ meets Q —and hence X —in five points (possibly with multiplicity). Furthermore, any line in P passing through r_1 , r_2 , or r_3 meets X in that point in addition to the five points where it meets Q . Finally, any of the three lines $\overline{r_i, r_j}$ meets X in those two points in addition to its five points of intersection with Q . Since there are three such lines in each of the ten planes P of \mathcal{A} and since each such line consists of the intersection of two of those planes, we arrive at the count of fifteen 7-secants. \square

7. A GENERAL-TYPE SURFACE WITH SYMMETRIES

Let \mathcal{A} be the plane arrangement constructed in Example 4.12. Recall that \mathcal{A} , and thus $V_{\mathcal{A}}$ has $D = \langle \rho, \sigma : \rho^5 = \sigma^2 = e, \rho\sigma = \sigma\rho^4 \rangle$ as a group of symmetries.

Consider the perfect matching $E_5 = \{ii'\}_{i \in \mathbb{Z}/(5)} \subset E(\Gamma_{\text{Pete}})$. Recall that each of the other perfect matchings are obtained as

$$E_j = \rho^j \tau E_5$$

for $j \in \mathbb{Z}/(5)$. In Section 5, we attached to each E_j a quintic polynomial $q_j \in I_{\mathcal{A}}$. It is easy to check that the first five quintic polynomials, q_0, \dots, q_4 are linearly independent. Let W be the vector space they span; it is clear that D acts on W .

We now denote by ζ a nontrivial 5th root of unity. Let w_1 and w_2 be eigenvectors for the action of ρ with eigenvalues ζ and ζ^2 respectively. Then, for each $i \in \{1, 2\}$, the quintics w_i and $\sigma(w_i)$ span a two-dimensional subspace W_i of W which is invariant under D . Take two general quintic polynomials in either W_1 or W_2 . Then the surface X linked, via these two quintics, to $V_{\mathcal{A}}$ is smooth. Thus, X is a smooth D -invariant general type surface in \mathbb{P}^4 .

As in Example 4.12, let V be the underlying vector space of \mathbb{P}^4 with basis $\{e_i\}_{i \in \mathbb{Z}/(5)}$. The action of D is given by $\rho(e_i) = e_{i+1}$ and $\sigma(e_i) = e_{5-i}$. In other words, V is the standard permutation representation of D . Recall that V contains the trivial representation of D , which is spanned by the vector $\sum_{i \in \mathbb{Z}/(5)} e_i$. Let p be the corresponding point of \mathbb{P}^4 . We can check that each of the quintic hypersurfaces used to link $V_{\mathcal{A}}$ to X contains this point. Thus p lies in the complete intersection $X \cup V_{\mathcal{A}}$. The point p cannot lie in any plane in \mathcal{A} since if p lies in $P \in \mathcal{A}$, then $\rho^i(P)$ would also contain p , which contradicts our assumption. Therefore $p \in X$. Thus, we have proved:

Theorem 7.1. *There exists a smooth surface $X \subset \mathbb{P}^4$ as in Theorem 6.2 and Theorem 6.6, admitting an action of the dihedral group on five elements which is free except at the single fixed point $p \in X$.*

REFERENCES

- [1] Hirotachi Abo, Holger P. Kley, and Chris Peterson, available at <http://www.uidaho.edu/~abo/programs.html>.
- [2] Anders Björner, Irena Peeva, and Jessica Sidman, *Subspace arrangements defined by products of linear forms*, J. London Math. Soc. **71** (2005), no. 2, 273–288.
- [3] William Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete 3. Folge, vol. 2, Springer-Verlag, 1984.
- [4] Daniel Grayson and Michael Stillman, *Macaulay 2, a software system for research in algebraic geometry*, available at <http://www.math.uiuc.edu/Macaulay2/>.
- [5] Robin Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, no. 52, Springer-Verlag, New York, NY, 1977.
- [6] Craig Huneke and Bernd Ulrich, *General hyperplane sections of algebraic varieties*, J. Alg. Geometry **2** (1993), 487–505.

- [7] Juan Migliore, *Introduction to liaison theory and deficiency modules*, Progress in Mathematics, no. 165, Birkhäuser, Boston, MA, 1998.
- [8] Christian Peskine and Lucien Szpiro, *Liaison des variétés algébriques I*, Invent. Math. **26** (1974), no. 4, 271–302.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IDAHO, MOSCOW, ID 83844

E-mail address: `abo@uidaho.edu`

URL: `http://www.webpages.uidaho.edu/~abo`

DEPARTMENT OF MATHEMATICS, COLORADO STATE UNIVERSITY, FORT COLLINS, CO 80523

E-mail address: `kley@math.colostate.edu`

URL: `http://www.math.colostate.edu/~kley`

DEPARTMENT OF MATHEMATICS, COLORADO STATE UNIVERSITY, FORT COLLINS, CO 80523

E-mail address: `peterson@math.colostate.edu`

URL: `http://www.math.colostate.edu/~peterson`