

Naturally Occurring Toxicants as Etiologic Agents of Foodborne Disease

Food Toxicology
 Instructor: Gregory Möller, Ph.D.
 University of Idaho

Learning Objectives

- Examine the etiology (causation) of human disease related to naturally-occurring foodborne toxicants.
- Understand the chemical complexity of foods.
- Explore goiter, tropical ataxia neuropathy (TAN), tropical amblyopia, lathyrism, and their linkage to foodborne toxicants.
- Review a range of natural food toxicants that are involved in human disease.

2

Complexity of Food

Nutrients

- Carbohydrates
- Proteins
- Lipids
- Minerals
- Vitamins

Non-nutrients

- Food Additives
- Naturally Occurring Secondary Chemicals
- Contaminants
- Processing Chemicals

3

Non-nutrient Chemicals in Different Foods

• Cheddar Cheese	160
• Orange juice	250
• Banana	325
• Tomato	350
• Wine	475
• Coffee	625
• Cooked beef	625

4

Natural Toxicants and Human Disease

- Goitrogens
- Cyanogenic glycosides
- Lathyrism
- Lectins
- Alkaloids
- Protease inhibitors
- Vasoactive amines

5

Goitrogens

- Contribute to growth of goiters
- Compounds in Cruciferae
 - *Brassica* species (cabbage, kale, turnips)
 - Seeds only - not leaves
- Combined w/ iodine deficiency




6

Food Toxicology

Cruciferae → Glucosinolates

- A class of about 100 naturally occurring thioglucosides that are characteristic of the Cruciferae and related families.
- Diets of people in many parts of the world include considerable amounts of Cruciferous crops and plants.
 - Processed radish and wasabi in the Far East
 - Cabbage and traditional root vegetables in Europe and North America.
 - Rapeseed, kale, swede and turnip may also contribute since they are extensively used as animal feed stuffs.
 - Glucosinolates in crops, such as oilseed rape (*Brassica napus*) and Brassica vegetables, is undesirable because of the toxicological effects of their breakdown products.
 - Breakdown products include nitriles, isothiocyanates, thiocyanates, epithionitriles and vinyl oxazolidinethiones.




Glucosinolates

7

Food Toxicology

Goitrogens

- Goitrogenic compounds (goitrin) are formed from breakdown of glucosinolates by thioglucosidase
- Goitrogenic metabolites
 - Nitriles; Thiocyanates; Oxazolidine



A thiocyanate


Goitrin
| 5-vinyl-2-oxazolidin-2-thione

8

Food Toxicology

Goitrogens

- Thyroid gland secretes thyroxine (TY), triiodothyronine (TII), thyroglobulin (TH)
- Controlled by hypothalamus and pituitary
- Hypothalamus produces thyrotropin-releasing hormone (TRH)
- Stimulates pituitary to release thyroid-stimulating hormone (TSH)



9

Food Toxicology

Goitrogens


- TSH promotes uptake of iodine, synthesis of TH and release of TY and TII which feed back to reduce TSH
- TY and TII hormones also affect
 - Oxygen consumption, cardiovascular function, neuromuscular activity, cholesterol metabolism, cerebral function
 - Growth and development

10

Food Toxicology

Goitrogens

- Goitrin and thiourea inhibit TY synthesis
- Thiocyanates, oxazolidine and nitriles inhibit uptake of iodine by thyroid
- Lack of iodine causes thyroid to enlarge and hypervascularate to trap more iodine = goiter




11

Food Toxicology

Goitrogens - Testing

- Weight and histology of thyroid
- Growth rate of patient/animal
- Iodine content of blood and thyroid
- Feed compound - measure uptake of radioactive iodine




12

Food Toxicology

Cyanogenic Glycosides

<p><u>Plant</u></p> <ul style="list-style-type: none"> • Bitter almond • Cassava root • Sorghum • Lima bean 	<p><u>Glycoside</u></p> <ul style="list-style-type: none"> • Amygdalin • Linamarin • Dhurrin • Linamarin
---	--




13 Cassava

Food Toxicology

Cassava Root

- The cassava (manioc, yucca; *Manihot esculenta*) is a woody shrub of the Euphorbiaceae (spurge family) that is extensively cultivated as an annual crop in tropical and subtropical regions for its edible starchy tuberous root, a major source of carbohydrate.
 - Cultivars contain varying amounts of CGs
 - Cultivar, drought, and food preparation significant for CGs




14

Food Toxicology

Cyanogenic Glycosides

- Toxic chemical = hydrogen cyanide
- HCN released when plant is chewed or chopped
- Releases 2 enzymes normally separate
 - Beta-glucuronidase
 - Hydroxynitrile lyase
- Act synergistically to release HCN




15 Cyanogenic Glycoside

Food Toxicology

Cyanogenic Glycosides

- Can reduce CN by chopping or grinding in water
- Cassava flour is made from boiled or fermented root
- Intestinal bacteria may also be able to break down cyanogenic glycosides to HCN



Food Toxicology

Cyanogenic Glycosides

Amygdaline (Laetrile)

- Anticancer compound
- Apricot pits
- Not approved in US
- Several deaths in foreign countries




17

Food Toxicology

Acute Cyanide Poisoning: Mechanism

- Shuts down cellular respiration - energy metabolism in mitochondria
- Binds ferric ion on cytochrome oxidase in Krebs cycle
- Min lethal dose 0.5-3.5 mg HCN/kg bw



18

Food Toxicology

Acute Cyanide Poisoning

Symptoms

- Muscular paralysis
- Mental confusion
- Respiratory distress
- Rapid onset

19

Food Toxicology

Acute Cyanide Poisoning

Treatment

- Nitrite or amyl nitrite
- Converts hemoglobin (Fe^{+2}) to methemoglobin (Fe^{+3})
 - Draws CN^- away from cytochrome oxidase
- Add thiosulfate to form thiocyanate

Ferrous Heme Ferric Heme

Food Toxicology

Chronic Cyanide Poisoning

- Occurs in areas of cassava in diet
- Not well understood
- Two disorders
 - Tropical Ataxia Neuropathy (TAN)
 - Tropical Amblyopia

Cassava
Manihot esculenta

21

Food Toxicology

Chronic Cyanide Poisoning

Tropical Ataxia Neuropathy (Konzo)

- Atrophy of optic nerve, ataxia, mental disorders
- More prevalent - West Africa
- High prevalence of goiter
- Low levels of S-containing aa
- Elevated plasma thiocyanate (goitrogen)

Konzo, a paralysis disease caused by poisoning from cyanide in cassava flour

22

Food Toxicology

Chronic Cyanide Poisoning

Dietary Modifiers

- No goiter with adequate iodine
- Malnutrition increases neural effects
- Protein-deficient diets
 - Lack S-containing amino acid to convert CN^- to thiocyanate

23

Food Toxicology

Chronic Cyanide Poisoning

Tropical Amblyopia

- Atrophy of optic nerve
 - Blurred vision, blindness
- Africa and South America where cassava is staple in diet
- Reproducible in lab animals

24

Lathyrism

- Consumption of peas – esp. *Lathyrus sativus*
 - Grass pea, blue sweet pea, chickling vetch, Indian pea, Indian vetch, white vetch, almorta or alverjón (Spain), guaya (Ethiopia), and khesari (India).
 - Primarily restricted to areas in Asia/Africa
- Well-known neurodegenerative disease
 - Enzyme inhibitor (BAPN)
 - Neurotoxic amino acid (ODAP)
- Hardy plant, drought resistant



25

Lathyrism

Two forms disease

- Osteolathyrism - animals
- Neurolathyrism - humans




26

Osteolathyrism

Animals

- Bone deformations
- Weakness in artery wall and connective tissues
- Beta-L-glutamylaminopropionitrile (BAPN)



27

Osteolathyrism

Mechanism

- BAPN inhibits lysyl oxidase enzyme
- Lysyl oxidase is needed to crosslink collagen strands for strength
- Collagen is main component of connective tissue and bones

28

Neurolathyrism

Humans


- Chronic consumption of *L. sativus*
- Paralysis of legs followed by general weakness and muscle rigidity
- Young man disease
- Sudden onset
 - Calf muscle spasms
- No animal model

29

Neurolathyrism

Humans

- *B-N-oxalyl-L-alpha,B*-diaminopropionic acid (ODAP)



30

Neurotoxicity


Humans

- Etiologic agent may be ODAP
- Only found with *L. sativus* species
- ODAP interferes with normal function of nerve synapse
 - Inhibits uptake of glutamic acid
- No animal model to study

31


Cholinesterase Inhibitors

- Found in a variety of plants - potato, tomato, eggplant
- Western African calabar bean (prototype)
 - Physostigmine
 - Natural carbamate



32

Cholinesterase Inhibitors



- Solanine - most studied - potato
- Glycoalkaloid - 20-100 mg/kg wet wt
- >200 mg/kg banned by FDA
- Lenape potato variety incident
 - > 300 mg/kg

33

Cholinesterase Inhibitors

Solanine

- Greatest concentrations
 - Peel, around sprouts
- Natural/artificial light increases levels
 - Russet Burbank - 250-700 mg/kg – 5 days
 - Green due to chlorophyll
 - Toxicity marker



34

Cholinesterase Inhibitors

Solanine

- Documented human toxicity and death
- Potato sprouts, sprouted potatoes, greened potatoes
- Gastric pain, nausea, vomiting, hyperesthesia, increased & accelerated respiration
 - 2/6 deaths
- Concentration unknown



Cholinesterase Inhibitors

Solanine - Case Study

- 420 mg/kg total alkaloid content
- Approx 50% solanine = 200 mg/kg
- Need to consume 1 kg potato for toxicity due to only solanine
- May act with other glycoalkaloids
 - Chaconine
- Animal LD₅₀ 500-1000 mg/kg

36

Vasoactive Amines


- Highest in cheese (aged), beer, wine
- Lower levels in banana, tomato, avocado, spinach, orange
- Spoiled meat (see food allergy)
- "Pressor Amines" (catecholamines)
 - Cause vasoconstriction, hypertension



37


Vasoactive Amines

- Meat/fish - bacterial action
 - Putrescine, cadaverine
- Banana/avocado
 - Dopamine, tyramine
- Catecholamine neurotransmitters
 - Norepinephrine, dopamine, serotonin



Vasoactive Amines: Problems

- Monoamine oxidase (MAO)
 - Widely distributed in body
 - Breaks down vasopressive amines
- MAO inhibitors - treat clinical depression
- Co-exposure to vasopressive amines in food
 - Mostly tyramine



39

Vasoactive Amines: Tyramine Mechanisms

- Indirect action
- Displaces normal catecholamines from granules in nerves
- Leads to hypertension
- Can be severe in presence of MAO inhibitors



40

Vasoactive Amines

Levels in Food - mg/kg

- Tyramine*****
 - Cheese 20-2000
 - Avocado 25
- Serotonin
 - Banana pulp 30
 - Avocado 10
- All others < 10, all sources

41

Vasoactive Amines

Symptoms

- Hypertension
 - Mild to severe
- Migraine headache
- Rare - intracranial bleeding/death

42

Pyrrolizidine Alkaloids

- A problem in food animal forage and human subpopulations exposed acutely in a toxic incident or chronically via cultural foods.
- High levels in some plants >5% dry wt
- 100 different compounds

43

Pyrrolizidine Alkaloids

- Most human exposure from herbal tea or crop contamination
 - Low level exposure from milk/meat
 - Bush tea in Jamaica
- Decreased use as herbal medicine
 - Comfrey - wound dressing (+other)
- Carcinogenic and liver toxin in animals
 - Epoxidation




Comfrey

44

PA Hepatotoxicity: Veno-Occlusive Disease

- Occlusive lesions are produced in the centrolobular hepatic veins (obliterating endophlebitis).
- Subendothelial edema, narrowing and occlusion of the lumina, atrophy, and necrosis of liver cells with portal hypertension are the results.



Veno-occlusive disease
Fibrous occlusion of acentrolobular venule

45

Tajikistan, 1992

- Due to a military blockade there was a late wheat harvest.
- Weeds were able to thrive in wheat fields.
- The harvest contained large amounts of the seed of *Heliotropium lasocarpium*.
 - The first cases occurred 6 weeks after contaminated bread was consumed.
- By the spring of 1993 there had already been more than 3900 cases.
- Patients at Stage I had abdominal pain, nausea, vomiting and asthenia.
- In Stage II hepatomegaly followed.
- Stage III was characterized by ascites.
- In the final stage there is hepatic encephalopathy.
- Case fatality ratio was 1.3% and increased with age.

46

Protease Inhibitors

- Anti-nutritional compounds
 - Legumes, grains, potatoes, eggplant, onion
- Inhibit gastric enzymes that break down proteins
 - e.g. trypsin (protease)
- All toxicity studies in animals
 - Human relevance unclear

47

Protease Inhibitors

- Pancreatic hypertrophy
 - Pancreatic hypersecretion of amino acid-rich proteins
 - aa deficiency
 - Growth retardation
- Similar mechanism of lathyrism

48

Other Possible Natural Toxicants

- Caffeine
- Spices
- Licorice, nutmeg, saffras
- Phytoalexins




49

Vicine and Convicine

- Fava bean alkaloids
- Causes favism in people who have an inherited absence of the enzyme glucose-6-phosphate dehydrogenase (G6PD) in their red blood cells
 - Headaches, dizziness, nausea, yawning, then
 - Vomiting, abdominal pain, and fever.
 - At this point, symptoms either spontaneously subside,
 - Or acute hemolytic anemia via oxidative stress occurs

50


Vicine and Convicine



51

Lectins

- Proteins or glycoproteins that bind carbohydrates
- Cause cells to agglutinate
- Basis for blood-type assays (hemagglutination)
- Decrease nutrient absorption from intestinal cells



52

Lectins: Sources

- Plant (800 species) and animal tissues
- Black beans, soybeans, lima beans, kidney beans, peas, lentils



53

Lectins: Problem

- Cause growth retardation from consumption of raw product
- Some very toxic (legume-specific)
 - Ricin - castor bean - rat LD50 0.05 mg/kg
 - Kidney bean - 0.5% rat diet two wks
- Growth retardation
 - 0.5-1% of diet black bean and soybean
- Heating destroys toxicity



54

Lectins: Mechanism of Action

- Unknown; complex
- Prevents nutrient uptake
- Gut flora involved
 - Germ-free animals show less effects
- Immune component?